

filters during the period between 1982 and 1984. The light curves show a stillstand near the phase,  $\phi = 0.8$  and the slight amplitude variations in the light curve which have been noted by Cherewick and Young(1975).

Using maximum times obtained from 70 photometric data and 20 spectroscopic data, we derived the period of light variation,  $p=0.20102977$  days and the rate of period change,  $\dot{p}=2.2\text{sec/century}$ . The (O-C) diagram shows a cyclic variation with a period of about 27 years and an amplitude of 17.9 minutes.

### **Fourier Analysis of T Tauri Light Variations**

Kwon, Suk Minn

*Department of Astronomy, Seoul National University*

In order to investigate the characteristics in the short-term light variations of T Tauri type stars, we have monitored the T Tauri on November 15, 1984 and January 11, 1985 for an hour each night. We have applied standard time-series analysis to the data. The power spectra of the T Tauri light variation increases toward low frequency in a power-law of  $P(f) \propto f^{-\gamma}$ . Preliminary interpretation of the power-law frequency spectrum suggests that the short-term light variations of the T Tauri may be originated from many hot clumpy regions of the stellar envelope.

### **Decomposition of Surface Brightness Profile of Barred Galaxies**

Ann, Hong Bae

*Department of Earth Science, Pusan National University*

We propose an objective method of decomposition of surface brightness profile of barred galaxies based on Kormendy's(1977) iterative method. This method can be applied to the barred galaxies with different degree of inclination. Some preliminary results of decomposition of several barred galaxies including nearly face-on galaxy NGC 4643 will be presented.

秋季學術大會

일시 : 1985年 11月 16日—17日

장소 : 전 북 대 학 교

韓國天文學會 創立 20週年 紀念

<招 請 講 演>

### **Star Count and the Galactic Galactic Structure**

Ishida, Keiichi

*Kiso Observatory, University of Tokyo*

It has been said that our Galaxy is composed of the two populations of the celestial objects, which

correspond to the different formation stages. The scenario was constructed by combining two kinds of observational data. One is the knowledge of kinematical characteristics, chemical abundances, and ages of various types of nearby stars. The other is the ones of the overall photometric structure of various types of external galaxies.

However, it has been known that all the nearby stars are not classified into two distinct populations but into some more of intermediate populations. In recent time, we have been accumulating photometric data for number of spiral and other types of galaxies, and they are decomposed into two components of disk and spheroid. However, again, some of the external galaxies are decomposed into three components including intermediate component which is called thick by some investigators.

The characteristics of the intermediate population will give us important key to know the formation process of our Galaxy. Star count data of UBV color bands for a large number of stars in a wide area of sky in high accuracy give a certain quantitative estimation of the intermediate population and metallicity gradient with the distance from the galactic plane.

The intermediate component thus identified has the local normalization factor 2% of the disk stars and scale height 1kpc. The stars of this component contribute 20% of the observed number of stars at  $15 \leq V \leq 17$  in the North Galactic Pole region. Metallicity gradient is found to be  $d[\text{Fe}/\text{H}]/dz = -0.5 \pm 0.1 \text{ kpc}^{-1}$  at  $z \leq 2$  kpc in logarithmic scale by assuming the solar abundance in the galactic plane.

## **Monochromatic Imagery of Galactic and Extragalactic Nebulae with the CFHT Camera**

Hua, Chon Trung

*Canada-France-Hawaii Telescope Corporation*

Hereafter are reported some results of morphological studies of galactic and extragalactic extended sources obtained with the image photon counting system (IPCS) at the Cassegrain focus of the 3.6m Canada-France-Hawaii Telescope (CFHT) on Mauna Kea, Hawaii. The observations were carried out using narrow-bandpass interference filters centered at the H alpha and N II 6583 taking into account the radial velocity of the objects in question. The processing of the data was made using the CDCA software facilities made available at the Laboratoire d'Astronomie Spatiale (LAS-Marseille) based on the same techniques previously described. Thanks to the high spatial resolution combined with the low-level light sensitivity of the IPCS, a faint nebulosity has been detected in H alpha emission surrounding the "double" galaxy I Zw18, spreading towards the two flares already reported by Zwicky. Other results concern the separation in several H II regions inside the nucleus of M101 (NGC 5457), monochromatic observations of NGC 3310 and of two small planetary nebulae, IC 351, and IC 2165. The absolute calibration of the device enables an assessment of fluxes emitted by different objects at the wavelengths under consideration.