

F325Identification of Osmo-inducible loci
in *Salmonella typhimurium*.안광수*, 방일수, 이영록, 박용근
고려대학교 이과대학 생물학과

Most microorganisms adapt and survive to osmotic stress conditions by inducing of specific adaptative system. The purpose of this study is to identify genes participating on osmo-inducible adaptive system in *Salmonella typhimurium*. By using MudJ(km, lac) operon fusion technique, we have identified two osmo-inducible lac operon fusions, *osib1011*, *osic1011*, in *Salmonella typhimurium*. Induction fold of these lac operon fusions to high osmotic condition was 3-5 folds(*osib1011*) and 15-20 folds(*osic1011*), respectively, and the β -galactosidase assay was performed in several osmotic conditions. In growth rate, B1011 decreased and C1011 was about the same compared with wild type. Both loci are located about 54-57 min. on the chromosome.

F326Construction of Mini-shuttle Vector pKU12 and
Restriction Enzyme Map이진규*, 황세현, 강형일, 박용근, 이영록
고려대학교 이과대학 생물학과

Mini-shuttle vector, pKU12 was constructed using binary vector pKB1 which was recombined with pBR322 and pKU10 isolated from *E. coli* and *Pseudomonas putida* KU816, respectively. The pKU12 having ampicillin resistance gene as an antibiotics marker has recognition sites for 7 restriction enzymes and its size is about 5.9 kbs. The restriction map of pKU12 was basically constructed by these restriction enzyme patterns. The pKU12 has ability to be transformed in *E. coli*, *P. putida*, and *Acinetobacter calcoaceticus* KU48. To investigate the usefulness of pKU12 in cloning of various genes concerned with degradation of recalcitrant aromatic compounds, *xyIE* gene was cloned by means of this mini-shuttle vector pKU12. It was confirmed that this vector was useful in cloning of various degrading genes for aromatic compounds.