

E331

HUMAN CYTOMEGALOVIRUS REPLICATION AND  $\text{Ca}^{2+}$   
RESPONSE IN CELL LINES OF NEURONAL ORIGIN

강경희, 이찬희  
충북대학교 자연과학대학 미생물학과

HCMV replication and  $\text{Ca}^{2+}$  response in human neuroblastoma (NB) cells were investigated. NB cells were permissive for HCMV multiplication with a delay of one day compared to virus multiplication in human embryonic lung (HEL) cells. The delay of HCMV multiplication appears to be correlated with a delay in the  $\text{Ca}^{2+}$  response. The cytoplasmic free calcium concentration ( $[\text{Ca}^{2+}]_i$ ) began to increase at 12 h p.i. in HCMV-infected NB cells, while  $[\text{Ca}^{2+}]_i$  increase in HCMV-infected HEL cells was observed as early as 3 h p.i. On the whole, the level of the increase in  $[\text{Ca}^{2+}]_i$  in NB cells was about 30% of that in HEL cells. These data suggest that the lower level of the increase in  $[\text{Ca}^{2+}]_i$  seems to result from the less efficient replication of HCMV in NB cells relative to that in HEL cells. When NB cells and HEL cells were co-cultured, the observed increase in  $[\text{Ca}^{2+}]_i$  was greater than expected. This synergistic effect did not require direct contact of the two types of cells, suggesting that factor(s) released from HCMV-infected cells is responsible for the synergistic effect on  $[\text{Ca}^{2+}]_i$ . Treatment with TPA of HCMV-infected NB cells resulted in a stimulatory effect on HCMV-induced  $[\text{Ca}^{2+}]_i$  increase. In conclusion, NB cells are permissive for HCMV replication and the delay in  $\text{Ca}^{2+}$  response may be a consequence of the lower responsiveness of NB cells to HCMV infection.

E332

Purification and Characterization of an Acetate specific  
Esterase from *Streptomyces coelicolor* A3(2)

오충훈\*, 김재현  
단국 대학교 자연 대학 미생물학과

*Streptomyces coelicolor* A3(2) had two isozymes of esterase when stained with  $\alpha$ -naphthyl acetate on non-denatured polyacrylamide gel electrophoresis. A comparison of the  $\alpha$ -naphthyl acetate specific esterase(ACE-EST) activity between *Streptomyces coelicolor* A3(2) and B6 strain containing hydrogen peroxide resistant gene revealed that the esterase activity in *Streptomyces coelicolor* A3(2) was lower than in B6 strain that was grown in YEME treated with 0.8 mM  $\text{H}_2\text{O}_2$ . An ACE-EST was purified 23-fold, with an overall yield of 12 % through ammonium sulfate, DEAE-sephadex, gel filtration and preparative polyacrylamide gel electrophoresis. The molecular weight of the enzyme was 32,000 dalton when determined by SDS-polyacrylamide gel electrophoresis and 96,000 dalton when determined by gel filtration. The esterase shown the maximal activity in Tris-HCl buffer at pH 8.0 and 35-40°C.