F813

A Screen for Genetic Loci Required for Body-wall Muscle Development During Embryogenesis in X-chromosome of *Caenorhabditis elegans*

Dug Kyu Lee*, Ji Yeon Shin* and Joo Hong Ahnn Department of Life Science, Kwangju Institute of Science & Technology, Kwangju, Korea

We have screened available chromosomal deficiencies for genetic loci whose zygotic expression is required for body-wall muscle cells during embryogenesis in Caenorhabditis elegans. In previous studies, it has been shown that no signs of muscle development were detected in nullo-X embryos during embryogenesis. Based on this observation, we are interested in identifying potential loci on X chromosome required for muscle development. Several chromosomal deficiency strains have been analyzed for their terminal phenotypes in order to map such myogenic loci on X chromosome. We have collected homozygotic embryos and observed the terminal phenotypes of the arrested embryos by Normarlski microscopy. These homozygotic embryos will be stained with monoclonal antibodies against muscle specific myosins in order to assess their muscle formation.

F814

Genomic in situ Hybridization of parental genomes in Lycoris flavescens M. Kim & S. Lee (Amaryllidaceae)

Lee, Byong-Soon* and Mu-Yeol Kim¹
Dept. of Biology, Jeonju University and Dept. of Biology,
Chonbuk National University

Genomic in situ hybridization(GISH) was used to discriminate the parental genomes of the Lycoris flavescens M. Kim & S. Lee (2n=19, 3M+4T+11A+1sm), a naturally occurring hybrid between L. sanguinea var. koreana (2n=22, 22A) and L. chinensis (2n=16, 6M+8T+2sm). For identifying parental genomes in interspecific hybrid, biotinylated genomic DNA of L. sanguinea var. koreana or L. chinensis hybridized preferentially in situ to mitotic chromosomes of L. flavescens as a probe conjugated to fluoresceinated avidin. The allopolyploid of L. flavescens was showed by the homology between 11A chromosomes of L. sanguinea var. koreana and 3M+1sm+4T chromosomes of L. chinensis.