

Lung mucus: A clinician's view

Won Dong Kim, M. D.

Respiratory Division, Asan Medical Center,
College of Medicine, University of Ulsan, Seoul, Korea

In ancient times, respiration's perceived purpose was only to cool the blood, and nasal mucus was thought to be a discharge from the brain. Later it was demonstrated that fluids cannot travel from the brain to the nose. It appears that credit for pointing out the medical significance of bronchial secretions and the biological value of the exocrine function of the lungs belongs to Laennec. He described the "chronic idiopathic pituitous catarrh" known today as bronchorrhea, which is characterized by paroxysms of expectoration. The importance of all this is that airway secretions, and their alterations, became one of the cardinal signs of many respiratory diseases [1].

Secretion of mucus

Respiratory mucus represents the products derived from secretion of the submucosal tracheobronchial glands and the epithelial goblet cells, whereas tracheobronchial secretions include the mucus plus other fluid and solute derived from the alveolar surface and the circulation. Sputum is a pathologic substance consisting of bronchial mucus contaminated by saliva, transudated serum proteins, and inflammatory and desquamated epithelial cells [2]. Sputum represents an increase in normal secretions and thus is pathologic.

Lucass and Douglas [3] postulated a double layer of mucus in the bronchial periciliary sol phase in which the cilia beat and a superficial blanket of gel which is moved forward by the tips of the cilia. The gel phase contains the bronchial glycoprotein, some serum proteins and protein bound to the bronchial glycoprotein; the sol phase contains any soluble components of bronchial secretion along with serum proteins. Mucus is a complex mixture of water, proteins and glycoproteins, lipids, and salts. Mucous glycoproteins or mucins are important constituents of airway mucus and are responsible for its viscoelastic properties. Respiratory mucus is usually cleared by airflow and ciliary interactions and sputum is generally cleared by cough. Airway clearance of mucus depends on the physical properties of the mucous gel, serous fluid properties, and ciliary function.

Respiratory mucus has three important roles in mucociliary clearance of particulate matter deposited within the respiratory tract; as an antibacterial substance (respiratory mucus contains immunoglobulins; lactoferrin, which chelates iron necessary for the growth of some bacteria; and lysozyme, which destroys bacteria); as a humidifier of inspired air and as a preventative of excessive fluid loss from the airway surface by dint of its hydrophilic nature [4].

Amounts of mucus secreted onto the airways surface in healthy lungs are small. Lavage of airways from normal volunteers recovers little mucin and other macromolecules. The total amount of surface liquid (mucus) in the conducting airways is

governed by the rate of mucous secretion, on the one hand, and the clearance of mucus by epithelial reabsorption, evaporation, ciliary transport, and cough transport.

Hypersecretion of mucus

Airway hypersecretion of mucus, if not accompanied by increased mucous clearance can lead to the accumulation of airway mucus. Increased volume of airway mucus has several functional consequences. Undesired effects include airflow obstruction, enhanced deposition of inhaled particulate matter in the tracheobronchial tree, and cough. However, the increased amount of airway mucus may also have a protective role by serving as a physical barrier against biologically active inhalants (mechanical screen), preventing bacteria colonizing the airways from adhering to the epithelium (biological screen), and inactivating cytotoxic products released from luminal leukocytes (chemical screen) [5].

Normal subjects may secrete substantial amounts of mucus only if they inhale noxious fumes (e.g., ammonia) or irritating substances, such as hypertonic saline, or acquire an acute respiratory tract infection. For laboratory animals as well as humans, chronic irritation or injury induces a state of mucous hypersecretion. Alterations in the quality or quantity of airway mucus, as encountered in many respiratory tract disorders, can result in impaired mucociliary clearance.

Clinical significance of hypersecretion of mucus

Accumulation of mucus in the airway tree may be caused by an increased volume of mucus produced, and perhaps also by decreased clearance due to defects in the ciliary clearance apparatus. Mucus hypersecretion may be chronic, but increased volumes are produced in exacerbations of chronic obstructive pulmonary disease, during attacks of asthma and in bronchiectasis and cystic fibrosis patients with chronic bacterial colonization.

Intraluminal mucus accumulation in the airways associated with mucous hypersecretion creates a clinical problem in almost all pulmonary and some nonpulmonary diseases. Airway mucous hypersecretion is an important determinant in the prognosis and clinical features of various pulmonary diseases such as chronic bronchitis, bronchiectasis and bronchial asthma, in addition to cystic fibrosis.

Hypersecretion of mucus contributes to the morbidity of airway diseases by predisposing patients to respiratory infections, contributing to airflow obstruction and patients' discomfort, significant association between chronic production of mucus and an increased risk of mortality [6].

Mucus hypersecretion and respiratory infections

Bacterial microorganisms are usually found as a component of sputum in chronic bronchitis. Their origin is either through inhalation or aspiration from the ambient air or oro-nasopharynx, or as part of the exudate from infected areas of the bronchi. The increase of secretions mixed with the products of inflammation would adversely affect mucociliary clearance. With the delayed clearance of bacteria, infection of the bronchial tissue occurs with establishment of a resident population of pathogens. Conversely, the

possibility have been addressed that bacteria and their products could contribute to mucous hypersecretion by directly stimulating mucus-secreting cells.

Airway hypersecretion and airway obstruction

Chronic bronchitis is a condition associated with chronic productive cough that lasts for at least 3 months of the year for 2 successive years. Thus, airway hypersecretion is a principal clinical feature in chronic bronchitis. The common association of chronic bronchitis and chronic limitation of airflow leads us to assume that hypersecretion of mucus results in limitation of airflow. Chest physiotherapy has been shown to facilitate sputum expectoration and, therefore, improvement in pulmonary functions [7,8]. It is also noteworthy that in the largest population-based studies, presence of chronic mucus hypersecretion is associated with excess FEV₁ decline [9].

Some epidemiological studies have suggested that chronic bronchitis and chronic obstructive impairment may be considered as separate entities, rather than as manifestations of a single disease [10,11]. Reid [12] had established that the major site of mucous gland hypertrophy was in the large bronchi in chronic bronchitis. On the other hand, Hogg et al [13] showed that the dominant site of irreversible airflow obstruction due to primary airway disease lay in the peripheral airways of less than about 3 mm diameter. Hence the major sites of obstruction and hypersecretion appeared to be in different sized airways, a finding which supported the epidemiological evidence for distinguishing the hypersecretory and the obstructive components of bronchial (and bronchiolar) disease.

However, goblet cells can be present in the peripheral airways (bronchioles), where they are not normally present, and it is referred to as goblet cell metaplasia. In chronic bronchitis with airflow limitation, the most specific bronchiolar change is goblet cell metaplasia, which can cause mucus plugging and may displace the surface layer of surfactant, allowing the airways to close more easily [14]. Thus increased mucus production from goblet cells in the more peripheral airways may contribute to obstruction in chronic obstructive pulmonary disease. Actually, for the measurements of intraluminal mucus, Aikawa et al [15] employed fixation of the lung by immersion instead of intrabronchial instillation of formalin, to prevent washout of mucus from the airways. They showed a large amount of mucus in the airways, especially in the peripheral airways, in chronic bronchitis with airflow limitation.

Clinically, it is often observed that patients with bronchial asthma expectorate a significant amount of sputum, especially during an asthmatic attack, and are subsequently released from dyspnea after the expectoration of a large amount of sputum. Moreover, a large amount of mucous plugging in airways with an increase in secretory cells is frequently observed in autopsied lungs of patients with bronchial asthma [16]. These suggest that mucous accumulation (mucous plugging) in airways resulting from mucous hypersecretion contribute to airway obstruction in addition to bronchoconstriction and mucosal edema in asthma.

Airway hypersecretion and mortality

Some epidemiological reports indicated that, after controlling for the degree of lung function, airway hypersecretion or phlegm was not significantly associated with death caused by chronic obstructive pulmonary disease, and they suggested that the

hypersecretory disorder is an innocent disease [10]. Nevertheless, other investigators observed a significant association between phlegm (or airway hypersecretion) and all causes of mortality [17].

Saetta et al [18] has reported that asthmatic patients who died suddenly and unexpectedly showed lumen occlusion by mucous plugs, plasma exudate, and inflammatory cells in the peripheral airways of autopsied lungs.

Turner-Warwick et al [19] found that half the idiopathic pulmonary fibrosis patients from their study expectorated a significant amount of sputum. Hiwatari et al [20] revealed that 44% of idiopathic pulmonary fibrosis patients who survived over 1 year had mucous hypersecretion and that idiopathic pulmonary fibrosis patients who suffer from mucous hypersecretion have a shorter survival rate than do those without mucous hypersecretion. These findings indicate that idiopathic pulmonary fibrosis is sometimes accompanied by a persistent mucous hypersecretion and bronchial gland hyperplasia.

Correlation between physical properties of sputum and clinical features

Airway clearance of mucus depends on the physical properties of the mucous gel, serous fluid properties, and ciliary function, as well as interactions between mucus and airflow or mucus and cilia. In cystic fibrosis sputum hydration is decreased and cough clearability is decreased, probably owing to alterations in the surface properties of the mucus resulting from impaired chloride transport into the periciliary fluid layer [21].

Several investigations have demonstrated that the degree of airway obstruction is inversely related to the physical properties of the sputum (i.e., the thicker the sputum, the greater the decrease of ventilatory capacity) [22,23,24].

Clinical assessment of sputum

Sputum is the abnormal material produced by, or expectorated from, the bronchopulmonary system. Production of sputum by a patient is indicative of an ongoing pathological process in the lung. Observation of chronic bronchitic sputum can reveal important clinical information concerning the type and level of the inflammatory process, the physical properties of the material, the extent of bronchial mucosal damage, and the identification of pathogenic microorganisms that may be present [25].

An useful general principle is that the volume of sputum expectorated usually reflects the severity of the chronic bronchitis. The volume should be obtained over a set period. A routine collection that includes a 24-h period is most valuable, since this accommodates the variability among bronchitics in terms of when each raises their sputum. The range of volumes expectorated in chronic bronchitis varies from 1 to 100 ml/day, with a rare individual producing more. The average range is 5-40 ml/day when patients are clinically stable, and may double or triple when acute infection is present. Bronchorrhea has been defined as a condition in which more than 100 ml of sputum is produced within 24 hr. It may be associated with tuberculosis, alveolar cell carcinoma, chronic bronchitis, or asthma: or it may be idiopathic.

The color and estimations of the specimen's degree of purulence can also provide some valuable information concerning the pathology in the bronchopulmonary system. The purulence of the specimen depends on the degree of yellow to green coloration. Macroscopic purulence is due to the content of polymorphonuclear leukocytes and deoxyribonucleoprotein fibers arising from disintegrated cells, particularly from

polymorphs. Cell counts being particularly useful on mucopurulent sputum from asthmatic patients since the pus may consist largely of eosinophils.

The usual implication of purulence in the literature is that this finding is synonymous with the presence of bronchopulmonary infection. Robertson [26] clearly related green coloration to the presence of myeloperoxidase released from degenerated polymorphonuclear neutrophil, an enzyme also prevalent in eosinophils. This coloration strongly suggests stasis in the airways of secretions containing polymorphonuclear leukocytes, which allows time for these cells to break down and release their peroxidases. The more peroxidases released, the greener the specimen.

Evaluation of the sputum cellular populations is of considerable value in differentiating chronic bronchitis from bronchial asthma, and in determining the etiology of acute exacerbations in chronic bronchitics. The polymorphonuclear neutrophils are the preponderant inflammatory cells in chronic bronchitis, and the frequency of eosinophils rarely exceed 5% during clinically stable periods. In bronchial asthma, polymorphonuclear neutrophils are also usually the preponderant polymorphonuclear cells, but the average percentage of eosinophils ranges from 20 to 40%, depending on corticosteroid dependency, but may be as high as 90%.

The cytological and Gram stain findings can distinguish the likely cause of the exacerbation in chronic bronchitis. The common bacterial species recovered from the sputum of chronic bronchitics during acute bacterial exacerbations are *Haemophilus influenzae*, *Moraxella catarrhalis*, *Streptococcus pneumoniae*, and *Haemophilus parainfluenzae*. Less commonly, *Pseudomonas aeruginosa* and *Klebsiella pneumoniae* can be found.

Management of mucus hypersecretion

The aim of therapy for bronchial hypersecretion has usually been to alter the rheological properties of bronchial mucus, to reduce the degree of airways obstruction, to enhance the function of the mucociliary escalator, and to promote expectoration. To reduce the amount of luminal mucus in the airways, three major strategies have been employed. The first is to inhibit mucous hypersecretion by pharmacological agents. The second is to stimulate ciliary activity and improve mucociliary interaction, thereby augmenting ciliary clearance; this is generally also accomplished with pharmacological agents. The third approach is to enhance cough clearance, either by changing the rheological properties of luminal mucus by pharmacological means or by moving mucus physically from smaller to larger airways from where it can be eliminated by cough clearance.

Dehydration of a patient with severe chronic bronchitis is associated with difficulty in expectorating small amounts of viscid sputum and rehydration usually eases expectoration. It is not clear whether this effect is on secretory gland activity or on tissue fluid transudate. Blanshard [27] showed a reduction in sputum viscosity following an increased fluid intake. These findings are in keeping with the clinical observation that the dehydrated patient in respiratory failure has difficulty in clearing retained secretions, which can be alleviated by intravenous fluids and presumably reflects the dependence of the secretory structures on adequate tissue fluid. Chodosh [28] also demonstrated that extra oral water increased the water content and decreased the apparent viscosity of sputum in chronic bronchitis, without increasing the volume expectorated. Therefore, adequate hydration is recommended as an essential measure, although the evidence for

efficacy of oral or parenteral water is limited.

Practically the management of mucus hypersecretion can be undertaken in two main ways: firstly improved clearance by physical methods, including cough, chest physiotherapy and the more recent method of high frequency oscillation, and secondly by pharmacological methods [29]. Pharmacological methods will be discussed later by other author.

Physical methods of clearance

Cough is the immediate reserve mechanism when mucociliary clearance is overwhelmed by copious secretions, as in chronic bronchitis. Encouragement should be given to cough from a high lung volume and perhaps augment cough with the forced expiration technique. Cough is significantly effective in removing secretions from central lung regions, in accordance with theory.

Chest physiotherapy consists of postural drainage where secretions are drained by the help of gravity, percussion with clapping of the chest wall producing a shock wave transmitted through the chest and thought to loosen bronchial mucus, and vibration with chest shaking, a further means of transmitting phasic energy, usually in expiration. Chest physiotherapy is significantly effective both in removing secretions from peripheral to more central regions and in promoting their final removal from the lungs.

References

1. Lenfant C. - Introduction. In: Takishima T, Shimura S, Eds. *Airway Secretion*. New York, Dekker. 1994.
2. Keal EE. - Physiological and pharmacological control of airway secretion. In: Brain JD, Proctor DF, Reid LM, Eds. *Respiratory defense mechanisms-Part I*. New York, Dekker. 1977; 357-401.
3. Lucas AM, Douglas LC. - Principles underlying ciliary activity in the respiratory tract, II. A comparison of nasal clearance in man, monkey and other animals. *Arch Otolaryngol*. 1934; 20:518-541.
4. Gallagher JT, Richardson PS. - Respiratory mucus: Structure, metabolism and control of secretion. In: Chantler EN, Elder JJB, Elstein M, Eds. *Mucus in health and Disease*. New York, Plenum Press. 1982; 335-350.
5. Wanner A. - Possible control of airway hypersecretion. In: Takishima T, Shimura S, Eds. *Airway secretion*. New York, Dekker. 1994; 629-646.
6. Larivee P, Rieves RD, Levine SJ, Shelhamer JH. - Airway inflammation and mucous hypersecretion. In: Takishima T, Shimura S, Eds. *Airway secretion*. New York, Dekker. 1994; 469-511.
7. Cochrance GM, Webber B, Clarke SW. - Effects of sputum on pulmonary function. *Br Med J*. 1977; 2:1181-1183.
8. May DB, Munt PW. - Physiologic effects of chest percussion and postural drainage in patients with stable chronic bronchitis. *Chest*. 1979; 1:1645-1648.
9. Sherman CB, Xu X, Speizer FE, Ferris BG Jr, Weiss ST, Dockery D.W. - Longitudinal lung function decline in subjects with respiratory symptoms. *Am Rev Respir Dis*. 1992; 146:855-859.
10. Fletcher C, Peto R. - The natural history of chronic airflow obstruction. *Br Med J*. 1977; 1:1645-1648.
11. Peto R, Speizer FE, Cochrane AL. - The relevance in adults of airway flow obstruction, but not of mucus hypersecretion, to mortality from chronic lung disease. *Am Rev Respir Dis*. 1983; 128:491-500.
12. Reid L. - Measurement of the bronchial mucous gland layer : A diagnostic yardsticks in chronic bronchitis. *Thorax*. 1960;15:132-141.
13. Hogg JC, Macklem PT, Thurlbeck WM. - Site and nature of airway obstruction in chronic obstructive lung disease. *N Engl J Med*. 1968; 278:1355-1360.
14. Macklem PT, Proctor DF, Hogg JC. - The stability of peripheral airways.

- Respir Physiol 1970; 8:191-203.
15. Aikawa T, Shimura S, Sasaki H, Takishima T, Yaehashi H, Takahashi T. - Morphometric analysis of intraluminal mucus in airways in chronic obstructive pulmonary disease. *Am Rev Respir Dis* 1989;140:477-482
 16. Messer JW, Peters GA, Bennett WA. - Cause of death and radiologic findings in 304 cases of bronchial asthma. *Dis Chest*. 1960; 38:616-624.
 17. Annesi I, Kauffmann F. - Is respiratory mucus hypersecretion really an innocent disorder? A 22-year mortality survey of 1,061 working men. *Am Rev Respir Dis*. 1986; 134:688-693.
 18. Saetta M, Stefano AD, Rosina C, Thiene G, Fabbri LM. - Quantitative structural analysis of peripheral airways and arteries in sudden fatal asthma. *Am Rev Respir Dis*. 1991, 143:138-143.
 19. Turner-Warwick M, Burrows B, Johnson A. Cryptogenic fibrosing alveolitis : Clinical features and their influence on survival. *Thorax*. 1980; 35:171-180.
 20. Hiwatari N, Shimuras, Sasaki T, Aikawa T, Asndo H, Ishihara H, Sekizawa K, Sasaki H, Takishima T. - Prognosis of idiopathic pulmonary fibrosis with mucus hypersecretion. *Am Rev Respir Dis*. 1991; 143:182-185.
 21. King M, Rubin BK. - Rheology of airway mucus, In: Takishima T, Shimura S, Eds, *Airway Secretion*, New York, Dekker. 1994;140:477-482
 22. Chodosh S, Medici TC, Ishikawa S, Enslein K,- Relationship of physical sputum characteristics and ventilatory capacity in chronic bronchial disease. *Chest*. 1973;63:56S-58S
 23. Keal E, Reid L, - Physico-chemical properties of sputum in reversible airways obstruction. *Chest* 1973; 63:58S-59S
 24. Lopez-Bidriero MT. - Individual and group correlations of sputum viscosity and airways obstruction. *Bull Physiopathol Respir* 1973;9:339-346
 25. Chodosh S. - Sputum production and chronic bronchitis. In: Takishima T, Shimura S, Eds. *Airway secretion*, New York, Dekker. 1994; 579-628.
 26. Robertson AJ. - Green sputum. *Lancet*. 1952; 1:12-15.
 27. Blanshard G. - The viscosity of sputum. *Arch Middlesex Hosp*. 1955; 5:222-241.
 28. Chodosh S. - Objective sputum changes associated with glyceryl guaiacolate in chornic bronchial disease. *Bull physiopathol Respir*. 1973; 9:452-456.
 29. Clarke SW. - Management of mucus hypersecretion. *Eur J Respir Dis*. 1987; 71(suppl.153):136-144.