

PSYCHO-PHYSICS

HUNG-KUK OH

School of Mechanical and Industrial Engineering
Ajou University Soowon Korea 442-749

ABSTRACT

Conventional atom model must be criticized on the following four points.

- (1) Natural motions between positive and negative entities are not circular motions but linear going and returning ones, for examples sexual motion, tidal motion, day and night etc.
- (2) Potential energy generation was neglected when electron changes its orbit from outer one to inner one. The $h\nu$ is the kinetic energy of the photo-electron. The total energy difference between orbits comprises kinetic and potential energies.
- (3) The structure of the space must be taken into consideration because the properties of the electron do not change during the transition from outer orbit to inner one even though it produces photon.
- (4) Total energy conservation law applies to the energy flow between mind and matter because we daily experiences a interconnection between mind and body.

New atomic model (the crystallizing π -bonding) was proposed in the journal of material processing technology since 1997 for the explanation of the mechanical behaviors in terms of physics. Π -ray physics proved that the electrons can come out from in the nucleus and modern chemistry corresponds to the π -bonding by the nuclear electrons. The π -bonding structure of the protons outside the nucleus is that electrons move between protons of the different atoms.

The perception step and the characteristic frequency in signal transduction is due to the accumulation of the π -rays outside of the proton before their absorption to the nucleus.

1. THE CRYSTALLIZING Π -BONDING

The crystallizing π -bonding [Ref.1, Ref.2] produces two π -rays of one wave length during the electron's going and receiving trip between two protons and makes an electron, a positron, a neutrino and an antineutrino disappear at the end of the trip as in Fig.1.

During the electron's going trip between two protons two π -rays of one wave length starts to produce.

During the electron's receiving trip the two π -rays of one wave length finishes to produce.

The two π -rays of one wave length are supplied and absorbed to the proton. The π^0 mesons produce implosion bonding between proton and neutron in this case.

There are four kinds of π -bondings between the protons, closed π -bonding, open π -bonding, covalent π -bonding and ferromagnetic π -bonding.

The closed π -bonding makes the protons fixed and only one electron moves between two protons of the attended ones. It produces two π -rays during the going and returning trip. Conventional metallic bonding belongs to this bonding.

The open π -bonding is an instantaneous bonding between two protons. The one of the two protons supplies an electron. It produces π -rays and then disbonds soon after. The signal transduction of the neuron belongs to this one.

The covalent π -bonding makes two electrons move between two protons and does not produce π -rays. It belongs to conventional covalent bonding.

The ferromagnetic π -bonding makes electrons circulate towards only one direction via the closed π -bonding protons and then produces magnetic moment. But it does not produce π -rays.

Conventional van der Waals bonding corresponds to the open π -bonding and the conventional hydrogen bonding also belongs to the open π -bonding. Conventional ionic bonding gives electrons to the other atom.

2. THE CRYSTALLIZING π -BONDING INSIDE THE NUCLEUS AND ENERGY CONSERVATION

The bonding structure of the protons inside the nucleus is also the crystallizing π -bonding as in Fig.2[Ref.3].

The electrons inside the nucleus were not possible in the conventional atomic model. π -ray physics said that they can exist in the nucleus satisfactorily [Ref.4], however.

In view of energy conservation law kinetic energies and potential energies of a positron, an electron, a neutrino and an antineutrino must supply the potential and kinetic energies of the produced π -gamma rays. They will be absorbed to the proton soon after the π -bonding and then

the strong interaction will happen due to the absorption.

3. THE CRYSTALLIZING π -BONDING OUTSIDE THE NUCLEUS AND ENERGY CONSERVATION

The bonding structure of the protons outside the nucleus is also the crystallizing π -bonding as in Fig.2[Ref.3]. The electrons move between protons of the different atoms and make π -rays of much lower energy. In view of energy conservation, following energy calculations are needed.

$$\begin{aligned} &\text{Kinetic energy of an electron after beta decay} \\ &=3 \text{ Mev}=3 \times 1.602 \times 10^{-19}(\text{J}) \times 10^6=4.8 \times 10^{-13}(\text{J}) \end{aligned} \quad (1)$$

$$\begin{aligned} &\text{Kinetic energy of a positron after beta decay} \\ &=3 \text{ Mev}=4.8 \times 10^{-13}(\text{J}) \end{aligned} \quad (2)$$

$$\begin{aligned} &\text{Potential energy of an electron after beta decay} \\ &=m_e C^2=9.109 \times 10^{-31}(\text{kg}) \times \{3.0 \times 10^8(\text{m/sec})\}^2=8.19 \times 10^{-14}(\text{J}) \end{aligned} \quad (3)$$

$$\begin{aligned} &\text{Energy of gamma ray} \\ &=h\nu=6.626 \times 10^{-34}(\text{J}\cdot\text{s}) \times 10^{21}(\text{1/sec}) \\ &=6.626 \times 10^{-13}(\text{J}) \end{aligned} \quad (4)$$

$$\begin{aligned} &\text{Energy of visible ray} \\ &=h\nu=6.626 \times 10^{-34}(\text{J}\cdot\text{s}) \times 10^{14}=6.626 \times 10^{-20} \end{aligned} \quad (5)$$

The number of its going and receiving trips between protons of the different atoms gets to about 10^7 to 10^8 before their disappearance of the elementary particles. The π -rays are accumulated outside of the proton and then will be absorbed into two π -gamma rays in the nucleus, which is followed by beta decay.

4. PERCEPTION STEP AND CHARACTERISTIC FREQUENCY IN SIGNAL TRANSDUCTION (COLOR, TASTE AND FLAVOUR) IN PSYCHOPHYSICS

The accumulation of the π -rays outside of the proton before their absorption to the nucleus makes the perception step appeared in psycho-physics.

The characteristic frequency in signal transduction is determined by the frequency and implosivity of the π -ray transduced through the perception neuron.

The color, taste and flavour are sensed from the difference of the frequency and implosivity of the transferred π -ray signal.

Conventional neuron signal can not give any specific information about the signal's origin.

The characteristic signal analysis from the π -ray physics will open new area in neuron science in the future.

Conventional physiology could not find out any cancer signal, which is produced from any abnormal trans-membrane.

If we have any technology, which can change the cancer signal into any other sound signal, the cancer will be conquered probably.

5. CONCLUSIONS

- (1) The crystallizing π -bonding produces two π -rays of one wave length during the electron's going and receiving trip between two protons.
- (2) The modern chemistry corresponds to the π -bonding by the nuclear electrons
- (3) The bonding structure of the protons outside the nucleus is that electrons move between protons of the different atoms.
- (4) The accumulation of the π -rays outside of the proton makes the perception step and the characteristic frequency in signal transduction in psycho-physics.

REFERENCES

1. HUNG-KUK OH, "Conventional Metallic Bonding and Three-Dimensional Crystallizing π -Bonding", Journal of Materials Processing Technology, 69(1997) page 134-147
2. HUNG-KUK OH, "Vortex of Electrons, π -Bonding of Atoms and Superconduction", Journal of Materials Processing Technology, 7411-3 (1998) page 126-130
3. HUNG-KUK OH, "Physics of Plasticity", Journal of Materials Processing Technology, 97 (2000) 19-29
4. HUNG-KUK OH, "NUCLEAR ELECTRONS", Proceedings of Society for Applied Micro Magnetic Energy, June 2000 page 11-19




Fig.1 Crystallizing π -bonding




Fig.2 Sequential arrangement of three-dimensional crystallizing π -bonding units of protons in the atoms of the periodic table