

Department of Pharmacy, College of Pharmacy, Catholic University of Taegu-Hyosung

*Rhododendron yedoense* Max. et Regel var. *poukhanense* Nakai (Ericaceae) is a deciduous and latifoliate shrub growing in Korea and Japan. The flower of this plant has been used as a depressant, but is very toxic. The roots of this plant have been reputed to be effective as hair-growing agents in the traditional medicine. The phytochemical research of this plant has never been reported. So, the roots of this plant were extracted with 95% MeOH. MeOH Ext. was subsequently fractionated into four parts: chloroform, ethylacetate, n-butanol and water fractions. In the present work, chromatographic separation of the chloroform fraction has yielded three triterpenoids corresponding to  $\alpha$ - or  $\beta$ -amyrins. Their structures were established by chemical and spectral evidences.

[PD2-2] [ 04/21/2000 (Fri) 14:50 - 15:50 / [1st Fl, Bldg 3] ]

### Asimitrin, a new bioactive Annonaceous acetogenin possessing an unusual non-adjacent bis-THF ring from *Asimina triloba* seeds

Woo MH<sup>o</sup>, Seo KM, Kim DH

Department of Pharmacy, College of Pharmacy, Catholic University of Taegu-Hyosung

The Annonaceous acetogenins are powerful inhibitors of mitochondria NADH: ubiquinone oxidoreductase and of the ubiquinone-linked NADH oxidase that is peculiar to the plasma membranes of tumor cells. The end result of both of these mechanisms is ATP deprivation. Thus, these compounds offer excellent potential for development as new antitumor, immunosuppressive, pesticidal, antiprotozoal, antifeedant, and antimicrobial agents. Exclusive in the tropical plant family, Annonaceae, the paw paw tree (*Asimina triloba* Dunal) is a temperate representative distributed abundantly in the eastern regions of North America. From the seeds, we have recently identified twenty novel and thirteen known bioactive Annonaceous acetogenins. Extended effort in these investigations has most recently led us to the isolation of a novel acetogenin, asimitrin. Asimitrin belongs to a new type of non-adjacent bis-THF ring acetogenin (one THF ring not possessing any flanking hydroxyls and the other bearing only a flanking hydroxyl) and 1,2-diol in long hydrocarbon chain.

[PD2-3] [ 04/21/2000 (Fri) 14:50 - 15:50 / [1st Fl, Bldg 3] ]

### Novel Triterpenoids from *Rhus javanica*

Lee IS, Jung KY, Park SH, Oh SR, Ahn KS, Hwang KH, Jung HJ and Lee HK

Immunomodulator Research Lab., Korea Research Institute of Bioscience and Biotechnology

*Rhus javanica* L. (Anacardiaceae) is a tall and broad leaf tree and distributed in Korea, Japan and China. Barks and leaves of this plant have been used in dysentery and diarrhea remedies in Korean herbal medicines. In a recent date, the production of tannic acid by cell cultures, the prophylactic efficacy against Herpes simple virus type and the antineoplastic effect were studied from this plant. However, reports of various chemical components were rare. Thus, in the course of studies on chemical constituents, we isolated three new dammarane triterpenes from the stem bark of this plant. Also, the known triterpene, semialatic acid, and the known steroid, stigmast-4-en-3-one were first isolated from this plant. Semialatic acid and stigmast-4-en-3-one had been reported previously from *Rhus semialata* and *Typha latifolia*, respectively. Based on the chemical and various NMR techniques (HMQC, 1H-1H COSY and HMBC), the structures of three new triterpenes were elucidated as semialactone, isofouquierone peroxide and fouquierone. We report here the structural assignments of three new triterpenes and the revised 13C NMR data of semialatic acid.