

The amyloid precursor protein (APP) can be processed via several alternative processing pathways. Alpha-secretase processing by cleavage within the amyloid beta-peptide domain of APP is highly regulated by several external and internal signals including G protein-coupled receptors, protein kinase C and phospholipase A2. Stimulation of m1 and m3 muscarinic acetylcholine receptors (mAChR), which are coupled to phosphoinositide hydrolysis and protein kinase C activation, has been shown to increase the release of soluble amyloid precursor protein (α APPs). There have been several reports indicating that Gq protein-coupled receptors including mAChRs (m1, m3), metabotropic glutamate receptors, and bradykinin receptors, regulate α APPs secretion. However, there are no direct evidence for the exact roles of G proteins. In the present study, to examine the regulation of Gq protein-linked muscarinic receptor-mediated α APPs release, we transiently transfected the different G α carboxyl-terminal peptide (G α q, G α i), which have shown a novel dominant-negative strategy (Gilchrist et al.), in SH-SY5Y cells expressing abundant m3 muscarinic receptors endogenously. In wide type cells, increase in α APPs released by normal metabolism of APP was detected in control medium in a time-dependent manner, and the α APPs release was stimulated by carbachol, a muscarinic agonist, and phorbol 12-myristate 13-acetate (PMA), a PKC activator. The carbachol-induced increase in α APPs release was blocked by EGTA, a Ca²⁺ chelator, indicating a Ca²⁺-dependent mechanism. On the other hand, PMA-induced α APPs releases was Ca²⁺-independent. Furthermore, to examine the regulation of α APPs secretion by upstream cellular signals, dominant-negative G α carboxy-terminal peptide-expressing SH-SY5Y cells were examined, and the results are discussed.

[PA1-42] [10/19/2000 (Thr) 10:00 – 11:00 / [Hall B]]

The role of intracellular Ca²⁺ increase and prooxidant production in the expression of ferritin light chain by sulfur amino acid deprivation in hepa1c1c7 cells

Kim HJ^{O*} and Kim SG

College of Pharmacy, Seoul National University (*A doctoral candidate from Duksung Women's University)

Sulfur amino acid deprivation (SAAD) induces oxidative stress through depletion of glutathione content. Ferritin synthesis increases in response to oxidative stress conferring resistance to subsequent insults. However, the molecular mechanisms for the expression of the ferritin gene by oxidative stress have not been studied yet. In the present study, change in intracellular calcium content was determined as part of the complete studies on the expression of ferritin light chain (FLC) gene by SAAD in hepa1c1c7 a murine hepatoma cell line. Confocal microscopy showed that intracellular calcium level was 1.5-fold increased after SAAD up to 80 sec, which extended for the next 200–300 sec, followed by returning to control level. The elevation of calcium by SAAD was prevented by GSH, methionine, cystine or cysteine, indicating that change in the redox-state might control the cellular calcium level. Furthermore, either verapamil or thapsigargin was active in inhibiting the increase in cellular calcium by SAAD, raising the notion that the calcium increase by SAAD might result from the influx of calcium via Ca²⁺ channel as well as the release from endoplasmic reticulum. SAAD increased the oxidation of dichlorofluorescein. Treatment of cells with verapamil or deferoxamine, or deficiency of extracellular calcium prevented prooxidant production by SAAD. Hence, elevation of intracellular calcium by SAAD was responsible for the oxidative stress. Northern blot analysis revealed that SAAD increased the mRNA level of FLC, which was inhibited by either EGTA or deferoxamine. Taken together, these results provided evidence that increases in intracellular calcium and oxidative stress by SAAD might lead to the enhanced expression of FLC mRNA in Hepa1c1c7 cells.

[PA1-43] [10/19/2000 (Thr) 10:00 – 11:00 / [Hall B]]

The changes of catecholamines and indolamines of rat brains by extremely low

frequency magnetic field exposure.

Jeong JH, Choi SH, Sung JH, Min YS, Kim JH, Huh IH, Sohn UD

Chungang Univ.

It has been shown that extremely low frequency time-varying magnetic field (MF) modulate the function of brain. We, therefore, were aimed at observing whether MF affects the central nerve system. We have studied the level of catecholamines and indolamines in rat brain using HPLC-ECD analysis system. The rats were exposed to sham or MF during 1, 2 and 3 days. After exposure, the parts of brain (cortex, hippocampus, striatum, cerebellum and thalamus) were isolated at the same time of day in order to escape the circadian rhythm of level in catecholamines and indolamines. The isolated brain samples were sonicated in 0.1 M perchloric acid and then centrifuged for the HPLC-ECD analysis to detect norepinephrine, DOPAC, dopamine, HIAA, HVA and serotonin. Exposure of rats to MF produced the increase of the level of norepinephrine, HVA and HIAA in striatum. In thalamus, norepinephrine also increased but dopamine decreased. These data suggests that exposure of extremely low frequency time-varying magnetic field to rats changes neurotransmitters such as norepinephrine or serotonin as well as their metabolites

[PA2-1] [10/19/2000 (Thr) 10:00 - 11:00 / [Hall B]]

Peroxynitrite scavenging and cytoprotective activity of 2,3,6-tribromo-4,5-dihydroxy benzyl methyl ether from marine alga *Symphyclocladia latiuscula*

Yoon JS^o, Choi HR, Park HJ, Choi JS, Choi WC and Chung HY

*Department of Pharmacy, ‡Department of Biology, Pusan National University, Pusan 609-35, and †Faculty of Food Science and Biotechnology, Pukyong National University, Pusan 608-737

Peroxynitrite (ONOO⁻), formed from the reaction of superoxide (·O₂⁻) and nitric oxide (·NO), is a cytotoxic species that can oxidize several cellular components such as proteins, lipids, and DNA. It has been implicated in diseases such as Alzheimer's disease, rheumatoid arthritis, cancer, and atherosclerosis. Due to the lack of endogenous enzymes responsible for ONOO⁻ inactivation, developing a specific ONOO⁻ scavenger is considerably important. The aim of this study was to evaluate the ability of marine natural products to scavenge ONOO⁻ and to protect cells against ONOO⁻. Methanolic extracts of 17 marine alga were tested for their ONOO⁻ scavenging activity. Among them, *Symphyclocladia latiuscula* showed the potent scavenging activity. CH₂CH₂ fraction of the methanol extract of *S. latiuscula* was highly effective for ONOO⁻ scavenging activity. Further analysis of the active fractionated extract identified 2,3,6-tribromo-4,5-dihydroxy benzyl methyl ether (TDB) as a potent ONOO⁻ scavenger. The data demonstrated that TDB led to decrease ONOO⁻-mediated nitration of tyrosine through electron donation. TDB showed the significant inhibition on nitration of bovine serum albumin (BSA) and low-density lipoprotein (LDL) by ONOO⁻ in a dose-dependent manner. They also provided cytoprotection from cell damage induced by ONOO⁻. TDB can be developed as an effective peroxynitrite scavenger for prevention of involved diseases.

[PA3-1] [10/19/2000 (Thr) 10:00 - 11:00 / [Hall B]]

Antiestrogenic effect of Conjugated linoleic acid on several estrogen-like compounds in MCF-7 human breast cancer cell

Shon MY^o, Park SJ, Oh SM, Kwon YI, Chung KH

College of pharmacy, Sungkyunkwan University