

induced edema a oral dose of 25mg/kg and 32.3±5.6% at PMA-induced edema at a topical dose of 0.5mg/ear, respectively. Cinnamic acid at a oral dose of 25 mg/kg has anti-inflammatory activity, but its potency has less than sinapinic acid : 18.7±2.2% at carrageenan-induced edema and 10.1±1.08% at PMA-induced edema, respectively. Sinapinic acid has the most active analgesic activity in HAc-induced writhing as compared with control (25.5±4.5%), but cinnamic acid has less effective than other cinnamic acid derivatives(19.2±3.7%). All cinnamic acid derivatives at a oral dose of 25mg/kg has inhibitory activity of inflammation and pain, but their activity have less than that of dexamethasone at a oral dose of 2mg/kg : 85.3±9.4% at carrageenan-induced edema and 65.8±7.2% at PMA-induced edema, and ibuprofen at a oral dose of 25mg/kg in the PMA-induced edema (67.0±4.6%) and HAc-induced writhing(45.1 ± 3.4%). These results show that anti-inflammatory activity cinnamic acid derivatives have the more hydroxyl and methoxyl of benzene ring, the more anti-inflammatory and analgesic activity.

[PB2-2] [10/20/2000 (Fri) 15:30 - 16:30 / [Hall B]]

Structure Activity Relationship of Cinnamic acid Derivatives on IgE -mediated Asthma in Non-anesthetized Guinea pigs

Park JH^o, Jang YW, Yoon JY, Cho JH, Kim CJ

Department of Pathophysiology, College of pharmacy, Chung-Ang University

The phenylpropanoids, C6-C3 compounds, have been reported that they have many biological activities, such as anti-inflammatory, anti-platelet aggregation, anti-viral activity and anti-oxidant activity. To investigate the anti-asthmatic activity of cinnamic acid derivatives in guinea pigs with IgE-mediated asthma, the specific airway resistance (sRaw) and tidal volume (TV) in immediate and late-phase asthmatic responses (IAR, LAR) induced by aerosolized-ovalbumin (OA) challenge in OA-sensitized guinea pigs were respectively determined by the double-chambered plethysmograph (HSE 850). And also recruitment of leukocytes, especially eosinophils into the lung, protein, histamine and phospholipase A2 (PLA2) activity in the bronchoalveolar lavage fluid (BALF) were analysed. Sinapinic acid at a oral dose of 25 mg/kg inhibited significantly the sRaw in IAR as compared with control (280±40% to 96±26%) and LAR (110±26% to 33±8%), respectively, and also suppressed not only the recruitment of the inflammatory cells, especially eosinophils, into the lung but also histamine release, protein exudation and PLA2 activity in BALF. Ferulic acid and caffeic acid at a oral dose of 25 mg/kg had significant anti-asthmatic activity, but their activity were less potent than that of sinapinic acid. The anti-asthmatic activity of cinnamic acid derivatives are same as cromolyn and dexamethasone. These results show that cinnamic acid derivatives have the more hydroxy and methoxy group of benzene ring, the more anti-asthmatic activity.

[PB2-3] [10/20/2000 (Fri) 15:30 - 16:30 / [Hall B]]

Cytotoxic Effect of Urushiol on Human Stomach Cancer Cells

Kim SH, Lee JH, Im EO, Jung NC¹, Na CS¹, Kim KW², Jung HY, Kim ND

Department of Pharmacy, ²Department of Molecular Biology, Pusan National University, Pusan, Korea, ¹Division of Special Purpose Tree, Forestry Research Institute, Suwon, Korea

Urushiol is mixture compound isolated from the sap of Korean lacquer tree (*Rhus vernicifera* Stokes). In the previous study, we evaluated the induction of apoptosis in MCF-7 human breast cancer cells. We observed that urushiol induced apoptosis through p53-dependent pathway. In this study, the ability of urushiol to induce gastric cancer cell death was investigated in the human gastric cancer cell lines, MKN-45 and MKN-28. The p53 statuses of the two cell lines are different. MKN-45 expresses heterozygous (wt/mt) but MKN-28 expresses mutant type of p53 (mt/mt). The cytotoxicity of urushiol measured by MTT assay. The IC 50 values were about 15 and

20 µg/ml on MKN-45 and MKN-28 cells, respectively. The expression levels of p53 and p21 proteins were measured by Western blotting after treatment of urushiol for 48 hours. The level of p53 proteins was not regulated significantly by urushiol on both cell lines. However, the p21 protein was slightly upregulated compared with the control groups. A cyclin-dependent kinase inhibitor p21 (WAF1/CIP1) protein, a key regulatory protein of the cell cycle, may have contributed to cell cycle arrest in urushiol treated stomach cancer cells. Thus, it is probable that urushiol mediated cell cycle arrest in MKN-45 and MKN-28 cells. However, further studies about several apoptosis-related proteins and DNA ladder formation are needed to know the mechanism of urushiol-mediated cell death in human gastric cancer cells.

[PB2-4] [10/20/2000 (Fri) 15:30 - 16:30 / [Hall B]]

Inhibitory effect of Ban-Myo(*Mylabris phalerata*) on tumor metastasis

Yoon TJ¹, Choi JS¹, Kang IC¹, Huh JE¹, Lee SJ¹, Lee EO¹, Lee SW,¹ Lee J², Kim SH⁰¹

Graduate School of East-West Medical Science, Kyung Hee University¹, R & D Center, DaeWoong Pharm².

We investigated the suppressive effect of solvent fractions from Ban-Myo(*Mylabris phalerata*) on tumor metastasis. Butanol(BuOH) fraction of Ban-Myo(BM) showed strong cytotoxic activity to B16-BL6 melanoma cell with IC50 value of 100µg/ml, while crude extract and ethylacetate(EtOAc) fraction of BM didn't even at concentration of 500µg/ml. Crude extract and butanol (BuOH) fraction induced apoptosis in B16-BL6 melanoma cells by FACS analysis. In experimental metastasis model developed by B16-BL6 melanoma cells, crude extract(500µg/mouse), ethylacetate(EtOAc) and butanol (BuOH) fractions significantly inhibited lung tumor colonization as compared with PBS-treated control group. Crude extract, EtOAc and BuOH fractions of BM also upregulated IL-1 β and TNF-alpha expression in murine peritoneal macrophages. Taken together these results, we suggest that crude extract, EtOAc and BuOH fractions of BM have apoptosis related cytotoxicity, anti-metastatic activity, activation of immune cells such as macrophages.

[PB2-5] [10/20/2000 (Fri) 15:30 - 16:30 / [Hall B]]

Study on Antitumor Effect of Kamicheungyeolhaedogtang(KCHT)

Kim K¹, Han KY², Kim DH¹, Choo JH², Ha ES², Kim CW², Kim SH²

Oriental Medical College, Taejon University¹, Graduated School of East-West Medical Science, Kyung Hee University²

To evaluate the antitumor activity and antimetastatic effects of Kamicheungyeolhaedogtang (KCHT), studies were done experimentally. KCHT is a oriental prescription for treatment of cancer. Its extract exhibited a significant cytotoxicity against A549, SK-MEL-2, SK-OV-3 and B16-BL6 cell lines and significantly inhibited DNA topoisomerase I. The T/C% was 145.8% in KCHT treated group in S-180 bearing ICR mice. It suppressed lung metastasis by B16BL/6 pathohistologically and CAM angiogenesis by 42% of control. These results suggest that KCHT extracts has antitumor and anti-metastatic effects.

[PB2-6] [10/20/2000 (Fri) 15:30 - 16:30 / [Hall B]]

Effects of aging and dietary restriction on the extracellular matrix proteins of rat tissues