

Hepatoprotective effects of flavonoid compounds is dependant on the antioxidant and detoxificant enzyme activity

Kim YG⁰, Kang MJ, Kim DH, Lee KT

College of Pharmacy, Kyung Hee University, Seoul 130-701, Korea

Abstract We used primary cultured rat hepatocytes injured by carbon tetrachloride as a model to screen for hepatoprotective effect of natural flavonoid compounds. To determine the protection effect of these compounds we measured the activity of GPT, LDH, and the levels of GSH and MDA.

CCl₄-induced liver toxicity on primary cultured rat hepatocytes was seen significantly elevation GPT, LDH, MDA and decrease GSH level.

Four flavonoid compounds showed anti-hepatotoxic effect by decrease GPT, LDH activity and MDA level and preserve GSH level. Moreover we measured radical scavenging effect, detoxifying enzyme (glutathione S-transferase, quinone reductase) and antioxidant enzyme (SOD, Catalase) activity.

Based on these results, we suggest that hepatoprotective activity of these four compounds preserve antioxidant status by scavenging reactive oxygen species and increasing detoxifying enzyme activity.

[PC1-20] [10/20/2000 (Fri) 15:30 - 16:30 / [Hall B]]

EDTA Attenuate Kalopanaxsaponin A-Induced Apoptosis in U937 Human Leukemia Cell.

Jang JW⁰¹, Park HJ², Kwon SH², Choi JW³, Lee KT¹

¹College of Pharmacy, Kyung Hee University, Seoul 130-701, ²Department of Botanical Resources, Sangji University, Wonju 220-702, ³College of Pharmacy, Kyung Sung University, Pusan 608-736, Korea

In previous study we screened that kalopanaxsaponin A induced apoptosis through inhibition of PTK, Bcl-2, c-myc, topoisomerase II- α and activation of Bax, PKC- α and caspase-3. This research was proposed to describe what was the crucial factor for apoptosis induced by kalopanaxsaponin A. Thereby we used NAC(N-acetyl-L-cysteine), vitamin E(α -tocopherol), vitamin C(ascorbic acid) and EDTA(ethylenediaminetetraacetic acid) to study the role of extra-cellular/intra-cellular ROS(reactive oxygen species) and metals. Only pre-treated EDTA blocked kalopanaxsaponin A-induced apoptosis by propidium iodide staining. Whereas antioxidant such as NAC, vitamin E and vitamin C did not show any effect. These result suggest that metals are associated with kalopanaxsaponin A-induced apoptosis through multi target signal transduction in U937 human leukemia cell.

[PC1-21] [10/20/2000 (Fri) 15:30 - 16:30 / [Hall B]]

Antitumor Activities of Manassatin A and B by Induction of Cell Differentiation and Apoptosis in Human Leukemia HL-60 cells

Seo BR, Jo HM, Choi JH, Lee KT, Ahn BT*, Jeong TS*, Bok SH*.

College of Pharmacy, Kyung Hee University, Cardiovascular disease substance research lab*, KRIBB

Differentiation and apoptosis inducing effects of manassatin A and B isolated from the *Saururus cernuus* were determined. 2.5µg/ml MNS-A and MNS-B showed cytostatic effect by Trypan Blue Exclusion Test. At over 5µg/ml these compounds were found to be a potent inducer of differentiation in human promyelocytic leukemia HL60 cells as assessed by the surface antigens of CD14 and CD66b, reducing nitroblue tetrazolium. Moreover, they showed apoptosis inducing effect through PI staining and DNA fragmentation at lighter concentration. These results suggest that MNS-A and MNS-B induce not only differentiation but also apoptosis in human leukemia cells.

[PC1-22] [10/20/2000 (Fri) 15:30 - 16:30 / [Hall B]]

Synthesis of Type IV Collagen and Laminin during Proliferation and Differentiation of 3T3-L1 Cells by Glycyrrhizin

Jeon H, Eun JS, Yum JY, Lim JP, Lim KH, Seo ES, Chae BS

College of Pharmacy, Woosuk University

We showed that the adipose conversion of 3T3-L1 cells was stimulated by Glycyrrhizin. To investigate effects of Glycyrrhizin on the basement membrane proteins during proliferation and differentiation of 3T3-L1 cells, we have analyzed synthetic amounts of basement membrane components such as type IV collagen and laminin. Glycyrrhizin stimulated the synthesis and secretion of type IV collagen from both 3T3-L1 preadipocytes and adipocytes. The synthesis and secretion of laminin was not affected by Glycyrrhizin. The continuous addition of Glycyrrhizin stimulated cell growth and increased cell density at confluence 1.2-fold. These results suggest an important role for type IV collagen in adipocyte differentiation

[PC1-23] [10/20/2000 (Fri) 15:30 - 16:30 / [Hall B]]

Effects of 19 naturally occurring prenylated flavonoids on arachidonic acid metabolizing enzymes: Cyclooxygenases and lipoxygenases

Chi YS, Jong HG, Son KH, Chang HW, Kang SS, Kim HP

College of Pharmacy, Kangwon National University, Department of Food and Nutrition, Andong National University, College of Pharmacy, Yeungnam University, Natural Products Research Institute, Seoul National University, Korea

Prenylated flavonoids are chemical entities having isoprenyl, geranyl, 1,1-dimethylallyl and/or lavendullyl moiety to their flavonoid backbone structure. In this study, effects of 19 naturally occurring prenylated flavonoids isolated from medicinal plants on cyclooxygenase (COX)-1, -2, 5-lipoxygenase (LO) and 12-LO were investigated using ¹⁴C-arachidonic acid as a substrate. The homogenates of bovine platelets, polymorphonuclear leukocytes and aspirin-pretreated lipopolysaccharide (LPS)-induced RAW 264.7 cells were used as COX-1, -2, 5- and 12-LO enzyme sources. Among 19 prenylated flavonoids, morusin, kuwanon C, sanggenon B, sanggenon D and kazinol B inhibited COX-2 activity (IC₅₀ = 73 - 100 µM), but the potencies were far less active than NS-398 (IC₅₀ = 2 µM). In contrast, many prenylated flavonoids such as kuraridin, kuwanon C and sophoraflavanone A inhibited COX-1 activity, among which kuraridin, kuraridinone and sophoraflavanone G having C-8 lavendullyl moiety showed potent and comparable activity (IC₅₀ = 0.1-1 µM) with indomethacin (IC₅₀ = 0.4-1.3 µM). Most prenylated flavonoids tested inhibited 5-LO activity, but only kuwanon C, papyriflavonol A and sophoraflavanone G showed the inhibitory activity against 12-LO at low concentration. Our results suggest that the position and nature of prenyl substitution greatly influence in vitro biological activities of these molecules.