

The flowers of *Prunus persica* Batsch have been used for skin disorders in East Asia from ancient time. In this investigation, the ethanol extract from this plant materials was prepared and evaluated for the protective effects on solar ultraviolet (UV)-induced skin damages including UVB-induced erythema in guinea-pigs, ear edema in ICR mice and skin carcinoma in SKH-1 mice. The *P. persica* extract clearly inhibited UVB-induced erythema formation dose-dependently when topically applied (IC₅₀ = 0.5 mg/cm²). And it also inhibited UVB-induced ear edema (49% inhibition at 3.0 mg/ear). In addition, the extract greatly reduced UVB-induced skin tumor formation (42% reduction of tumor incidence and 71% reduction of total number of tumors in 28 weeks) at the initial dose of 5.0 mg/cm²/day and the maintenance dose of 2.5 mg/cm²/day). From the extract, four kaempferol glycoside derivatives were successfully isolated and their contents were measured with HPLC. Among the derivatives, the content of multiflorin B was highest and this compound inhibited UVB-induced erythema formation, indicating that multiflorin B is one of the active principles of the extract. All these results suggest that the flowers of *P. persica* extract may be useful for protection against UV-induced skin damage when topically applied.

[PC1-29] [10/20/2000 (Fri) 15:30 - 16:30 / [Hall B]]

Cloning of nitric oxide synthase from *Staphylococcus aureus*

Park JH⁰, Yang SY¹, Lee HY¹, Han JH, Lee HW

College of Pharmacy, SungKyunkwan University, 1Departments of Pharmacology, College of
Medicine, Konyang University

Recently, we have reported the existence of bacterial nitric oxide synthase (NOS) in the prokaryotic cell, *Staphylococcus aureus* (*S. aureus*). The bacterial NOS in *S. aureus* requires the existence of cofactors (FMN, FAD, NADPH, BH₄ and calcium/calmodulin) and is induced by methanol. We compared the cofactor binding sites of murine iNOS with total genomic sequence of *S. aureus* and found seven matching sequences. Using homologous primers, we amplified 0.85kb DNA fragment from genomic DNA library of *S. aureus*. Screening of genomic DNA library produced twelve positive clones. Four DNA fragments (2, 4.5, and two 6 kb) were selected by Southern blot analysis and ligated into mammalian expression vector, pcDNA3.1.

[PC1-30] [10/20/2000 (Fri) 15:30 - 16:30 / [Hall B]]

Detection of Proteins in Sodium Dodecyl Sulfate -Polyacrylamide Gel Electrophoresis Using a Counter Ion Dye Staining Technique with Coomassie Brilliant Blue R / Neutral Red

Choi Jk⁰, Yoo GS, Lee JK

Chonnam National University, College of Pharmacy, Analytical Biochemistry Lab.

Coomassie Brilliant Blue R (CBBR) staining method has been used the most widely in the detection of proteins in Sodium Dodecyl Sulfate-Polyacrylamide Gel Electrophoresis (SDS-PAGE). However, it requires long staining time and can't be visualized in course of staining. Here we describe fast, sensitive and simple staining method that can be visualized in the course of staining employing a counter ion dye staining technique. Acidic dye, CBBR(0.005%) was mixed up with basic dye, Neutral Red (NR) (0.0005%). After electrophoretic separation in SDS-PAGE, gels were fixed with 40%MeOH/7%HAc for 30min. Then, gels were stained in 0.005%CBBR-0.0005%NR, in 30% MeOH/7%HAc staining solution for 30min-1hr. It was preferable to wash gels for 10 minutes after