

antithrombotic agents *in vitro* and *ex vivo*, and its antithrombotic effect *in vivo*. Chungpesagan-tang, Rhei Rhizoma and Scutellariae Radix potently inhibited ADP-, collagen-induced rat platelet aggregation in a dose-dependent manner *in vitro*. Chungpesagan-tang and most of its ingredients did not affect such coagulation parameters as APTT, PT and TT in human plasma. However, Rhei Rhizoma potently protected plasma clotting. Chungpesagan-tang treated group was significantly inhibited platelet aggregation in orally administered rats *ex vivo*. Puerariae Radix and Rhei Rhizoma also showed significant prolongation of tail bleeding time in conscious mice. Chungpesagan-tang showed significantly protection from death due to pulmonary thrombosis in mice.

[PC2-5] [ 10/20/2000 (Fri) 15:30 – 16:30 / [Hall B] ]

### Purification and Characterization of novel salt-inducible heparinase II from *Bacteroides stercoris* HJ-15

Hong SW<sup>0,1</sup>, Kim BT<sup>1</sup>, Kim DH<sup>1</sup>, Kim YS<sup>2</sup>

<sup>0,1</sup>College of Pharmacy, Kyung Hee University, <sup>2</sup>Natural Products Research Institute, Seoul National University

*Bacteroides stercoris* HJ-15, which is human colon gram-negative rod cell, has been known to degrade heparin, acharan sulfate and chondroitin sulfate. Recently we found to produce new five heparinases: one heparinase I, three heparinase II, one heparinase III. Among three heparinase II, one is a salt-inducible enzyme. In the present study, we tried to purify the enzyme. *Bacteroides stercoris* HJ-15 was cultured in 100L of tryptic soy broth containing hepain, collected, sonicated and centrifuged at 18,000rpm for 70min at 4°C. From the supernatant, the salt-inducible heparinase II was purified to homogeneity by QAE-cellulose, DEAE cellulose, CM Sephadex C-50, hydroxyapatite and phosphocellulose. The specific activity of the purified heparinase II was 81.3  $\mu\text{mole}/\text{mg}$ . Its molecular weight was 94KDa. The optimal activity was observed to be pH 7.3 containing 50mM KCl.  $\text{Cu}^{+2}$ ,  $\text{Ni}^{+2}$ ,  $\text{Co}^{+2}$ ,  $\text{Fe}^{+2}$  and  $\text{Zn}^{+2}$  inhibited the enzyme activity. These findings suggest that the biochemical properties of the purified enzyme should be different from the previously purified heparinase II.

[PC2-6] [ 10/20/2000 (Fri) 15:30 – 16:30 / [Hall B] ]

### Purification and Characterization of *Staphylococcus epidermidis* Urease

Min SH<sup>0</sup> and Lee MH

Microbiology Lab., College of Pharmacy, Catholic University of Taegu

The multi-subunit urease from *Staphylococcus epidermidis* ATCC12228 was purified 1,127-fold by ion exchange and gel filtration chromatography. The enzyme activity was highest when the cells were grown up to late exponential phase in LB supplemented with 50 $\mu\text{M}$  nickel chloride.

The specific activity of the purified urease was 993.8  $\mu\text{mol}$  urea degraded  $\text{min}^{-1}$ . The enzyme was shown to have a  $M_r$  of 409,000 and consist of three different subunits of 73.3, 22.7, and 13.8 kDa. The Michaelis constant ( $K_m$ ) was 8.9 mM urea. Activity gel-staining of the native urease with urea and cresol red revealed two major peptide bands.

[PC2-7] [ 10/20/2000 (Fri) 15:30 – 16:30 / [Hall B] ]

### Expression control of the *dnak* operon in *Streptococcus pneumoniae*