

carbohydrate of RTA significantly decreased uptake by liver and kidney, and resulted in prolonged circulating half-life. These results suggest that RTA prepared by carbohydrate-directed PEGylation would be more effective when constructed as immunotoxin for tumor targeting.

[PE1-31] [10/19/2000 (Thr) 15:00 – 16:00 / [Hall B]]

Comparative re-evaluation of protein denaturation for PLG microspheres prepared by W/O/W multiple emulsification processes.

Song SH^o, Shin TH, Kim JH, Cho SW, Choi YW

College of Pharmacy, Chung-Ang University

The purpose of this study was to re-evaluate the degree of protein denaturation during microencapsulation with poly (lactide-co-glycolide) (PLG) polymer. Ovalbumin(OVA), a model antigen, was entrapped in PLG microspheres by W/O/W multiple emulsification method in various conditions, i.e. with or without addition of polyvinylpyrrolidone(PVP) as a stabilizer for primary emulsification. PLG microspheres were suspended in PBS solution(pH 7.4) and incubated with shaking at 37.5°C. Release medium was collected periodically and analyzed using the micro-BCA protein assay method and ELISA method. In preliminary study with SDS-PAGE, typical bands for OVA showed that the primary structure of protein was not affected significantly. However, there was a decrease in immunoreactivity of OVA. In order to express the degree of protein denaturation, antigenicity ratio(AR) was introduced as follows: amount of immunoreactive of OVA / total amount of OVA released × 100(%). Especially the addition of primary emulsification stabilizer greatly influenced on protein denaturation : i.e. about 65% of AR with stabilizer versus less than 35% of AR without stabilizer. Therefore, in order to obtain information on structural integrity of protein, it is better to re-evaluate the protein denaturation employing immunoreactivity measurement for antigen entrapped in PLG microspheres.

[PE1-32] [10/19/2000 (Thr) 15:00 – 16:00 / [Hall B]]

Intranasal Delivery of Poly(ethylene glycol) Conjugated Salmon Calcitonin in Rats

Park MO^o, Lee SK, Kim BM, Yoon YS, Lee IB, Lee BH, Na DH, Lee KC

Drug Targeting Laboratory, College of Pharmacy, SungKyunKwan University

Activated PEG molecules (2000 and 5000 Da) were attached to salmon calcitonin via covalent linkage. Mono-PEG-sCT was then separated by size exclusion chromatography and characterized by CE and MALDI-TOF mass spectrometry. Six Male Sprague-Dawley rats (220–280 grams) per group were used. sCT (2 IU/kg) and PEG-sCT (2 IU/kg) were dissolved in saline solution and directly applied onto the nasal mucous membrane of rat by using micropipet (50 ul/rat) under anesthesia, respectively. Blood was withdrawn at a certain time intervals and centrifuged to collect the plasma samples. Serum calcium concentration was measured by o-cresolphthalein complexone complex assay using UV spectrophotometer. Serum calcium concentrations following nasal placebo administration remained around 100 % of the basal calcium levels during 6 hours. Nasal administration of native sCT (2 IU/kg) resulted in a slight decrease in serum calcium with a maximum decrease of less than 10 % at 30 min and returned to the control level within two hours. Significantly prolonged decrease of serum calcium were observed over six hours after nasal administration of PEG-sCTs. It was also observed that PEG5000-sCT had the less hypocalcemic effect than PEG2000-sCT. Therefore, these results indicate that PEG attachment to sCT can be investigated as a novel nasal delivery system of therapeutic peptides.

[PE1-33] [10/19/2000 (Thr) 15:00 – 16:00 / [Hall B]]

Effect of Enhancers on the in vitro Skin Permeation of Ciclopirox from novel Soft