## 5-3 Five New Species of Korean Amynthas (Oligochaeta, Megascolecidae)

\*Yong Hong and Samuel W. James<sup>1</sup>
Korean Institute for Biodiversity Research, Chonbuk National University

<sup>1</sup>Department of Life Sciences, Maharishi University, Fairfield, Iowa, USA

Most Korean earthworm species within the Megascolecidae belong to the genus *Amynthas*. This group is known to be diverse and abundant in litter layers and soils in forests.

The present study deals with 5 new species from Korea. Korean Amynthas vary in numbers of spermathecal pores; two pairs, three pairs, four pairs, five pairs, and none. Spermathecal pores of the species with 2 pairs are situated at the following segments: 5/6 and 6/7, 6/7 and 7/8, 7/8 and 8/9. The 5 new species aforementioned are 5/6-6/7 and presumed to be endemic to Korea.

## S-4 Origin of Hawaiian Drosophilids inferred from alcohol dehydrogenase gene sequences

Jecheol Park and Won Kim School of Biological Sciences, Seoul National University, Seoul 151-742, Korea

The evolution of Hawaiian drosophilids (including drosophiloid and scaptomyzoid species) has been thoroughly studied from various points of view, including morphology, behavior, cytology, and molecular phylogenetics. In spite of these studies, the origin of Hawaiian drosophilids is still controversial.

To clarify the origin of Hawaiian drosophilids, we determined Adh gene sequences for species of the genus Drosophila and related genera. The phylogenetic tree obtained from minimum evolution method with Juckes-Cantor distance showed that Hawaiian drosophilid lineage is more closely related to the virilis-repleta lineage than to the immigrans-Hirtodrosophila lineage. Within Hawaiian drosophilid lineage, Hawaiian Drosophila formed a monophyletic cluster separated from scaptomyzoid and quinaria section of subgenus Drosophila. In topology, our data shows that Hawaiian drosophilids may have the Oriental origin(s). However, we failed to find closely related convincing outgroup(s) for the Hawaiian specific clade and to resolve origin for Hawaiian and non-Hawaiian scaptomyzoid species.