

Z303 **Studies of Changes of Ca²⁺-channels Distribution In the Activated Mouse Ova**

Yeon-Soo Chang^{*}, In-Ha Bae

성신여자대학교 자연과학대학 생물학과

In muscle and neuronal cells, calcium channels have been classified by electrophysiological and pharmacological properties into (1) voltage-dependent Ca²⁺-channel ① P/Q-type Ca²⁺-channel ② N-type Ca²⁺-channel ③ L-type Ca²⁺-channel ④ T-type Ca²⁺-channel ⑤ R-type Ca²⁺-channel.

The immunocytochemical method was used to identify the existence of voltage-dependent Ca²⁺-channels in parthenogenetically activated 2-cell embryos by ethanol and SrCl₂ treatment. These 2-cell embryos were obtained by exposure to 6% ethanol for 6min and to 10mM SrCl₂ for 2h. P/Q-type Ca²⁺ channels and L-type Ca²⁺-channels have been identified. Whereas, three type of Ca²⁺-channel P/Q-type, N-type, L-type have been identified in 2-cell embryos fertilized in vivo. In the present study, activation by ethanol was faster than those by SrCl₂. However, there was difference in DAB staining of the embryos between ethanol and SrCl₂ treatment(87.7% and 54.1%). Intensity of staining was also different between ethanol- and SrCl₂-treated group. However, it has not been known why there was some difference in DAB staining and staining intensity in the present study.

Z304 **Developmental Expression Patterns of Mouse GM3 Synthase (mST3Gal V) during Embryogenesis**

Min-Young Ji *¹, Hee-Gon Choi¹, Dong-Hoon Kwak¹, Young-Ja Choi²,
Kyu-Yong Jung³, Su-Il Do⁴ and Young-Kug Choo¹

원광대학교 생명과학부¹, 원광보건대학 방사선과², 원광대학교 의과대학³,
대전생명공학연구소 분자 당연구실⁴

Gangliosides are ubiquitous membrane components in mammalian cells and are suggested to play important roles in various functions, such as cell-cell interaction, adhesion, cell differentiation, growth control and signaling. Among the gangliosides, GM3 has the simplest carbohydrate structure and all ganglio-series gangliosides are synthesized from GM3. GM3 is synthesized by CMP-NeuAc:lactosylceramide 2,3-sialyltransferase (GM3 synthase; ST3Gal V). In order to understand the developmental expression of ST3Gal V, We have investigated the spatial and temporal expression of mST3Gal V mRNA during the mouse embryogenesis [embryonic (E) days; E9, E11, E13, E15] by *in situ* hybridization. Specific signal was detected in neural tube (E9), neuronal differentiation, brain and liver development (E11, 13, 15). These results will provide further information for understanding the cellular control of recognition events mediated by carbohydrate groups in developmental embryogenesis.