

Direct Observation of CO to H₂ Conversion Factor in the Orion B Molecular Cloud: an Analysis of CO Absorption Line in the FUV Region

Dae-Hee Lee ¹, Kyoung-Wook Min ², Kwang-Sun Ryu ², Hyun-Soo Chung ³, W. Van Dyke Dixon ⁴, Mark Hurwitz ⁴

¹Department of Physics, Korea Advanced Institute of Science and Technology

²Satellite Technology Research Center, Korea Advanced Institute of Science and Technology

³Korea Astronomy Observatory

⁴Space Sciences Laboratory, University of California, Berkeley

We analyze the CO absorption line transition at 1076-1077Å from the spectrum of HD 37903, which is observed by Berkeley Extreme and Far-Ultraviolet Spectrometer (BEFS) on the ORFEUS telescope. HD 37903 is a bright UV source star behind the southern part of the Orion B molecular cloud and generating the reflection nebula NGC 2023. Concealed by the strong wings of H₂ absorption lines, it is hard to observe CO absorption lines in the FUV region as far. The great abundance of CO molecules in the Orion B molecular cloud makes it possible to obtain the column density of CO toward HD 37903 in this study ($N[\text{CO}] = 3.0 \times 10^{17} \text{ cm}^{-2}$). Based on the pre-derived H₂ column densities, we verify that the obtained CO column density is reliable according to the χ^2 - and F-test. In consequence of the direct measurement of CO column density, we obtain the CO to H₂ conversion factor $[\text{CO}]/[\text{H}_2]$ in the southern part of the Orion B molecular cloud, which is $[\text{CO}]/[\text{H}_2] = 4 \times 10^{-4}$. We obtain the three other H₂ tracers, which are the integrated CO and ¹³CO emission intensities, $W(\text{CO})$ and $W(^{13}\text{CO})$, and the ¹³CO LTE column density, $N^*(^{13}\text{CO})$, to investigate the physical conditions of the Orion B molecular cloud. The 14 m telescope in the Taeduk Radio Astronomy Observatory (TRAO) is used to observe the radio emission of CO(1-0) at 115.271 GHz and ¹³CO(1-0) at 110.201 GHz toward HD 37903.