

Molecular Biological Characterization of a New Crystal Protein Gene from *Bacillus thuringiensis* KSK-1182

Ming Shun Li, Jong Yul Roh, In Hee Lee, Jin Hee Chang,

Ho San Kim, Yeon Ho Je and Kyung Saeng Boo

Graduate School of Agricultural Biotechnology, Seoul National University,
Suwon, Korea

A new *Bacillus thuringiensis* strain (KSK-1182), having high toxicity to *Spodoptera exigua* larvae, was found to have a *cryIF*-like new gene besides *cryIAa*, *cryIAb*, *cryIAc*, *cryIE* and *cry2A* genes. Southern hybridization using a 2 kb fragment containing the *cryIF*-like PCR product was performed and a *HindIII*-*XhoI* 8 kb fragment was cloned. Through subcloning and construction of *ExoIII*-deleted mutants, a 5.3 kb fragment containing the full gene (named as the *cryX*) was sequenced. The *cryX* full gene was composed of 3,513 bp encoding 1,171 amino acid-long sequence. The δ -endotoxin of *cryX* contained all 8 blocks which played an important role in insecticidal mechanism. And its expression-regulating region of promoter, ribosome binding site, and terminator also existed but it had some different sequence from those of known *cry* genes. Through comparison of nucleotide and amino acid sequences between *cryX* and known *cry* genes, *cryX* showed only 77.6% and 73% homology to those of *cryIHa1*, the most closely related, and it was concluded that it is a new δ -endotoxin gene. At present, we are trying to express the *cryX* full gene in *B. thuringiensis* Cry-B strain for characterization of the CryX protein.