

inhibitory activity was examined in vivo, GRR, GRA, GRP inhibited the increase of blood glucose level in an oral administration of glucose in KKAY mice. In the mechanism study, GRR and GRP upregulated PPAR $\gamma$  protein expression, GRR and GRA upregulated GLUT4 protein expression when compared to the control group. sGLT1 mRNA levels were not different among the study groups. In conclusion, Ginseng Radices can ameliorate the hyperglycemia in KKAY mice possibly through up-regulations of PPAR  $\gamma$  protein in epididymal fat and glucose transporter protein in quadriceps muscle.

[PA1-34] [ 04/20/2001 (Fri) 10:30 – 11:30 / Hall 4 ]

### Antiangiogenic Effect of Gardenia Fruits

Joo MH, Kim SH, Kim MH, Song YS, Park EH,

College of Pharmacy, Sookmyung Women's University

Angiogenesis is known to be a complex process involving endothelial cell migration, proliferation, invasion and tube formation. Inhibition of these processes may have implications in various angiogenesis-mediated disorders. It is very desirable to find natural products containing anti-angiogenic activity. To investigate the effect of gardenia fruits, their ethanolic extract was prepared and dried in vacuo. It was used to test its effect in the chick chorioallantoic membrane(CAM) assay. The extract was found to contain the significant anti-angiogenic activity. Its anti-angiogenic activity appeared to be dependent on concentration. These findings suggest another usefulness of gardenia fruits.

[PA1-35] [ 04/20/2001 (Fri) 10:30 – 11:30 / Hall 4 ]

### Antidiabetic effect and mechanisms of DKY in KKAY mice

Kim YY<sup>o</sup>, Kim YL, Han KS, \*Park HS, \*Lee JS, and Chung SH

College of Pharmacy, Kyunghee University, Seoul 130-701, Korea, \*Research & Development center, Samchundang Pharm. co., Seoul 150-037, Korea

Dongryongkangdangyoungjung (DKY), composed of 36 herbs including *Cordyceps Sinensis*, has been used as an antidiabetic prescription in China. We have studied the antidiabetic effect and mechanism of DKY in male hyperglycemic and hyperinsulinemic KKAY mice. Oral administration of DKY at 500, 1000 mg/kg/day lowered plasma glucose from a week after treatment and the hypoglycemic activity was retained for 4 weeks, also Insulin and HbA1c level were markedly reduced dose dependently. But there was no significant difference at 2000 mg/kg/day level as compared to the control. In the mechanism study, PPAR $\gamma$  mRNA and protein in epididymal fat were increased in DKY-treated group, which is similar to those of the group treated with rosiglitazone, agonist for PPAR $\gamma$ . We have tried to investigate GLUT4 mRNA and protein expressions in quadriceps muscle, PEPCK activity and mRNA expression in liver. We found no correlation between mRNA and protein expression. These results suggest that DKY-induced activation of PPAR $\gamma$  directly correlate with antidiabetic action.

[PA1-36] [ 04/20/2001 (Fri) 10:30 – 11:30 / Hall 4 ]

### Antihypertensive and antihyperglycemic effects of Ginseng Radix Palva and Ginseng Radix Alba in SHR with diabetes

Han KS<sup>o</sup>, Kim YL, Kim JM, Youn SH, \*Park SH and Chung SH

Kyunghee University, School of Pharmacy, Seoul 130-701, Korea, \* Ilwha central research institute

The aim of this study was to investigate a potential antihypertensive and antihyperglycemic effects of Ginseng Radix palva (GRP) and Ginseng Radix Alba (GRA) in spontaneously hypertensive rat (SHR) injected with streptozotocin (STZ). Animals were divided into four groups containing 6 rats each as follows : Group 1, SHR drinking a tap water (hypertensive control) ; Group 2, SHR-STZ (hypertensive and hyperglycemic control) ; Group 3, SHR-STZ administered with 500 mg/kg/day of GRA ; Group 4, SHR-STZ administered with 500 mg/kg/day of GRP. Ginseng Radices were administered to 6 week-old SHR-STZ rats for 4 weeks and blood pressure (SBP, DBP), heart rate and blood glucose were determined at 2 weekly. After 4 weeks, serum creatinine, urinary albumin excretion(UAE) and glomerular TGF-beta mRNA and protein levels were compared between experimental groups. Ginseng Radices treated groups improved hypertensive and hyperglycemic abnormalities when compared to the SHR-STZ control.

[PA1-37] [ 04/20/2001 (Fri) 10:30 - 11:30 / Hall 4 ]

### **G protein and MAP kinase mediates sphingosylphosphorylcholine-induced feline ileal smooth muscle cell contraction.**

Lee TS<sup>o</sup>, Shin CY, Lee YP, Park SY, Kim TH, Kang HY, Huh IH and Sohn UD

Department of Pharmacology, College of Pharmacy, Chung Ang University

It has been shown that sphingosylphosphorylcholine (SPC) produced pertussis toxin sensitive contraction of feline ileal smooth muscle cells (Lee et al., 1999). We studied further mechanism of the contraction that might involve the activation of a certain Gi protein and phospholipase and kinases-dependent pathway using Western blot and GTP $\gamma$ S binding techniques. Western blotting using G protein antibody revealed that subtypes of Gi1, Gi3 and Go existed in feline ileum. Gi3 antibody treatment in permeabilized single cells isolated with collagenase decreased SPC-induced contraction. In addition, [<sup>35</sup>S]GTP $\gamma$ S binding study showed that Gi3 subtype activated significantly after SPC treatment, because SPC elevated the binding of Gi3 protein with [<sup>35</sup>S]GTP $\gamma$ S. To examine the relation of MAP kinase and SPC-induced contraction, we pretreated PD 98059, MEK inhibitor, and SB202190, p38 MAP kinase inhibitor. The contraction was blocked by PD 98059, but not by SB 202190. PKC inhibitor, chelerythrine, and PLC inhibitor, neomycin also decreased the contraction. However, co-treatment of PD98059 and chelerythrine showed no significant difference. When we examined MAP kinase activation using phospho MAP kinase antibody, SPC increased the band of phospho-MAP kinase. We suggest that SPC-induced contraction is mediated via pathway involving Gi3 protein coupled to PLC, resulting in the activation of PKC and p42/44 MAP kinase in feline ileal smooth muscle cells.

[PA1-38] [ 04/20/2001 (Fri) 10:30 - 11:30 / Hall 4 ]

### **The effect of extremely low frequency magnetic fields on NO levels in brain regions of rats**

Jeong JH<sup>o</sup>, Sung JH, Kum C, Kim JS, Kim HR, JH Kim, HJ Song, Huh IH, Sohn UD

Department of Pharmacology, College of Pharmacy, Chung Ang University

It has been shown that the extremely low frequency magnetic fields (MFs) produce the biological effects in the brains. MFs affect nitric oxide (NO) production, which plays a major role in neurotransmission and pathogenesis in the brain. To confirm the relevance of NO and MFs in the brain, we investigated the concentration of NO in various brain areas (cortex, cerebellum, hippocampus thalamus and striatum) of rats following exposure to MFs. Rats were exposed for 2 or 5 days to MFs (60Hz, 10, 15, 20 gauss, G) or sham in a laboratory exposure facility. 20 G MFs exposure for 2 or 5 days caused a continuous increase in NO release of hippocampus, thalamus and striatum, whereas no changes were obtained in cerebellum and cortex. Low strength in MFs did not alter the