

MRRC and College of Pharmacy, Wonkwang University, Iksan 570-749, Korea, \*Professional Graduate School of Oriental Medicine, Wonkwang University, Iksan 570-749, Korea

Bioassay-guided fractionation of the various solvent extracts of *Rhodiola sacra* furnished two hepatoprotective and two inactive compounds (compounds 1-4) *in vitro*. The structures of 1-4 are identified as cinnamyl alcohol, kaempferol, daucosterol and salidroside, respectively, by comparison of spectral data with those of literature. Compounds 1 and 2 showed better hepatoprotective effects against tacrine-induced cytotoxicity in Hep G2 cells than silymarin, as a positive control (EC<sub>50</sub>, 38.6 µg/ml). The EC<sub>50</sub> values of 1 and 2 are 29.9 and 9.62 µg/ml, respectively.

[PD2-33] [ 04/20/2001 (Fri) 13:30 - 14:30 / Hall 4 ]

### Antimutagenic activity and cytotoxicity of *Rumex acetosa* L.

Lee NJ<sup>01</sup>, Lee KH<sup>1</sup>, Lee DU<sup>2</sup>, Park SS<sup>2</sup>, Ryu SY<sup>3</sup>

<sup>1</sup>College of Pharmacy, Pusan National University, <sup>2</sup>Department of Biochemistry, Dongguk University, <sup>3</sup>Korea Research Institute of Chemical Technology

Antimutagenic compounds act by either inactivating mutagens or interfering in the process of mutagenesis, which lead to their antimutagenicity by reducing the frequency or rate of spontaneous or induced mutation. Antimutagenic agents may prevent cancer because they can either destroy mutagens in or out of body cells or block mutagens which damage DNA and cause mutations in cells. As a part of our continuing search for anticancer agents from natural sources, we have investigated the antimutagenic activities of the total extract of whole plant of *Rumex acetosa* L. (Polygonaceae) and its fractions by Ames test using NPD as a mutagen for *S. typhimurium* TA98 and NaN<sub>3</sub> for *S. typhimurium* TA100.

The most active fraction was a methylene chloride fraction, showing 96.6% antimutagenic activity against NPD and 61.6% activity against NaN<sub>3</sub> at a concentration of 1.0mg per plate. Cytotoxicity of the methanol extract and its fractions against five cultured human tumor cell lines, A549(non small cell lung), SK-OV-3(ovary), SK-MEL-2(melanoma), XF498(central nerve system) and HCT-15(colon) was examined *in vitro*. Among the tested samples, the methylene chloride fraction was most effective, exhibiting IC<sub>50</sub> values of 13.17, 13.46, 18.73, 18.35 and 17.62µg/ml against above cell lines, respectively. These results suggest that the methylene chloride fraction possesses potent antimutagenic and/or anticancer constituents.

[PD2-34] [ 04/20/2001 (Fri) 13:30 - 14:30 / Hall 4 ]

### Inhibitory effects of Hydrolyzable tannins on Melanin Biosynthesis in B16 mouse melanoma cell line

Cho SM, Lee YA, Kim KH, Lee JH, Kim JH, Kwon YM, Lee MW

College of Pharmacy, Chung Ang University, Seoul 156-756, Korea

For the utilizing of hydrolyzable tannins in the whitening-effect cosmetics, inhibitory effects of hydrolyzable tannins on melanin biosynthesis in B16 mouse melanoma cell line were determined. These hydrolyzable tannins showed inhibitory effect against tyrosinase previously. Hydrolyzable tannins especially, 2,3-(S)-HHDP-D-glucose and pedunculagin inhibited melanin biosynthesis in a

dose-dependent manner.

[PD2-35] [ 04/20/2001 (Fri) 13:30 – 14:30 / Hall 4 ]

### Virus-cell fusion inhibitory activity for the polysaccharides from various Korean edible clams

Woo E.-R.<sup>01</sup>, Kim WS<sup>2</sup>, and Kim YS<sup>2</sup>

College of Pharmacy, Chosun University, Kwang-ju 501-759, Korea<sup>1</sup>, Nat. Prod. Res. Inst. Seoul National University, Seoul 110-460, Korea<sup>2</sup>

In order to find potent virus-cell fusion inhibitory components from Korean edible clams, thirteen prepared polysaccharides were introduced to syncytia formation inhibition assay, which is based on the interaction between the HIV-1 envelope protein gp120/41 and the cellular membrane protein CD4 of T lymphocytes. Among them, Meretrix petechialis showed a potent virus-cell fusion inhibitory activity. Fusion index (FI) and percent (%) fusion inhibition of the polysaccharide of this clam were  $0.21 \pm 0.02$ , and  $67.52 \pm 4.09$  at 100  $\mu\text{g/mL}$ , respectively. It exhibited almost equivalent virus-cell fusion inhibitory activity to that of dextran sulfate which was used as a standard control.

[PD2-36] [ 04/20/2001 (Fri) 13:30 – 14:30 / Hall 4 ]

### Antioxidative compounds from Clerodendrii folium

Ham I, Chung M.Y, Lee S.J, Kim H.H, Sohn U.D, and Whang W.K

ChungAng University, College of Pharmacy

*Clerodendron trichotomum*(Verbenaceae) has been used for arthritis, rheumatism, and hypertension as a folk medicine.

In order to evaluate anti-oxidative activity, its fractions(H<sub>2</sub>O, 30%, 60%, 100Fr.) were measured with DPPH method. It was revealed that 30% and 60% MeOH fractions have significant antioxidative activity.

From 30% MeOH, five phenolic compounds were isolated by column chromatography and elucidated two flavonoid glycosides and three phenyl propanoids compounds.

[PD2-37] [ 04/20/2001 (Fri) 13:30 – 14:30 / Hall 4 ]

### Anti-oxidative activity of Ban-Lan-Gen

Jang HJ<sup>0</sup>, Kim EJ, Yang KS

College of Pharmacy, Sookmyung Women's University

Ban-Lan-Gen is the dried root of *Isatis indigotica* Fort.(Cruciferae) and one of the most commonly-used traditional chinese medicines for antipyretic, antiviral, and detoxifying purpose in china.

In order to evaluate anti-oxidative activity, Ban-Lan-Gen was fractionated and