

Park WonBong<sup>0</sup> Kim JongHyun Rhim JeeYoung Jung SeungHee Moon YouSun Lee GyuYim Lyu SuYun  
Shim ChangSup

College of Natural Science, Seoul Women's University, Seoul 139-774, Korea

Aloe and propolis have been used for thousand of years in folk medicine for several purposes. They possess several biological activities such as anti-inflammatory, antifungal, antiviral and regenerative. Although the antibacterial activity of propolis has already been demonstrated, very few studies have been done on clinical relevance in dentistry. We investigated antimicrobial activity of ethanol extract of aloe and propolis and water soluble fraction of propolis against oral pathogens. We also investigated the synergistic effect of aloe and propolis. Three microorganisms were used as follows: Streptococcus mutans, Enterococcus faecalis, Enterococcus hirae. The antimicrobial activity was tested by serial broth dilution method, and the antimicrobial activity was expressed by minimal inhibitory concentration (MIC). To investigate the synergistic effect of aloe and propolis, the extract of propolis diluted serially was used for each strain. The ethanol extract of aloe showed weak antimicrobial activity, while both of ethanol extract and water-soluble fraction of propolis inhibited greatly all microorganisms tested. However, the ethanol extract of aloe enhanced significantly antimicrobial activity of propolis against oral pathogens.

[PC2-11] [ 10/19/2001 (Fri) 09:00 - 12:00 / Hall D ]

### **A NAT for reliable HCV RNA screening of blood**

Hong SeungHee, Jung SaRah, Park SuJin, Yoo SiHyung<sup>0</sup>, Kim SunNam, Lee KiHong, Kang HyeNa, Shin InSoo, Choi SeungEun, Lee SeokHo, Hong SeungHwa

Biologics Evaluation Department, Korea Food and Drug Administration

Current methods to detect hepatitis C virus (HCV) are based on its antibody detection. Viral contamination of a donated blood can potentially escape such detection during the window phase of infection, when no antibody is present or the level of antibody is too low to detect. Application of nucleic acid amplification technology (NAT) for the direct detection of HCV can actually reduce the window period and contribute to the safeguard of blood and blood products. The objective of this study was to develop a highly sensitive in-house method for the HCV RNA screening via NAT. The superiority of the protocol being developed was compared with commercial methods as control, and was applied to blood samples. In this study, five sets of primers were designed and then two sets of primers were selected for RT-PCR. We have found that QIAamp viral RNA isolation kit is the most efficient extraction kit for these systems after several PCR conditions such as annealing temperature, reverse transcription temperature, MgCl<sub>2</sub> concentration, etc. were optimized. In order to determine the positive cut-off point, a diluted series of the WHO HCV International Standard (96/790) were tested under these conditions, and the detection limit was calculated to be 5 IU . In order to validate the specificity of this analytical procedure, we are performing the test using HCV RNA negative plasma pools which has already been confirmed by European Medicine Evaluation Agency (EMEA).

[PC2-12] [ 10/19/2001 (Fri) 09:00 - 12:00 / Hall D ]

### **Immunological characterization and localization of the alcohol- dehydrogenase in Streptococcus pneumoniae**

Hyog-young Kwon<sup>0</sup>, Youn-Jin Park, Suhk-Neung Pyo, Dong-Kwon Rhee

College of Pharmacy, Sungkyunkwan University

Heat shock proteins serve as chaperone by preventing the aggregation of denatured proteins and promote survival of pathogens in harsh environments. In bacteria, ethanol shock induced the major chaperone GroEL and DnaK, but in Streptococcus pneumoniae, it induced neither GroEL nor DnaK but

alcohol dehydrogenase (ADH). In this study, ADH gene encoding a 104-kDa (p104) protein was identified and characterized. The deduced amino acid sequence of pneumococcal ADH shows homology with other members of the ADH family, and particularly with *Entamoeba histolytica* ADH2 and *E. coli* ADH. *S. pneumoniae* adh is composed of 883 amino acids and its estimated isoelectric point is 6.09. Although ADH is conserved between *S. pneumoniae* and *E. coli*, immunoblot analysis employing antisera raised against pneumococcus ADH demonstrated no cross-reactivity with ADH analog in *Escherichia coli*, *Staphylococcus aureus* and human HeLa cells. Also secretion of ADH was demonstrated by subcellular fractionation and immunoblot analysis of proteins. These results suggest that *S. pneumoniae* ADH could be a highly feasible candidate for both diagnostic marker and vaccine.

Poster Presentations – Field C3. Cell Biology

[PC3-1] [ 10/19/2001 (Fri) 09:00 – 12:00 / Hall D ]

**Caspase-dependent apoptosis by naphthoquinone analog in HL-60 cells**

Mun JungYee<sup>o</sup> Chun YoungJin Kim mieyoung

Division of Biochemistry, College of Pharmacy, Chung-Ang University

Menadione has been known to exhibit a broad spectrum of antitumor activity in rodent and human cancer cells. Previous study showed that 2,3-dichloro-5,8-dihydroxy -1,4-naphthoquinone, one of the synthesized naphthoquinone analogs, have a anticancer effect on mouse leukemic L1210 and sarcoma-180 cells. Here we investigated the cellular effects and biochemical changes by naphthoquinone analog in human leukemic HL-60 cells. Naphthoquinone analog(NA) induced apoptotic cell death in HL-60 cells, which was shown by DNA ladder of fragments, a characteristic morphological change associated with apoptotic cells. NA induced the activation of caspases, release of mitochondrial cytochrome c into cytosol and upregulation of pro-apoptotic Bax protein but had no effect on anti-apoptotic proteins like Bcl-2 and Bcl-xL. The caspase inhibitor, z-VAD-FMK inhibited caspase activation and Bid cleavage by naphthoquinone analog but not cytochrome c release. These results show that naphthoquinone analog induces apoptosis through activating caspases and regulating Bcl-2 family proteins in HL-60 cells.

[PC3-2] [ 10/19/2001 (Fri) 09:00 – 12:00 / Hall D ]

**Studies on the Anticarcinogenic effects of Solanum tuberosum extracts**

Park YunJa<sup>o</sup> Bae SongJa

신라대학교 식품영양학과

Studies on the Anticarcinogenic effects of Solanum tuberosum extracts

Park YunJa, Bae SongJa  
Dept. Food and Nutrition, Silla university

In this study, we performed to investigate the effects of cytotoxicity and quinone reductase(QR) induced activity of Potato(*Solanum tuberosum*) peel extracts on several human cancer cells, such as HepG2, HeLa and MCF-7. We extracted the peel of *Solanum tuberosum*(STP) with methanol and the methanol extract(STPM) was partitioned with n-hexane(STPMH), ethylether(STPMEE), ethylacetate(STPMEA), n-