

A new lectin was purified from the larvae of *Allomyrina dichotoma* by physiological saline extraction, salt fractionation, anion exchange column chromatography on DEAE Sephadex A-50 and gel filtration column on Sephadex G-200. This purified lectin was designated as ADL. Several biochemical properties of ADL were characterized as follows: ADL showed single band on SDS-PAGE and agglutinated the erythrocytes of human and rabbit. Agglutination-ability was relatively stable at basic pH, temperature below 40°C, and was not affected by metal ions. This lectin was proved to be a glycoprotein containing 0.47% of sugars. The molecular weight of ADL was estimated to be 97,000 by SDS-PAGE. The gene expressions of 5 cytokines (IL-1, IL-2, IL-6, IFN $\gamma$ , TNF $\alpha$ ) from human peripheral blood mononuclear cells, stimulated with ADL, were investigated by RT-PCR and the productions of the cytokines were measured by ELISA. ADL induced the highest secretion of IL-2 at 8hr, TNF $\alpha$  at 4hr, and IFN $\gamma$  at 24hr, respectively. This lectin was proved to be a potent agglutinin for cancer cells such as HeLa, L929 and L1210.

[PD2-39] [ 10/19/2001 (Fri) 14:00 - 17:00 / Hall D ]

### **Antioxidative activity screening of Herbal drugs**

Kim SeongEun<sup>o</sup>, Son HyeJung, Kang EunYoung, Jang HyunJin, Yang KiSook

College of Pharmacy, Sookmyung Women's University

Recently, oxygen free radical injury and lipid peroxidation have been suggested as major causes of atherosclerosis, cancer, liver disease and the aging process. In order to search for antioxidants from the plants, MeOH extracts from about 80 herbal medicines were investigated. The DPPH radical scavenging activity and lipid peroxidation inhibitory activity of each extracts were measured. As a result, *Ulmus parvifolia macrocarpa*, *Trogopterus xanthipes*, *Myristica fragrans*, *Amomum tsao-ko* showed relatively strong antioxidative activities.

[PD2-40] [ 10/19/2001 (Fri) 14:00 - 17:00 / Hall D ]

### **Free Radical Scavenging and Hepatoprotective Compounds in vitro of Mori Ramulus**

Ko EunKyung<sup>o</sup>, Oh HyunCheol, Kim MiHee, Lee ByungHoon, Sohn DongHwan, An NyunHyeong, Kim YounChul

MRRC and College of Pharmacy, Wonkwang University, Professional Graduate School of Oriental Medicine, Wonkwang University

Isolation and structure elucidation of free radical scavenging and hepatoprotective compounds of Mori Ramulus was investigated. 1,1-Diphenyl-2-picrylhydrazyl (DPPH) was used for free radical scavenging activity, and protective effect against tacrine-induced cytotoxicity in human liver-derived Hep G2 cells was used for hepatoprotective activity. Assay-guided fractionation of an EtOH extract of Mori Ramulus furnished three compounds which are two prenylated flavonoids and a stilbene.

[PD2-41] [ 10/19/2001 (Fri) 14:00 - 17:00 / Hall D ]

### **Hepatoprotective constituents from *Hedyotis diffusa* and *Gardenia jasminoides***

Kim Chul Young, Yoon Kee Dong, Kim Jinwoong, Lee Eun Ju, Kim Seung Hyun, Kim Young Choong

College of Pharmacy, Seoul National University