

Seo Eun-Kyoung<sup>o</sup>, Kim Kyeong Ho, Kim Min Ki, Choi EunWook, Kim KiNam, Lee Hyun-Tai, Choi Hye-Young, Han Ah-Reum, Mar Woongchon

Natural Products Chemistry Laboratory, College of Pharmacy, Ewha Womans University, College of Pharmacy, Kangwon National University, Natural Products Research Institute, Seoul National University

A prenylated coumarin, osthenol and a sesquiterpene, bisabolangelone have been isolated as active principles with 5 $\alpha$ -reductase inhibitory effect from the roots of *Angelica koreana* Max. by bioassay-guided chromatographic fractionation. Osthenol showed a strong inhibitory effect on 5 $\alpha$ -reductase type I (IC<sub>50</sub> = 14.6  $\mu$ g/ml), and especially exhibited a highly potent inhibitory activity on 5 $\alpha$ -reductase type II with an IC<sub>50</sub> value of 0.1  $\mu$ g/ml, which is about 200 times more potent than the positive control, finasteride (IC<sub>50</sub> = 19.8  $\mu$ g/ml). Bisabolangelone also inhibited the activity of 5 $\alpha$ -reductase type II (IC<sub>50</sub> = 11.6  $\mu$ g/ml), indicating that these compounds are possible candidates for development of new drugs to treat prostate disease and other androgen-sensitive conditions. In addition, four compounds including isoxypeucedanin, oxypeucedanin hydrate, oxypeucedanin and isoimperatorin were also isolated and found to be inactive in the 5 $\alpha$ -reductase assay systems used in the present study.

[PD2-48] [ 10/19/2001 (Fri) 14:00 - 17:00 / Hall D ]

### Antibiotics Resistance Inhibition of *Staphylococcus aureus* SA2 by Some Natural Products

Kim HyeKyung<sup>o</sup>, Lee ChungKyu, Moon KyungHo

College of Pharmacy, Kyungsung University, Busan, Korea

Numerous antimicrobial agents or antibiotics were developed and introduced but they lose its activity soon owing to the development of resistance to antibiotics by microorganism. In order to maintain the effectiveness of antibiotics, the reduction of resistance also is thought to be valuable as well as developing newer and stronger antibiotics.

Some essential oil components of plants showed potent inhibitory activities against multi-drug resistant microorganisms such as *Staphylococcus aureus* SA2, which has resistances to 10 usual antibiotics including chloramphenicol(Cm). Acorenone showed the strongest resistant inhibitory activity at the level of 5 g/ml when combined with 50 g/ml of Cm. Carvone and dillapiol showed resistant inhibition at the level of 50~100 g/ml when combined with 100 or 50 g/ml of ampicillin or Cm, respectively.

[PD2-49] [ 10/19/2001 (Fri) 14:00 - 17:00 / Hall D ]

### Binding affinity of some herbal extracts on the muscarinic acetylcholine receptor subtype 1 (mAChR-M1)

Kim YoungSup, Kim JeoungSeob, Kim SeongKie, Heor Junghee, Lee ByungEui, Ryu ShiYong

Bio-Organic Division, Korea Research Institute of Chemical Technology, Taejon 305-343, Korea

The water extracts of 82 kind of Korean medicinal herbs were examined for the binding affinity on the recombinant human muscarinic acetylcholine receptor subtype 1 (mAChR-M1) produced from the CHO (Chinese Hamster Ovary) cell line. Among the tested, the extract of *Coptidis Rhizoma*, *Phellodendri Cortex*, *Hedyotis Herba* and of *Terminariae Fructus* were found to exhibit a significant competition with [<sup>3</sup>H] N-methyl-scopolamine for the specific binding to mAChR-M1 in a dose dependent manner, respectively.