

Metabolic Engineering of Nonmevalonate Pathway in *Escherichia coli* Enhances Lycopene Production

Seon-Won Kim¹ and J. D. Keasling²

¹Dept. of Food Science and Nutrition Gyeongsang National University Chinju, 660-701 Korea

²Dept. of Chemical Engineering UC Berkeley, CA 94720-1462 USA

Abstract

Isopentenyl diphosphate (IPP) is the common, five-carbon building block in the biosynthesis of all carotenoids. IPP in *Escherichia coli* is synthesized through the non-mevalonate pathway. The first reaction of IPP biosynthesis in *E. coli* is the formation of 1-deoxy-D-xylulose-5-phosphate (DXP), catalyzed by DXP synthase and encoded by *dxs*. The second reaction in the pathway is the reduction of DXP to 2-C-methyl-D-erythritol-4-phosphate, catalyzed by DXP reductoisomerase and encoded by *dxr*. To determine if one or more of the reactions in the non-mevalonate pathway controlled flux to IPP, *dxs* and *dxr* were placed on several expression vectors under the control of three different promoters and transformed into three *E. coli* strains (DH5 α , XL1-Blue, and JM101) that had been engineered to produce lycopene. Lycopene production was improved significantly in strains transformed with the *dxs* expression vectors. When the *dxs* gene was expressed from the arabinose-inducible *araBAD* promoter (P_{BAD}) on a medium-copy plasmid, lycopene production was 2-fold higher than when *dxs* was expressed from the IPTG-inducible *trc* and *lac* promoters (P_{trc} and P_{lac} , respectively) on medium-copy and high-copy plasmids. Given the low final densities of cells expressing *dxs* from IPTG-inducible promoters, the low lycopene production was probably due to the metabolic burden of plasmid maintenance and an excessive drain of central metabolic intermediates. At arabinose concentrations between 0 and 1.33 mM, cells expressing both *dxs* and *dxr* from P_{BAD} on a medium-copy plasmid produced 1.4 - 2.0 times more lycopene than cells expressing *dxs* only. However, at higher arabinose concentrations lycopene production in cells expressing both *dxs* and *dxr* was lower than in cells expressing *dxs* only. A comparison of the three *E. coli* strains transformed with the arabinose-inducible *dxs* on a medium-copy plasmid revealed that lycopene production was highest in XL1-Blue.

Introduction

Carotenoids are naturally occurring pigments synthesized as hydrocarbons (carotenes) or oxygenated derivatives (xanthophylls) by plants and microorganisms. Their major functions are protecting against oxidative damage by quenching photosensitizers, interacting with singlet oxygen (Krinsky, 1994), and scavenging peroxy radicals (Conn et al., 1992), thus preventing the accumulation of harmful oxygen species. Recently, carotenoids have been successfully synthesized in non-carotenogenic bacteria and yeast using recombinant DNA techniques (Misawa et al., 1991; Sandmann et al., 1990; Yamano et al., 1994).

One of the key factors for high-yield carotenoid production is a sufficient supply of precursors. All carotenoids are synthesized via a common metabolic precursor, isopentenyl diphosphate (IPP; C₅). Until

recently, the biosynthesis of IPP was generally assumed to proceed exclusively from acetyl-CoA via the classical mevalonate pathway (McCaskill and Croteau, 1998). Recent studies have demonstrated that mevalonate is not the biosynthetic precursor of IPP in all living organisms (Horbach et al., 1993; Rohmer et al., 1993). The existence of an alternative, mevalonate-independent pathway for IPP formation was characterized initially in several species of eubacteria (Rohmer et al., 1993; Rohmer et al., 1996) and a green alga (Schwender et al., 1996). The first reaction in the non-mevalonate pathway is the transketolase-type condensation reaction of pyruvate and D-glyceraldehyde-3-phosphate to yield 1-deoxy-D-xylulose-5-phosphate (DXP). A gene (*dxr*), which encodes DXP reductoisomerase responsible for the reduction of DXP to 2-C-methyl-D-erythritol-4-phosphate (MEP), the proposed second step in the nonmevalonate pathway, was cloned from *E. coli* (Takahashi et al., 1998). It may be possible to increase the supply of IPP by enhancing DXP synthase and DXP reductoisomerase enzymatic activities.

Materials and Methods

Bacterial strains, and plasmids

E. coli DH5 α , XL1-Blue and JM101 were used for lycopene production. Plasmids pTrc99A, pBluescript, and pBAD24 were used as vectors for *dxs* and *dxr* expression studies. Plasmid pAC-LYC04 expresses the *Erwinia herbicola crtE*, *crtB* and *crtI* genes necessary for lycopene biosynthesis in *E. coli*. This plasmid also contained the *Haematococcus pluvialis ipi* gene that encodes IPP isomerase.

PCR Cloning of *dxs* and *dxr* genes

The *dxs* and *dxr* genes were cloned by the PCR from genomic DNA of *E. coli*. Plasmid pTrc99A containing *dxs* and *dxr* were designated pTdxs and pTdxr, respectively. Plasmids pDdxs and pDdxs/r were pBAD24 containing *dxs*, and both *dxs* and *dxr*.

Growth and induction experiments

Bacterial growth was determined by measuring the optical density at a wavelength of 600 nm (OD₆₀₀). A seed culture was made by inoculating cells into 2 \times YT medium containing 100 μ g/ml ampicillin and 50 μ g/ml chloramphenicol and growing the cells overnight at 37°C. An aliquot of the seed culture was inoculated into 5 ml of 2 \times YT medium containing 100 μ g/ml ampicillin and 50 μ g/ml chloramphenicol to an OD₆₀₀ of 0.1, and incubated at 29°C for 24 h. IPTG and arabinose induction studies were initiated at an OD₆₀₀ of 0.8 (unless otherwise stated).

Results

Expression of *dxs* and *dxr* using pTrc99A

E. coli cells transformed with pAC-LYC04 are pigmented pink due to the accumulation of lycopene. *E. coli* DH5 α engineered to produce lycopene were transformed with either pTdxs, pTdxr, or pTrc99A (as a control) to determine the effect of *dxs* and *dxr* overexpression on lycopene biosynthesis. In the absence of induction, cells harboring pTdxs exhibited a two-fold increase in lycopene production over cells harboring pTrc99A (Fig. 1). When IPTG was added to the culture, cell growth and lycopene




Fig. 1. Effect of IPTG concentration on lycopene production and cell growth in *E. coli* DH5 α harboring pTdxs (solid bar), pTdxr (gray bar), or pTrc99A (open bar).




Fig. 2. Effect of IPTG concentration on lycopene production and cell growth in *E. coli* DH5 α harboring pBdxs (solid bar) or pBluescript (open bar).

production decreased, even in presence of 0.1 mM IPTG, below levels produced by cells harboring either pTdxr or pTrc99A. In contrast, regardless of the amount of inducer added to the culture, *dxr* expression had no significant effect on lycopene production or cell growth.

Expression of *dxs* using pBluescript

Because the *trc* promoter of pTrc99A was too strong to control expression of *dxs*, we introduced the *dxs* gene into pBluescript, which carries the weaker *lac* promoter. As with uninduced cells harboring pTdxs, cells harboring pBdxs produced more lycopene than cells harboring pBluescript (Fig. 2). In contrast to the results with the *trc* promoter, there was no difference in lycopene production or cell growth at any IPTG concentration used. As 2 \times YT medium is very rich and there is no *lacI* gene in pBluescript nor in *E. coli* DH5 α , the expression of *dxs* on pBdxs was fully induced in the absence of IPTG. Lycopene production by the pBdxs-containing cells was 3.5-fold higher than by the cells harboring pBluescript (control), whereas cell growth was similar. However, growth and lycopene production by the pBdxs-containing culture were lower than by *E. coli* DH5 α transformed with pTdxs but not induced (Fig. 1). These differences were probably not due to the expression of *dxs*, as cell growth and lycopene production in transformants harboring pBluescript were lower than that by transformants harboring pTrc99A. As pBluescript is a high-copy-number vector and pTrc99A is a medium-copy-number vector, the added metabolic burden from the higher copy number of pBluescript relative to pTrc99A may result in lower cell growth and lycopene production. Ruther et al. (1997) reported similar results.

Lycopene production in *E. coli* transformed with pDdxs or pDdxs/r

Because expression of *dxs* from pTrc99A and pBluescript was difficult to control, we introduced the *dxs* gene into pBAD24, a medium-copy-number plasmid containing the arabinose-inducible *araBAD* promoter (P_{BAD}). In contrast to the results obtained using the IPTG-inducible systems, the amount of lycopene produced by the cells increased with the arabinose concentration (Fig. 3). Furthermore, cell growth was unaffected by the amount of arabinose added to the culture. When induced with 13.3 mM




Fig. 3. Effect of arabinose concentration on lycopene production and cell growth in *E. coli* DH5α harboring pDdxs (solid bar), pDdxs/r (gray bar), or pBAD24 (open bar).




Fig. 4. Comparison of lycopene production (solid bars) and growth (open bars) in *E. coli* strains DH5α, XL1-Blue, and JM101 transformed with pBAD24 or pDdxs. Cultivation was carried out for 48 hours and 13.3 mM arabinose was added initially for induction.

arabinose, *E. coli* transformed with pDdxs produced 12.3 mg lycopene/l in 24 hours. This lycopene production is four times higher than the control and significantly higher than any other genetic construct. Lycopene production in the absence of inducer was slightly higher in strains harboring pDdxs than those harboring pBAD24, most likely due to leaky expression from P_{BAD}.

Since the amplification of *dxs* expression increased production of lycopene, we postulated that further enhancement of lycopene production could be limited by other reactions in the pathway. To determine if amplification of *dxr* expression could improve lycopene production in strains harboring pDdxs, the *dxr* gene was combined with the *dxs* of pDdxs to obtain pDdxs/r. At arabinose concentrations between 0 and 1.33 mM, cells expressing both *dxs* and *dxr* from P_{BAD} produced 1.4-2.0 times more lycopene than cells expressing *dxs* only. However, at higher arabinose concentrations lycopene production in cells expressing both *dxs* and *dxr* was lower than in cells expressing *dxs* only. (Fig. 3). There was no significant difference in the growth of cells harboring pDdxs or pDdxs/r at all arabinose concentrations.

Comparison of lycopene production in different *E. coli* strains transformed with pDdxs or pBAD24

E. coli strains DH5α, XL1-Blue, and JM101 harboring pAC-LYC04 were transformed with either pBAD24 or pDdxs to compare lycopene production in these strains. Lycopene production was highest in XL1-Blue whereas no significant differences in cell growth were observed for all these strains (Fig. 4).

References

- Conn, P. F., Lambert, C., Land, E. J., Schach, W., and Truscott, T. G. 1992. Carotene-oxygen radical interactions [published erratum appears in *Free Radic Res Commun* 1993; 18(4):253]. *Free Radic Res Commun* 16: 401-8.
- Horbach, S., Sahn, H., and Welle, R. 1993. Isoprenoid biosynthesis in bacteria: two different pathways? *FEMS Microbiol Lett* 111: 135-40.
- Krinsky, N. I. 1994. The biological properties of carotenoids. *Pure Appl Chem* 66: 1003-10.

- McCaskill, D. and Croteau, R. 1998. Some caveats for bioengineering terpenoid metabolism in plants. *TIBTECH* 16: 349-55.
- Misawa, N., Yamano, S., and Ikenaga, H. 1991. Production of beta-carotene in *Zymomonas mobilis* and *Agrobacterium tumefaciens* by introduction of the biosynthesis genes from *Erwinia uredovora*. *Appl Environ Microbiol* 57: 1847-9.
- Rohmer, M., Knani, M., Simonin, P., Sutter, B., and Salm, H. 1993. Isoprenoid biosynthesis in bacteria: a novel pathway for the early steps leading to isopentenyl diphosphate. *Biochem J* 295: 517-24.
- Rohmer, M., Seemann, M., Horbach, S., Bringer-Meyer, S., and Sarm, H. 1996. Glyceraldehyde-3-phosphate and pyruvate as precursors of isoprenic units in an alternative non-mevalonate pathway for terpenoid biosynthesis. *J Am Chem Soc* 118: 2564-6.
- Ruther, A., Misawa, N., Boger, P., and Sandmann, G. 1997. Production of zeaxanthin in *Escherichia coli* transformed with different carotenogenic plasmids. *Appl Microbiol Biotechnol* 48: 162-7.
- Sandmann, G., Albrecht, M., Schnurr, G., Knorz, O., and Boger, P. 1999. The biotechnological potential and design of novel carotenoids by gene combination in *Escherichia coli*. *Trends Biotechnol* 17: 233-7.
- Schwender, J., Seemann, M., Lichtenthaler, H. K., and Rohmer, M. 1996. Biosynthesis of isoprenoids (carotenoids, sterols, prenyl side-chains of chlorophylls and plastoquinone) via a novel pyruvate/glyceraldehyde 3-phosphate non-mevalonate pathway in the green alga *Scenedesmus obliquus*. *Biochem J* 316: 73-80.
- Takahashi, S., Kuzuyama, T., Watanabe, H., and Seto, H. 1998. A 1-deoxy-D-xylulose 5-phosphate reductoisomerase catalyzing the formation of 2-C-methyl-D-erythritol 4-phosphate in an alternative nonmevalonate pathway for terpenoid biosynthesis. *Proc Natl Acad Sci U S A* 95: 9879-84.
- Yamano, S., Ishii, T., Nakagawa, M., Ikenaga, H., and Misawa, N. 1994. Metabolic engineering for production of beta-carotene and lycopene in *Saccharomyces cerevisiae*. *Biosci Biotechnol Biochem* 58: 1112-4.