

### 3-3-11. Purification, Identification and Characterization of Pupal Major Haemolymph Protein of the Chinese Oak Silkmoth, *Antheraea pernyi*

Sang Bong Park<sup>1</sup>, Soohyun Kim<sup>2</sup>, Nam Sook Park<sup>3</sup>, Hung Dae Sohn<sup>3</sup>,  
Byung Rae Jin<sup>3</sup> and Sang Mong Lee<sup>1</sup>

<sup>1</sup>Department of Sericultural and Entomological Biology, Faculty of Agriculture,  
Miryang National University

<sup>2</sup>Biomolecule Research Team, Korea Basic Science Institute. <sup>3</sup>College of Natural  
Resources and Life Science, Dong-A University

A pupal major haemolymph protein of the wild silkmoth, the Chinese oak silkworm *Antheraea pernyi* (AMHP), purified and identified. The protein AMHP was purified by a simple preparative polyacrylamide gel electrophoresis (PAGE) and diffusive elution. The preparation was shown to be homogeneous by 7.5% native-PAGE. The native molecular weight of the AMHP was 450 kDa with a 80 kDa single subunit, suggesting hexamer. The protein contained high amounts (18.3%) of aromatic amino acids, phenylalanine (9.7%) and tyrosine (8.6%). Therefore, the protein was identified as a kind of a storage protein referred to as an arylphorin. The protein was stained by Schiff's reagent, suggesting a glycoprotein. The protein contained 4.9% (w/w) sugar and mannose and N-acetylglucosamine were major components. Also, degradation of protein was begun by heat treatment at 90°C for 20 minutes. From these results, it was concluded that the AMHP is a storage protein referred to as arylphorin which is purified from the pupal haemolymph of the wild silkmoth, *A. pernyi*.