

15-Deoxy- Δ 12,14-prostaglandin J2 (15-deoxy-PGJ2), a naturally occurring ligand activates the peroxisome proliferator-activated receptor- γ (PPAR- γ). It was known to have promoting ability of nerve growth factor(NGF)-induced neurite extension. However, it is not clear yet as to what signaling pathway is involved in its promoting ability of neurite extension. Since MAP kinase classes as well as transcription factors have been known to be implicated in neuronal cell differentiation, we investigated whether 15-deoxy-PGJ2 exert its ability to promote cell differentiation through up-regulation of MAP kinase classes and the activation of transcription factors. PC 12 cells treated with 15-deoxy-PGJ2 (0.2 to 1.6 μ M) alone showed measurable neurite extension and expression of neurofilament, markers of cell differentiation. However much greater extent of neurite extension and expression of neurofilament was observed in the presence of NGF (50 ng/ml). In parallel with its increasing effect on the neurite extension and expression of neurofilament, 15-deoxy-PGJ2 enhanced NGF-induced p38 MAP kinase expression and its phosphorylation in addition to the activation of transcription factor AP-1 in a dose dependent manner. Moreover, pretreatment of SD 203580, a specific inhibitor of p38 MAP kinase inhibited the promoting effect of 15-deoxy-PGJ2 (0.8 μ M) on NGF (50 ng/ml)-induced neurite extension. This inhibition correlated well with the ability of SB203580 to inhibit the enhancing effect of 15-deoxy-PGJ2 on NGF-induced the expression of p38 MAP kinase and activation of AP-1. These data demonstrate that activation of p38 MAP kinase in conjunction with AP-1 signal pathway may play an important role in the promoting activity of 15-deoxy-PGJ2 on the NGF-induced differentiation of PC12 cells.

[PB3-6] [04/19/2002 (Fri) 10:00 - 13:00 / Hall E]

PRESENILIN-2 MUTATION ALTERS NEURITE EXTENSION, APOPTOSIS AND TRANSCRIPTION FACTOR(NF-KB) ACTIVATION.

Seong Min Je^o, Song Youn Sook, Shin Im Chul, Oh Ki Wan, Kim Young Ku*, Hwang Dae Hyun*, Chung Soo Youn*, Yang Ki Hwa*, Hong Jin Tae

College of Pharmacy Chungbuk National University, *National Institute of Toxicological Research, KFDA

Alzheimer's disease (AD) is characterized by β -amyloid deposition and associated with loss of neuron cells in brain regions involved in learning and memory process. Many cases of early onset autosomal dominant inherited forms of AD are caused by mutation in the genes encoding presenilin-2 (PS-2). However, its pathogenic mechanisms in AD are not known. Here we report that expression of an AD-like human PS-2 mutation (N141I) in PC12 cells resulted in aberrant differentiation responses to nerve growth factor (NGF) and β -amyloid. NGF-induced neurite extension was significantly reduced in cells stably expressing mutant PS-2. Induction of apoptosis and apoptosis-associated gene expression by β -amyloid was markedly increased in cells carrying mutant PS-2. The DNA binding activity of the transcription factor NF-kB by β -amyloid was decreased in the cells carrying mutant PS-2. These finding shows that altered responsibility to neurotrophic (or neurotoxic) factors could a role in the pathogenesis of AD carrying PS-2 mutations.

[PB3-7] [04/19/2002 (Fri) 10:00 - 13:00 / Hall E]

L-DOPA-Induced Neurotoxicity Is Enhanced by Tetrahydropapaveroline in PC12 Cells

Lee JaeJoon^o, Kim Yu Mi, Yin shouyu, Yin Shouyu, Park HeeDong, Kang MinHee, Lee Myung Koo

College of Pharmacy, and Research Center for Bioresource and Health, Chungbuk National University

Tetrahydropapaveroline (THP) is formed in Parkinsonian patients receiving L-DOPA therapy and is detected in plasma and urine of the patients. THP and its derivatives were proposed to be candidates of dopaminergic neurotoxins related to the pathogenesis of Parkinsonism. In this study, we have investigated the effects of THP on dopamine content and L-DOPA-induced neurotoxicity in cultured rat pheochromocytoma (PC12) cells. PC12 cells were exposed to THP, L-DOPA, or a combination of the two for 24h or 48 h. THP at concentration range of 5-15 μ M decreased dopamine content in a concentration-dependent manner. L-DOPA at 20-100 μ M increased dopamine content in PC12 cells, but the increase in dopamine levels by L-DOPA was in part inhibited when L-DOPA was associated with 5-15 μ M THP. Exposure of PC12 cells up to 10 μ M THP or 20 μ M L-DOPA after 24h or 48h, neither affected cell viability, determined by MTT assay, nor induced apoptosis, by flow cytometry and TUNEL staining. However, at

concentrations higher than 15 μ M, THP showed a cytotoxicity through an apoptotic process. In addition, THP at 5–15 μ M significantly enhanced L-DOPA-induced neurotoxicity (L-DOPA concentration, 50 μ M). Treatment of PC12 cells with 15 μ M THP and 50 μ M L-DOPA, alone or in combination, also induced cell death via a mechanism which exhibited morphological and biochemical characteristics of apoptosis, including chromatin condensation and membrane blebbing. Exposure of PC12 cells to THP, L-DOPA and THP plus L-DOPA for 48 h resulted in a marked increase in the cell loss and percentage of apoptotic cells compared with exposure for 24 h. These findings indicate that the enhancing effects of THP on L-DOPA-induced neurotoxicity were time and concentration dependent. Furthermore, these results suggest that THP inhibits L-DOPA-induced increase in dopamine content and enhances L-DOPA-induced neurotoxic and apoptotic effects on PC12 cells.

[PB3-8] [04/19/2002 (Fri) 10:00 – 13:00 / Hall E]

Increase of intracellular Ca²⁺ and cytotoxicity induced by neuro-toxicants in PC12 cells carrying mutant presenilin-2

Shin Im Chul^o, Hwang In Young, Song Youn Sook, Park Cheol Beom, Oh Ki Wan, Lee Myung Koo, Kim Young Ku*, Hong Jin Tae

Department of Pharmacy, College of Pharmacy, Chungbuk National University, Cheongju, 361-763, Korea,*National Institute of Toxicological Research, KFDA

Many cases of early onset autosomal dominant inherited forms of Alzheimer's disease (AD) are caused by mutation in the genes encoding presenilin-2 (PS-2) on chromosome 1. It is characterized by amyloid deposition and associated with loss of neuron. However, molecular mechanisms underlying the role of PS-2 mutation in the pathogenic AD are not known. Pathophysiological elevation of intracellular calcium concentration in the neuron has been demonstrated as an important responsible factor in the neuronal cell death. In this study, we compared real-time alteration of intra-cellular calcium concentration and cellular response (cytotoxicity) in the pheochromocytoma cells (PC12) and PC12 cells carrying mutant PS-2 stimulated either by beta-amyloid and glutamate. Prolonged elevation of intra-cellular calcium concentration by glutamate and beta-amyloid was significantly enhanced in cells carrying mutant PS-2. With reverse correlation with the level of intra-cellular calcium concentration, significant decrease of cell viability and increase of the induction of apoptosis was found in the cells carrying mutant PS-2. This results showing that PS-2 mutation elevates intra-cellular calcium concentration and thereby render neurons vulnerable to neuro-toxic stimuli, suggested that perturb of intra-cellular calcium homeostasis could play a important role in the pathogenesis of AD.

[PB3-9] [04/19/2002 (Fri) 10:00 – 13:00 / Hall E]

15-DEOXY- Δ 12,14-PROSTAGLANDIN J₂, A LIGAND OF PEROXISOME PROLIFERATOR ACTIVATED RECEPTOR- γ INDUCE APOPTOSIS THROUGH PHOSPHORYLATION OF ERK PATHWAY IN NEUROBLASTOMA CELLS

Kim Eun Joung^o, Sheen Yhun Yhong*, Chung Soo Youn, Seong Min Je, Hong Jin Tae**

National Institute of Toxicological Research, KFDA, *Collage of Pharmacy, Ewha Womans University, **Collage of Pharmacy, Chungbuk National University

15-Deoxy- Δ 12,14-prostaglandin J₂ (15-deoxy-PGJ₂), a peroxisome proliferator-activated receptors (PPAR- γ) ligand, has been shown to stimulate the differentiation and induction of apoptosis in the several cancer cells including breast, prostate and lung cancer cells. In the previous our study, it was found that 15-deoxy-PGJ₂ inhibit cell growth through induction of apoptosis in neuroblastoma cells (SK-N-MC and SK-N-SH cells). Here we demonstrated possible molecular mechanisms underlying the induction of apoptosis by 15-deoxy-PGJ₂. 15-Deoxy-PGJ₂ dose (2–16 μ M) dependently induced apoptosis. Consistent with the induction of apoptosis, 15-deoxy-PGJ₂ reduced the expression of anti-apoptotic gene Bcl-2 but increased the expression of pro-apoptotic genes : caspase-3 and 9, and Bax. In parallel with the increasing of apoptosis, 15-deoxy-PGJ₂ increased the expression of phosphorylated ERK. Moreover,