

## Antioxidative Phenolic Compounds from the Cambodian Mushroom *Phelinus Linteus*

Lee SooYoung<sup>o</sup>, Ryu GeonSeek, Choi ByoungWook, Lee BongHo

Dep. of Chemical Technology, Hanbat National University

In continuing search for antioxidative compounds from natural sources, we found an antioxidative activity in the methanolic extract of the Cambodian mushroom *Phelinus Linteus*. After partitioning between  $\text{CHCl}_3$  and 30% MeOH, the former layer was purified by a series of ODS flash, Shephadex LH-20, silica column chromatography to give two antioxidative phenolic compounds, 4-(3,4-dihydroxyphenyl)-(E)-3-buten-2-one (1) and caffeic acid (2). Compounds 1 and 2 like stilbenes or flavonoids with a catechol moiety showed high radical scavenging effect on DPPH radicals with  $\text{RC}_{50}$  values of 7.9 and 8.7  $\mu\text{g}/\text{mL}$ , respectively.

[PD2-34] [ 04/19/2002 (Fri) 10:00 - 13:00 / Hall E ]

## An Antioxidative Dimeric C-Glucoside from *Ardisia japonica* Blume

Kim BongSeok<sup>o</sup>, <sup>1</sup> Ryu GeonSeek, <sup>2</sup> Choi ByoungWook, <sup>2</sup> Lee BongHo, <sup>2</sup> Lee NamHo, <sup>1</sup> Jung DukSang<sup>1</sup>

<sup>1</sup>Dep. of Chemistry, Cheju National University, <sup>2</sup>Dep. of Chemical Technology, Hanbat National University

In our ongoing researches for bioactive compounds from natural sources, our attentions were focused on potent antioxidants from *Ardisia japonica*. The methanolic extract of the plant was partitioned in accordance with the Kupchan's scheme. Chromatographic purification of the *n*-BuOH layer afforded an unprecedented dimeric C-glucoside (1). The molecular formula of 1 was established as  $\text{C}_{28}\text{H}_{30}\text{O}_{18}$  by the HRFAB mass and  $^{13}\text{C}$  NMR data, indicative of 13 degree of unsaturation. Its structure was elucidated by the interpretation of the NMR spectra and its stereochemistry determined by *J*-coupling constants and NOE experiments. Compound 1 exhibited moderate antioxidative activity with  $\text{RC}_{50}$  value of 4.6  $\mu\text{g}/\text{mL}$ .

[PD2-35] [ 04/19/2002 (Fri) 10:00 - 13:00 / Hall E ]

## Four Anti-tumor Sesquiterpenes from the Red Alga *Laurencia okamurae*

Park SooHee<sup>o</sup>, Ryu GeonSeek, Choi ByoungWook, Lee BongHo

Dep. of Chemical Technology, Hanbat National University

Some halogenated metabolites from marine organisms are believed to play the role of chemical defence substances against marine herbivores. Among the chemical constituents isolated from the red algae in genus *Laurencia* (Ceramiaceae, Rhodomelaceae) was an unusual tricyclic sesquiterpene named as aplysin. The co-occurrence of brominated and non-brominated aplysins in many natural sources prompted speculation that the debromo-analogues scavenged reactive halogens from the marine environment before they could inflict damage on the host. The aplysins are also believed to act as anti-feedant preventing the predatory advances of other marine organisms. As a part of our continuing searches for anti-tumor agents from Korean marine algae, we found a potent anti-tumor activity in the methanolic extract of *Laurencia okamurae*. We herein report the isolation, structural elucidation, and biogenesis of four aplysin-related sesquiterpenes. Among them, laurinterol (1) showed potent anti-tumor activity against A549, SK-OV-3, SK-MEL-2, XF498, and HT15 with  $\text{EC}_{50}$  values of 2.6-3.6  $\mu\text{g}/\text{mL}$ , respectively.

[PD2-36] [ 04/19/2002 (Fri) 10:00 - 13:00 / Hall E ]