

most frequently encountered in oligosaccharides. The results demonstrate that ELLA is a very useful tool for rapid estimation of the types and relative amounts of specific carbohydrate structures within intact glycoproteins. The microheterogeneity of some oligosaccharides was also studied by using lectins that bind to their terminal residues. These results are compared with the results from direct structural analysis of oligosaccharide using a liquid chromatography system.

[PE3-3] [04/19/2002 (Fri) 10:00 ~ 13:00 / Hall E]

Characterization and cytotoxicity of lectin from *Maackia fauriei*

Kim BumSoo, Kim YunJung^o, Kong KwangHoon, Kim HaHyung

College of Pharmacy, Chung-Ang University, Seoul 156-756, Korea, College of Natural Science, Chung-Ang University, Seoul 156-756, Korea

A lectin has been purified from the bark of the legume plant *Maackia fauriei*. *M. fauriei* agglutinin (MFA) demonstrated high homogeneity with the lectin from *M. amurensis* in its N-terminal amino acid sequence and amino acid composition, however, the carbohydrate-binding specificity was different. The hemagglutination activity of MFA with human erythrocytes was specifically inhibited by N-acetylneuraminic acid as well as by Neu5Aca2-3Galb1-4GlcNAc. The hemagglutination activity of MFA is stable at pH values from 4.04 to 7.34, and at temperatures below 45°C. MFA exerts cytotoxic effects on human breast cancer MCF-7 cells, human melanoma G-361 cells, and human liver cancer SNU-449 cells.

Poster Presentations - Field F1. Clinical Pharmacy

[PF1-1] [04/19/2002 (Fri) 10:00 ~ 13:00 / Hall E]

Urinary profiles of fatty acids and androgens in female patients with thyroid cancer

Kim KyungMee^o, Choi ManHo, Paeng KiJung, Chung BongChul

Bioanalysis and Biotransformation Research Center,
Korea Institute of Science and Technology, Seoul, Korea

In female thyroid cancer, the relationship and potential effect of 11 endogenous androgens and 8 polyunsaturated fatty acids (PUFAs) were quantitatively determined in the urine of patients with thyroid cancer and normal subjects using the gas chromatography-mass spectrometry with selected ion-monitoring (SIM) mode. The values of urinary PUFAs as the long chain fatty acid, the precursor of cholesterol, was not different between normal controls and patients with thyroid cancer. But, in female thyroid cancer, the level of docosahexaenoic acid (DHA) was increased significantly higher than those of normal subjects. The concentrations of adrenal androgens derived from the parent cholesterol were not different significantly in the two groups. However, urinary values of dihydrotestosterone (DHT, $P < 0.2$), androsterone (An, $P < 0.05$), and etiocholanolone (Et, $P < 0.05$) are different to the values of the other androgens. Especially, concentrations of An and Et were increased in patients with thyroid cancer than those of normal controls. These data indicate that fatty acids can be related to the 5 α -reductase, which is one of the enzyme in androgen metabolism.

[PF1-2] [04/19/2002 (Fri) 10:00 ~ 13:00 / Hall E]

Determination of Practical Dosing of Warfarin in Korean Outpatients with Mechanical Heart Valves

Lee Byungkoo^o Lee Ju Yeun, Jeoung Young Mi, Suh Ok Kyung, Shin Wan Gyoon, Lee Myung Koo

Department of Pharmacy, Seoul National University Hospital, College of Pharmacy, Seoul National University, College of Pharmacy, Chungbuk National University

Following implantation of heart valve prostheses, patients are prescribed life-long warfarin therapy to reduce the risk of thromboembolism. In the process of determining maintenance dose of warfarin, patient's INR variation is unavoidable.

The objective of this study was to suggest a steady-state warfarin dosing guideline for Korean outpatients with prosthetic heart valves.

A retrospective chart review was completed for all patients seen by the pharmacist-run anticoagulation service (ACS) at Seoul National University Hospital from March, 1997 to September, 2000. Patients who were at least 6 months post-valve replacement, were enrolled in the outpatient ACS and had at least one INR value less than 2.0 or greater than 3.0 were included.

Six hundred eighty-eight patients (1,782 visits) whose INR values have been previously stable while adhering to the dosing regimen, without any drug or foods that interact with warfarin, were identified and required dosing adjustment in order to achieve therapeutic INR. Of these aortic valve replacement patients (AVR, target INR: 2.0~2.5) were 151 (358 visits) and mitral valve replacement or double aortic-mitral valve replacement patients (MVR/DVR, target INR: 2.3~2.8) were 537 (1,399 visits).

To suggest the guideline for warfarin dosage adjustment, mean adjusted dose and standard deviation were calculated based on the data of the INR at the time of the event. AVR patients and MVR/DVR patients were evaluated separately.

We proposed two methods for the warfarin dosage adjustment: mg-based (Guideline I) and percentage-based (Guideline II) total weekly dose (TWD) to correct moderate INR deviation. The effectiveness of Guideline I vs Guideline II was evaluated. Guideline I was superior to Guideline II overall in patients with both AVR and MVR/DVR, but there was no statistically significant difference between two guidelines for patients receiving intermediate dose (23~47 mg TWD), the most commonly used dose.

[PF1-3] [04/19/2002 (Fri) 10:00 - 13:00 / Hall E]

Factors Affecting the Nontherapeutic INRs in Korean Outpatients with Prosthetic Heart Valves

Lee Byung koo^o Lee Ju Yeun, Jeoung Young Mi, Suh Ok Kyung, Shin Wan Gyoon, Myung Koo Lee

Department of Pharmacy, Seoul National University Hospital, College of Pharmacy, Seoul National University, College of Pharmacy, Chungbuk National University

Following implantation of heart valve prostheses, patients are prescribed life-long warfarin therapy to reduce the risk of thromboembolism. In Korean patients, maintaining a target INR of 2.0-3.0 can be challenging due to a variety of reasons. The objective of this study was to determine the causes of nontherapeutic INRs and to identify factors associated with nonadherence to warfarin therapy in Korean outpatients with prosthetic heart valves managed by a pharmacist-run anticoagulation service (ACS). A retrospective chart review was completed for all patients seen by the ACS at Seoul National University Hospital from March, 1997 to September, 2000. Patients were at least 6 months post-valve replacement were enrolled in the outpatient ACS and had at least one INR value less than 2.0 or greater than 3.0. All possible causes of nontherapeutic INRs were documented for each patient visit. The association of gender, age, duration of ACS and length of warfarin therapy to noncompliance was investigated by univariate analysis.

Eight hundred sixty-eight outpatients were seen by the ACS and 5,304 visits were identified. The reasons for nontherapeutic INRs were identified as follows: inadequate dosage adjustment (21%), nonadherence to dosing regimen (13%), drug/herbal interactions (12%), changes in diet (7%), and indeterminate cause (42%). Younger age, shorter duration of ACS and longer duration of warfarin therapy were associated with nonadherence.

In this retrospective study, nonadherence and interactions between diet and medications were found to be important factors influencing nontherapeutic INRs. Younger age and longer duration of warfarin therapy were associated with non-adherence whereas longer duration of enrollment in the ACS was associated with increased adherence to warfarin therapy.

Poster Presentations – Field F2. Social Pharmacy

No submitted abstract in the field F2 (Social Pharmacy)