

Food Organisms of Juveniles of *Tridentiger trigonocephalus* Inhabited at Intertidal Zone of the Western Coast of Korea

Jong-Yeon Kim · Soo-Gun Jo and Jong-Man Yoon

Faculty of Marine Life Science , Kunsan National University, Kunsan 573-701, Korea

ABSTRACT

To investigate the food organisms of the *Tridentiger trigonocephalus* during the juveniles stage, the stomach contents of fish, captured in the intertidal zone of Chungchongnam-do Sochon-gun Su-myon Dodun-ri between on early June from end of May 1999, were observed. Total length of the juveniles of *T. trigonocephalus* was 5.5mm~9.0mm size, and the participation rate of feeding was 68.5%. Main food organisms were such as copepods, amphipods, shrimp larvae, polychaete larvae, and these occupied more than dry weight 2%. Copepods among them dominated the most quantities by average 67.5%, and next, food organisms appeared much by order of polychaete larvae and shrimp larvae etc.. Therefore, most important food organisms of juvenile stage of *T. trigonocephalus* were copepods, polychaete larvae, shrimp larvae etc.

INTRODUCTION

Gobiidae being calculated that about 2,000 species are ranging in all over the world (Hoesse, 1985), is occupying very important location in nutrition rank of the ecosystem in coastal district and inside bay (Choi et al., 1996) .

The *T. trigonocephalus* of them is the small fish that live in the bottom, distributed much between on the shore tide pool and sunken rock in our country. In this research, distribute widely to western sea district along the coast intertidal zone, the feeding participation rate in the juveniles stage of *T. trigonocephalus* that is occupying ecologically important location was examined, and the kind and composition of food, the change of food according to the fish size, were examined from stomach contents.

MATERIALS AND METHODS

Fish fixed in neutral formalin solution 5% as soon as collect, and size of fish measured total length to 0.1 mm by standard. In order to investigate the appearance aspect of food organisms, the weight change of stomach contents was found going

through fullness of stomach. Also, fish size groups particularly, the dry weight ratio of each food organisms for whole stomach content was measured. Participation rate of feeding calculated as the rate of juveniles that feed food about numbers of total individuals. The appearance rate of food organisms was examined from the number of juveniles that each food organisms appears. To recognize the importance degree of each food organisms, multiplying the number composition of individuals of food organisms and the appearance rate, calculated the index of relative importance.

RESULTS AND DISCUSSION

If see the participation rate of feeding that ensue to total length, expressed the lowest participation rate from total length 5.5~6.0mm to 50.0%, and the highest participation rate from total length 8.5~9.0mm to 87.0%. If observe the participation rate of feeding by gathering time, showed level as much as average 71.6% in the morning, and while this decreased until noon time 54.2% becoming low gradually, after begin to increase again in the afternoon and show the maximum value by 83.5% on 3 o'clock P.M., showed low and by 52.4% on 6 o'clock P.M. the lowest value again gradually. Main food organisms observed in the stomach of *T. trigonocephalus* were such as copepods, amphipods, shrimp larvae, polychaete larvae, and these occupied more than dry weight 2%. Copepods among it dominated the most quantity by average 67.2%, and appeared much by polychaete larvae, order of shrimp larvae etc.. The composition rate of individual among food organisms of copepods is high and because the appearance rate is high, IRI represented the highest value by 5,409.6, and polychaete larvae is 646.1, shrimp larvae represented value of 362.9.

REFERENCES

- Choi, Y., I.S. Kim, B.S. Ryu. and J.Y. Park. 1996. Ecology of *Synechogobius hasta* (Pisces: Gobiidae) in the Kum River Estuary, Korea. J. Korean Fish. Soc., 29(1), 115~123.
- Hoese, D.F. 1985. Indopacific genera of gobiid fishes. Abstracts of Second International Conference on Indo-Pacific Fishes. Ichthyol. Soc. Japan pp. 60.
- Jenkins, G. P. 1987. Comparative diets, prey selection, and predatory impact of co-occurring larvae of two flounder species. J. Exp. Mar. Biol. Ecol., 110, 147~170.