

Acupuncture Stimulation for Motor Cortex Activities: Evidence from
3T Functional MRI Study

Bo-Young Choe¹ • Sei-Kwon Kang² • Myung-Ja Chu¹ • Sin-Soo Jeun³ • Seung-Schik Yoo^{1,4} •
Gi-Soon Choi⁵ • Sang-Dong Park⁵ • Eun-Chul Lim⁵ • Sung-Taek Chung⁶ •
Chang-Hyun Oh⁷ • Sun I Kim⁸ • Hyoung-Koo Lee¹ • Tae-Suk Suh¹

¹Departments of Biomedical Engineering, College of Medicine, The Catholic University of Korea,

²Young-Dong Severance Hosp. Yonsei Univ,

³Department of Neurosurgery, The Catholic University of Korea,

⁴Department of Radiology, Brigham and Womens Hospital, Harvard Medical School,

⁵Dong-Seo Hospital of Oriental Medicine, ⁶Medinus Co. LTD.,

⁷Department of Electronics and Information Engineering, Korea University,

⁸Department of Biomedical Engineering, Hanyang University

Purpose : To investigate whether or not acupuncture of GB34 produces a significant response of the modulation of somatomotor areas by functional magnetic resonance imaging (fMRI) study. The acupoint, GB34, located in the back of the knee, is known to be effective in recovering motor function after stroke.

Materials and Methods : Using 3T MRI scanner, functional MR imaging of the whole brain was performed in 12 normal healthy subjects during two stimulation paradigms; acupuncture manipulation on GB34 and sham points. This study investigates the activation of the motor cortex elicited by a soft and an intensified stimulation of GB 34. Three different paradigms were carried out to detect any possible modulation of the Blood Oxygenation Level Dependent (BOLD) response in the somatomotor area to motor stimulation through acupuncture.

Results : Group analysis from seven individuals showed that bilateral sensorimotor areas (BA 3, 4, 6 and 7) showed stimulation related BOLD signal contrast of approximately 6% whereas very few areas were activated when sham stimulation is given.

Conclusions : The present study shows that acupuncture fMRI study can be safely conducted in 3T MRI environment, and acupuncture stimulation in GB34 modulates the cortical activities of the somatomotor area in human. The present findings may shed light on the CNS mechanism of motor function by acupuncture and form a basis for future investigations of motor modulation circuits in the stroke patients.