

**Use of Metallothionein-Transgenic and Null Mice to
Determine the Role of Metallothionein in Cadmium
Toxicity**

Curtis D. Klaassen

University of Kansas Medical Center, USA

Acute Cd exposure produces hepatotoxicity, whereas chronic Cd exposure produces nephrotoxicity, hematotoxicity, immunotoxicity and bone damage. Previous experiments suggest that the low-molecular-weight, metal-binding protein metallothionein (MT) in liver protects against liver injury, but is responsible for the kidney injury observed after chronic Cd exposure. Thus, prior to the development of MT-transgenic and MT-knock-out mice models, MT's role was always assumed to be a toxicological paradox, hepatoprotection but nephrotoxicity. The development of MT-transgenic and MT-knockout mice models has reconfirmed MT's protective role against Cd-induced hepatotoxicity, but it has challenged MT's suggested role in Cd-induced nephrotoxicity. The recent data using these genetically altered mice models indicate that MT protects against not only the Cd-induced hepatotoxicity, but also nephrotoxicity, hematotoxicity, immunotoxicity, and bone damage.

Tumor Necrosis Factor- α -Null Mice Are Not Resistant to Cadmium Chloride-Induced Hepatotoxicity¹

Eric B. Harstad² and Curtis D. Klaassen

Department of Pharmacology, Toxicology, and Therapeutics, University of Kansas Medical Center, Kansas City, Kansas 66160

Received October 22, 2001; accepted December 18, 2001

Tumor Necrosis Factor- α -Null Mice Are Not Resistant to Cadmium Chloride-Induced Hepatotoxicity. Harstad, E. B., and Klaassen, C. D. (2002). *Toxicol. Appl. Pharmacol.* 179, 155–162.

Acute administration of cadmium results in hepatotoxicity. Recent reports indicate that Kupffer cells, the resident macrophages of the liver, participate in the manifestation of chemical-induced hepatotoxicity. Tumor necrosis factor- α (TNF- α) is a proinflammatory cytokine that is a major product of Kupffer cells and mediates the hepatotoxic effects of lipopolysaccharide (LPS). It has been speculated that cadmium also may exert its hepatotoxicity via the production of TNF- α by the Kupffer cells. Therefore, this study was undertaken to determine whether mice deficient in TNF- α are resistant to Cd-induced hepatotoxicity. TNF- α -null (TNF-KO) and wild-type (WT) mice were dosed ip with saline, LPS (0.1 mg/kg)/Gln (D-galactosamine, 700 mg/kg), or CdCl₂ (2.2, 2.8, 3.4, and 3.9 mg Cd/kg). Serum alanine aminotransferase (ALT) and sorbitol dehydrogenase (SDH) activities were quantified to assess liver injury. Caspase-3 activity was quantified to assess hepatocellular apoptosis. LPS/Gln treatment increased ALT (17-fold) and SDH (21-fold) in WT mice. In contrast, LPS/Gln-treatment did not significantly increase ALT or SDH in TNF-KO mice. LPS/Gln-treatment caused a 7.8-fold increase in caspase-3 activity in WT mice but did not increase caspase-3 in TNF-KO mice. Cadmium caused a dose-dependent increase in liver injury in both WT and TNF-KO mice. However, the liver injury produced by Cd in the TNF-KO mice was not different from that in WT at any dose. No significant increase in caspase-3 activity was detected in any of the Cd-treated mice. These data indicate that, in contrast to LPS/Gln-induced hepatotoxicity, TNF- α does not appear to mediate Cd-induced hepatotoxicity. © 2002 Elsevier Science (USA)

Key Words: cadmium; tumor necrosis factor; TNF; liver; hepatotoxicity; necrosis; apoptosis.

The heavy metal cadmium (Cd) is an industrial and environmental pollutant. It is toxic to several tissues, most notably causing hepatotoxicity with acute exposure and nephrotoxicity with chronic exposure. Histological evaluation of liver injury

reveals that acute Cd exposure causes swelling, congestion, pyknosis, karyorrhexis, apoptosis, and necrosis in the liver (Dudley *et al.*, 1982). However, few reports have definitively addressed the mechanism of toxicity at the molecular level. Cadmium has been shown to cause inhibition of electron transport (Diamond and Kench, 1974), DNA damage (Bagchi *et al.*, 1996), and lipid peroxidation (Stacey *et al.*, 1980; Harvey and Klaassen, 1983). These data describe some of the toxic effects of acute Cd, yet do not identify the mechanism of toxicity.

Several current reports implicate endogenous mediators in the pathogenesis of chemical-induced hepatic injury. The prevailing theory is that Kupffer cells, the resident macrophages of the liver, produce proinflammatory cytokines, such as tumor necrosis factor- α (TNF- α), and that this endogenous mediator contributes to chemical-induced hepatotoxicity. Normally, TNF- α participates in the regulation of inflammation and immunity to pathogens. However, unusually high amounts of TNF- α can result in host cell damage through its proinflammatory or cytotoxic properties. TNF- α can be directly cytotoxic to hepatocytes by inducing apoptosis and necrosis (Adamson and Billings, 1992; Tartaglia *et al.*, 1993). In addition, TNF- α initiates signal transduction pathways responsible for the production of cell adhesion molecules and chemotactic factors affecting hepatocytes and neutrophils (Witthaut *et al.*, 1994; Essani *et al.*, 1995). As a result, neutrophils adhere to damaged cells and release a host of cytotoxic intermediates, such as cytokines, proteases, and reactive oxygen species (Jaeschke *et al.*, 1996).

It is widely accepted that TNF- α is the endogenous mediator of toxicity for bacterial lipopolysaccharide (LPS) (Taniguchi *et al.*, 1997; Bopst *et al.*, 1998). The development of mice deficient in TNF- α (TNF-KO) has provided an animal model that can most accurately define specific functions of TNF- α (Marino *et al.*, 1997). Recent studies have shown TNF-KO mice to be resistant to LPS as well as LPS/D-galactosamine (LPS/Gln)-induced hepatotoxicity (Marino *et al.*, 1997; Taniguchi *et al.*, 1997; Bopst *et al.*, 1998).

TNF- α may also be involved in the manifestation of hepatotoxicity caused by nonbacterial hepatotoxins, such as carbon

¹ Supported by NIH Grant ES-01142.

² Supported by NIH Grant ES-07079.



tetrachloride (Czaja *et al.*, 1989; Laskin, 1996) and acetaminophen (Blazka *et al.*, 1995). Experiments involving suppression of Kupffer cell function with gadolinium chloride ($GdCl_3$) have shown that Kupffer cell activation appears to be involved in producing the hepatotoxicity caused by these chemicals (Edwards *et al.*, 1993; Laskin *et al.*, 1995; Michael *et al.*, 1999). TNF-KO mice are less sensitive to the hepatotoxic effects of carbon tetrachloride, demonstrating a role for TNF- α in the mechanism of chemical-induced liver injury (Morio *et al.*, 2001). In contrast, TNF-KO mice are not less sensitive to the hepatotoxic effects of acetaminophen (Boess *et al.*, 1998). These studies demonstrate that TNF- α can be involved in the mechanism of some chemical-induced as well as bacterial-induced hepatotoxicity.

Limited data indicate that TNF- α is also involved in Cd-induced hepatotoxicity. Suppression of Kupffer cells with $GdCl_3$ has been shown to decrease Cd-induced hepatotoxicity (Sauer *et al.*, 1997; Yamano *et al.*, 1998b). This observed decrease in sensitivity to Cd may be related to decreased TNF- α levels; however, $GdCl_3$ may have other protective actions in the liver. For example, $GdCl_3$ has been shown to suppress superoxide production, interfere with inducible nitric oxide synthase (iNOS) expression, and weakly induce metallothionein (Iimuro *et al.*, 1994; Roland *et al.*, 1996). A more specific method of elucidating the role of TNF- α in Cd-induced hepatotoxicity was to treat mice with anti-TNF- α antibodies prior to Cd administration. However, pretreatment with anti-TNF- α antibodies resulted in only slight protection against Cd-induced hepatotoxicity (Kayama *et al.*, 1995). Most recently, Yamano *et al.* (2000) reported that hepatic TNF- α protein concentrations were not increased after Cd administration. This evidence cannot be interpreted as conclusive proof that TNF- α is involved in Cd-induced hepatotoxicity. Consequently, the current study was designed to specifically examine the role of TNF- α in Cd-induced hepatotoxicity utilizing TNF-KO mice, which is the most conclusive model currently available. This study therefore tested the hypothesis that TNF-KO mice are resistant to the hepatotoxic effects of Cd.

MATERIALS AND METHODS

Chemicals. Cadmium chloride was obtained from Fisher Scientific Co. (Fair Lawn, NJ). Lipopolysaccharide from *Escherichia coli* 0127:B8 (TCA extract), D-galactosamine, and serum enzyme activity kits were purchased from Sigma Chemical Co. (St. Louis, MO). All other chemicals were reagent grade.

Animals. Wild-type (WT) and homozygous TNF- α -null (TNF-KO) mice (background C57BL/6 \times 129) were used throughout the study. Both strains of mice were a generous gift from Dr. Marino (Marino *et al.*, 1997) at Memorial Sloan-Kettering Cancer Center. Mice were housed in an AAALAC certified facility at $70 \pm 2^\circ F$ with a 12-h light/dark cycle and were fed laboratory mouse chow (Purina, St. Louis, MO) and water *ad libitum*. Male mice (8 weeks old, approximately 25–30 g) were dosed ip with saline or a combination of 100 $\mu g/kg$ lipopolysaccharide and 700 mg/kg LPS/Gln and hepatotoxicity was assessed 9 h later ($n = 12$). For the cadmium dose-response experiments,




FIG. 1. Hepatotoxicity following LPS/Gln administration in WT and TNF-KO mice. WT (solid bars) and TNF-KO mice (hatched bars) were administered LPS/Gln (100 μg LPS and 700 mg D-galactosamine ip). Blood and livers were collected 9 h later. Hepatotoxicity was assessed by measuring serum indices of hepatotoxicity (A) ALT and (B) SDH. (C) Liver sections were examined histologically and scored semiquantitatively for liver injury. (D) Apoptosis was assayed by quantifying caspase-3 activity. Data are presented as means \pm SE ($n = 12$). *TNF-KO mice exhibited significantly less toxicity than WT mice ($p < 0.05$).

hepatotoxicity was assessed at two different time points, 9 h after doses of 2.2, 2.8, 3.4, or 3.9 mg Cd/kg ip ($n = 12$) and 16 h after doses of 2.0, 2.4, 2.8, 3.2, 3.6, or 4.0 mg Cd/kg ip ($n = 8$). Mice were decapitated and blood was collected and processed for serum enzyme activity analysis. Livers were removed and a portion of the left lobe of the each liver was placed in 10%


ACUTE CADMIUM HEPATOTOXICITY AND TNF- α 

FIG. 2. Photomicrographs of liver sections after administration of saline or LPS/Gln in WT and TNF-KO mice. Wild-type and TNF-KO mice were administered saline or LPS/Gln (100 μ g LPS and 700 mg D-galactosamine ip). Livers were collected 9 h later, processed routinely, and stained with hematoxylin and eosin. Photomicrographs (200 \times) are representative of mouse liver from wild-type (A and B) and TNF-KO (C and D) mice 9 h after administration of saline (A and C) or LPS/Gln (B and D). Note that apoptotic bodies (arrows) are present after LPS/Gln administration.

neutral buffered formalin. The remainder of the liver was stored at -80°C until assayed for caspase-3 activity.

Serum enzyme activity assays. Biochemical evaluation of liver injury was performed by quantifying serum activities of alanine aminotransferase (ALT) and sorbitol dehydrogenase (SDH) spectrophotometrically, using Sigma test kits (Sigma Chemical Co.) according to the manufacturer's instructions.

Histopathology. Samples were taken consistently as cross-sections of the largest lobe of the liver. Liver sections, approximately 5 μm thick, were processed, stained with hematoxylin and eosin, and analyzed by light microscopy for liver injury. Grade of liver injury was analyzed semiquantitatively with six scores of severity: 0, none; 1, minimal (>2 foci of single cell necrosis per section); 2, mild (at least 5 areas of focal necrosis per section); 3, moderate (at least five foci of zonal necrosis per section); 4, severe (lobular damage, with many viable lobules per section); and 5, global (severe lobular damage, few areas of viability per section).

Caspase-3 activity assay. Liver samples were homogenized (1:5) w/v in lysis buffer (Tris-buffered saline with 1% Tween-20) using a Teflon pestle and a mortar. Homogenates were then centrifuged at 12,000g for 10 min. Supernatants were retained and stored at -80°C until assayed for caspase-3 activity. This assay was modified from Gillardon *et al.* (1997) by Harstad *et al.* (1999). Final assay mixture contained 150 μg protein, 5 mM DTT, 15 mM *N*-acetyl-Asp-Glu-Val-Asp-7-amino-4-trifluoromethyl coumarin (P-409; Biomol, Plymouth Meeting, PA), 1 μl of the protease inhibitor mixture (P-8340; Sigma

Chemical Co.), 25 mM HEPES, and 2 mM MgCl_2 in a total volume of 100 μl per reaction. After 1 h incubation at 37°C , fluorescence of the liberated fluorophore was quantified at 505 nm using a 96-well luminescence spectrometer (LS 50B; Perkin—Elmer, Norwalk, CT).

Statistics. Data from LPS/Gln-treated groups were analyzed using Student's *t* test. Data from Cd-treated groups were analyzed by two-way ANOVA.

RESULTS

LPS/Gln-induced hepatotoxicity in WT and TNF-KO mice. LPS/Gln-treatment produced a marked increase in serum ALT activity (43-fold) and SDH activity (33-fold) in WT mice, indicating extensive liver injury (Figs. 1A and 1B). Histological grade of liver injury was increased only in WT mice after LPS/Gln administration (Fig. 1C). LPS/Gln-treatment also increased caspase-3 activity in WT mice demonstrating hepatocyte apoptosis (Fig. 1D). Apoptosis and necrosis were confirmed by histology and the samples were scored semiquantitatively for liver damage. The biochemical measures of liver injury were confirmed by histological examination (Fig. 2). LPS/Gln

treatment of WT mice caused extensive liver injury, consisting of sinusoidal congestion, apoptosis, and necrosis (Fig. 2B). Endothelial cells were indiscernible, while parenchymal cells demonstrated morphology characteristic of both necrosis and apoptosis. Hepatic sinusoids were congested with erythrocytes and leukocyte infiltration was prominent. This morphology was evenly distributed throughout the liver. LPS/Gln administration to TNF-KO mice caused no observable change in histology compared to saline-treated controls (Fig. 2D). These results demonstrate that LPS/Gln treatment causes both apoptosis and necrosis in WT mice. In contrast, administration of LPS/Gln to TNF-KO mice did not result in liver injury by any of the four indices measured. Saline treatment had no deleterious effect in either WT or TNF-KO mice.

Cd-induced hepatotoxicity in WT and TNF-KO mice. Cadmium caused a dose-dependent increase in serum ALT (Fig. 3A) and SDH (Fig. 3B) activities in WT mice (Fig. 3, solid squares), indicating liver damage. Administration of 2.2 mg Cd/kg in WT mice resulted in minor increases (2-fold) in serum ALT and SDH, but no injury was apparent histologically. Dosages of 2.8, 3.4, and 3.9 mg Cd/kg increased serum ALT (16-, 19-, and 26-fold, respectively) as well as SDH (9-, 17-, and 25-fold, respectively) compared to saline control. The dose-related increase in serum enzyme activity paralleled increasing grades of liver damage (Fig. 3C). Cadmium treatment did not increase caspase-3 activity in WT mice at any dose, indicating a minimal amount of apoptosis (Fig. 3D). Histologically, Cd-induced hepatotoxicity in both WT and TNF-KO mice consisted of necrosis with no apoptosis observed histologically (not shown).

Cadmium also produced significant liver injury in TNF-KO mice (Fig. 3; open circles) 9 h after exposure. As in WT mice, administration of 2.2 mg Cd/kg did not result in hepatotoxicity. Dosages of 2.8, 3.4, and 3.9 mg Cd/kg increased serum ALT (4-, 17-, and 18-fold, respectively) as well as SDH (17-, 11-, and 17-fold, respectively) compared to saline control TNF-KO mice. Increases in ALT and SDH in TNF-KO mice were not significantly different from WT mice at any dose of Cd. The grade of liver injury increased with dose of Cd and paralleled the increases in serum enzyme activities. Cadmium treatment did not increase caspase-3 activity in TNF-KO mice. Histologically, it was not possible to discriminate between WT and TNF-KO mice after Cd administration. These data indicated that Cd-induced hepatotoxicity in TNF-KO mice was not significantly different from WT mice at any dose.

Nine hours after Cd administration, WT mice could not be histologically or biochemically discriminated from TNF-KO mice. However, it was determined that more extensive hepatotoxicity may demonstrate differential toxicity in the two mouse types. Thus, a similar experiment was conducted and liver injury was assessed at a later time (16 h after Cd administration) to determine whether the lack of difference was due

to a short exposure (Fig. 4). Cadmium produced more liver injury at 16 than at 9 h after Cd administration in WT mice. Serum ALT activity was increased (10-fold) at the dosage of 2.4 mg Cd/kg. Serum SDH activity was increased (9- and 7-fold) at dosages of 2.0 and 2.4 mg Cd/kg, respectively, compared to saline control in WT mice. Dosages of 2.8 and 3.2 mg Cd/kg in WT mice caused extensive liver damage, increasing serum ALT activity (36- and 29-fold, respectively) and SDH activity (19- and 16-fold, respectively). At dosages of 3.6 and 4.0 mg Cd/kg, the increases in serum ALT activity (36- and 29-fold, respectively) and SDH activity (20- and 16-fold, respectively) were different from saline control. The histological grade of liver injury increased in parallel to serum enzyme activities. Again, the dose-related increase in serum enzyme activity paralleled increasing grades of liver damage. Cadmium treatment did not cause an increase in caspase-3 activity at any dose, indicating that the absence of apoptosis at 9 h was not related to elapsed time.

TNF-KO mice were also sensitive to Cd-induced liver injury 16 h after Cd administration (Fig. 4). Dosages of 2.0 and 2.4 mg Cd/kg caused only minor injury, resulting in insignificant increases in ALT (2- and 2-fold, respectively) and SDH (2- and 2-fold, respectively) activities. Doses of 2.8 to 4.0 mg Cd/kg caused significantly more hepatotoxicity in TNF-KO mice. Serum ALT activity was increased 45- to 55-fold and serum SDH activity was increased 22- to 23-fold. As in WT mice, grade of liver injury in TNF-KO mice increased with dose and correlated with the increases in serum enzyme activities. Again, Cd treatment did not increase caspase-3 activity in TNF-KO mice. Histological examination confirmed the biochemical measures of liver damage in both WT and TNF-KO mice 16 h after Cd administration (Fig. 5). Dosages of 2.0 and 2.4 mg Cd/kg caused little apparent damage, consisting of only mild congestion and hydropic degeneration. Cadmium doses of 2.8 and 3.2 mg Cd/kg caused moderate liver damage. Cd-induced liver injury consisted primarily of focal necrosis as well as isolated single cell necrosis. Foci of necrosis were numerous but small, consisting of approximately 10–20 cells each. Congestion was more extensive and the presence of leukocytes was noted, although the leukocytes were primarily contained within the sinusoids. At dosages of 3.6 and 4.0 mg Cd/kg, the necrotic foci were larger and often spanned several lobules. Congestion was more severe and extended into the parenchyma as peliosis hepatis. Thus, after 16 h, Cd caused more significant liver damage in both WT and TNF-KO mice. However, the longer time point did not demonstrate that the liver injury produced by Cd in the TNF-KO mice was different from that in WT mice.

DISCUSSION

The aim of this study was to determine whether TNF- α is involved in cadmium-induced liver damage. Several indepen-

ACUTE CADMIUM HEPATOTOXICITY AND TNF- α




FIG. 3. Hepatotoxicity 9 h after administration of various dosages of Cd in WT and TNF-KO mice. WT (solid squares) and TNF-KO mice (open circles) were administered various doses of Cd (2.2, 2.8, 3.4, or 3.9 mg Cd/kg ip, $n = 12$). Blood and livers were collected 9 h later. Hepatotoxicity was assessed by measuring serum indices of hepatotoxicity (A) ALT and (B) SDH. (C) Liver sections were examined histologically and scored semiquantitatively for liver injury. (D) Apoptosis was assayed by quantifying caspase-3 activity. Data are presented as means \pm SE ($n = 12$).

dent researchers have provided strong evidence for involvement of TNF- α in liver damage produced by bacterial endotoxin (Marino *et al.*, 1997; Taniguchi *et al.*, 1997; Bopst *et al.*, 1998) and some evidence that TNF- α plays a role in acetaminophen- (Blazka *et al.*, 1995, 1996) and carbon tetrachloride-induced hepatotoxicity (Czaja *et al.*, 1989; DeCicco *et al.*, 1998; Rikans *et al.*, 1999; Morio *et al.*, 2001). These studies

have included models of Kupffer cell suppression, anti-TNF- α -antibody pretreatment, and TNF-KO mice. All of these data clearly support a role of TNF- α in hepatotoxicity produced by these chemicals. The role of TNF- α in Cd-induced hepatotoxicity is alluded to, but insufficiently defined.

TNF- α is widely accepted as the primary mediator of LPS/




FIG. 4. Hepatotoxicity 16 h following administration of various dosages of cadmium in WT and TNF-KO mice. WT (solid squares) and TNF-KO mice (open circles) were administered various doses of Cd (2.0, 2.4, 2.8, 3.2, 3.6, or 4.0 mg Cd/kg ip, $n = 8$). Blood and livers were collected 16 h later. Hepatotoxicity was assessed by measuring serum indices of hepatotoxicity (A) ALT and (B) SDH. (C) Liver sections were examined histologically and scored semiquantitatively for liver injury. (D) Apoptosis was assayed by quantifying caspase-3 activity. Data are presented as means \pm SE ($n = 8$).

HARSTAD AND KLAASSEN




FIG. 5. Photomicrographs of liver sections after administration of saline or 4.0 mg Cd/kg in WT and TNF-KO mice. Wild-type and TNF-KO mice were administered saline or 4.0 mg Cd/kg. Livers were collected 16 h after Cd administration, processed routinely, and stained with hematoxylin and eosin. Photomicrographs (200 \times) are representative of mouse liver from wild-type (A and B) and TNF-KO (C and D) mice 16 h after administration of saline (A and C) or 4.0 mg Cd/kg (B and D).

Gln-induced hepatotoxicity in WT mice. Therefore, LPS/Gln was chosen as a positive control for these experiments. LPS/Gln administration caused extensive liver damage, including both necrosis and apoptosis, in all areas of the livers of WT mice. However, necrosis and apoptosis were absent in TNF-KO mice after LPS/Gln treatment. These data are consistent with published data (Taniguchi *et al.*, 1997) and clearly confirm that TNF- α is a required component in the production of LPS/Gln-induced liver damage. In addition, this experiment demonstrates the validity of the TNF-KO mouse as an animal model that can define a role for TNF- α in the manifestation of chemical-induced hepatotoxicity.

Several studies have attempted to address the role of TNF- α in Cd-induced hepatotoxicity. Suppression of Kupffer cells prior to Cd administration protects rats against Cd-induced liver injury (Sauer *et al.*, 1997; Yamano *et al.*, 1998a), most likely in the absence of MT induction. While these data do not directly point to TNF- α , it strongly supports a role for Kupffer cells and, thus, suggests proinflammatory cytokines as mediators. More specifically, pretreatment of rats with anti-TNF- α

antibodies offers mild protection against Cd toxicity (Kayama *et al.*, 1995). Yamano *et al.* (2000) recently published a report that contrasted these data in demonstrating no induction of TNF- α upon Cd treatment. Rather, they showed that chemokines may be more important in the manifestation of Cd-induced hepatotoxicity.

In this study, we utilized the TNF-KO mouse to more conclusively determine whether TNF- α is involved in Cd-induced hepatotoxicity. Two separate time points were chosen to examine Cd-induced hepatic injury. Initially, the 9-h time point was chosen to correlate with earlier studies from this lab (Habeebu *et al.*, 1998). However, in this study, only moderate liver injury was observed 9 h after Cd treatment. This is most likely due to the relative resistance of the background strain (C57BL/6) for the TNF-KO mice to Cd-induced liver injury. Previous studies in this lab used a relatively more sensitive strain of mice (129Sv/Ola). Therefore, hepatotoxicity was also examined at a later time point (16 h) in order to cause more extensive toxicity and to possibly reveal whether there is a difference between WT and TNF-KO mice in susceptibility to

ACUTE CADMIUM HEPATOTOXICITY AND TNF- α

Cd-induced hepatotoxicity. Cadmium caused more hepatotoxicity after 16 than 9 h but still did not result in a significant shift in the dose response in TNF-KO in comparison to the WT mice. Therefore, it can be concluded from these data that TNF- α does not appear to be required for Cd-induced hepatotoxicity.

Previous experiments indicate that apoptosis may contribute to the hepatotoxicity of Cd (Habeebu *et al.*, 1998). In that study, apoptosis was observed maximally from 9 to 14 h in 129Sv/Ola mice. In the current study, Cd treatment did not cause an increase in apoptosis in either WT or TNF-KO mice (background C57BL/6) at any dose or time. These results may be interpreted as indicative of a strain difference in apoptosis in Cd-induced hepatotoxicity and require further investigation.

In summary, the present data demonstrate that, in contrast to LPS/Gln-induced hepatotoxicity, TNF- α does not appear to play a role in Cd-induced hepatotoxicity. These data do not exclude the involvement of other inducible proinflammatory cytokines and chemokines from the mechanism of cadmium-induced hepatotoxicity nor do they exclude a compensatory role of other proinflammatory cytokines in the TNF-KO mice after Cd administration. However, from these data, it can be concluded that, while TNF- α is clearly involved in LPS/Gln-induced hepatotoxicity, TNF- α does not appear to be important in Cd-induced hepatotoxicity.

ACKNOWLEDGMENTS

The authors thank Jeremy Affolter for his technical assistance. Additionally, the authors thank Dr. Michael Marino for generously providing WT and TNF-KO mice breeder pairs.

REFERENCES

- Adamson, G. M., and Billings, R. E. (1992). Tumor necrosis factor induced oxidative stress in isolated mouse hepatocytes. *Arch. Biochem. Biophys.* **294**, 223–229.
- Bagchi, D., Bagchi, M., Hassoun, E. A., and Stohs, S. J. (1996). Cadmium-induced excretion of urinary lipid metabolites, DNA damage, glutathione depletion, and hepatic lipid peroxidation in Sprague-Dawley rats. *Biol. Trace Elem. Res.* **52**, 143–154.
- Blazka, M. E., Elwell, M. R., Holladay, S. D., Wilson, R. E., and Luster, M. I. (1996). Histopathology of acetaminophen-induced liver changes: Role of interleukin 1 alpha and tumor necrosis factor alpha. *Toxicol. Pathol.* **24**, 181–189.
- Blazka, M. E., Wilmer, J. L., Holladay, S. D., Wilson, R. E., and Luster, M. I. (1995). Role of proinflammatory cytokines in acetaminophen hepatotoxicity. *Toxicol. Appl. Pharmacol.* **133**, 43–52.
- Boess, F., Bopst, M., Althaus, R., Polsky, S., Cohen, S. D., Eugster, H. P., and Boelsterli, U. A. (1998). Acetaminophen hepatotoxicity in tumor necrosis factor/lymphotoxin-alpha gene knockout mice. *Hepatology* **27**, 1021–1029.
- Bopst, M., Haas, C., Car, B., and Eugster, H. P. (1998). The combined inactivation of tumor necrosis factor and interleukin-6 prevents induction of the major acute phase proteins by endotoxin. *Eur. J. Immunol.* **28**, 4130–4137.
- Czaja, M. J., Flanders, K. C., Biempica, L., Klein, C., Zern, M. A., and Weiner, F. R. (1989). Expression of tumor necrosis factor-alpha and transforming growth factor-beta 1 in acute liver injury. *Growth Factors* **1**, 219–226.
- DeCicco, L. A., Rikans, L. E., Tutor, C. G., and Hornbrook, K. R. (1998). Serum and liver concentrations of tumor necrosis factor alpha and interleukin-1beta following administration of carbon tetrachloride to male rats. *Toxicol. Lett.* **98**, 115–121.
- Diamond, E. M., and Kench, J. E. (1974). Effects of cadmium on the respiration of rat liver mitochondria. *Environ. Physiol. Biochem.* **4**, 280–283.
- Dudley, R. E., Svoboda, D. J., and Klaassen, C. D. (1982). Acute exposure to cadmium causes severe liver injury in rats. *Toxicol. Appl. Pharmacol.* **65**, 302–313.
- Edwards, M. J., Keller, B. J., Kauffman, F. C., and Thurman, R. G. (1993). The involvement of Kupffer cells in carbon tetrachloride toxicity. *Toxicol. Appl. Pharmacol.* **119**, 275–279.
- Essani, N. A., Fisher, M. A., Farhood, A., Manning, A. M., Smith, C. W., and Jaeschke, H. (1995). Cytokine-induced upregulation of hepatic intercellular adhesion molecule-1 messenger RNA expression and its role in the pathophysiology of murine endotoxin shock and acute liver failure. *Hepatology* **21**, 1632–1639.
- Gillardot, F., Bottiger, B., Schmitz, B., Zimmermann, M., and Hossman, K. A. (1997). Activation of CPP-32 protease in hippocampal neurons following ischemia and epilepsy. *Brain Res. Mol. Brain Res.* **50**, 16–22.
- Habeebu, S. S., Liu, J., and Klaassen, C. D. (1998). Cadmium-induced apoptosis in mouse liver. *Toxicol. Appl. Pharmacol.* **149**, 203–209.
- Harstad, E. B., Hartley, D. P., Kolaja, K. L., and Klaassen, C. D. (1999). Activation of caspase-3 following cadmium chloride treatment in mice. *Toxicol. Sci.* **48**, 1674. [Abstract]
- Harvey, M. J., and Klaassen, C. D. (1983). Interaction of metals and carbon tetrachloride on lipid peroxidation and hepatotoxicity. *Toxicol. Appl. Pharmacol.* **71**, 316–322.
- Imuro, Y., Yamamoto, M., Kohno, H., Itakura, J., Fujii, H., and Matsumoto, Y. (1994). Blockade of liver macrophages by gadolinium chloride reduces lethality in endotoxemic rats: Analysis of mechanisms of lethality in endotoxemia. *J. Leukocyte Biol.* **55**, 723–728.
- Jaeschke, H., Smith, C. W., Clemens, M. G., Ganey, P. E., and Roth, R. A. (1996). Mechanisms of inflammatory liver injury: Adhesion molecules and cytotoxicity of neutrophils. *Toxicol. Appl. Pharmacol.* **139**, 213–226.
- Kayama, F., Yoshida, T., Elwell, M. R., and Luster, M. I. (1995). Role of tumor necrosis factor-alpha in cadmium-induced hepatotoxicity. *Toxicol. Appl. Pharmacol.* **131**, 224–234.
- Laskin, D. L. (1996). Sinusoidal lining cells and hepatotoxicity. *Toxicol. Pathol.* **24**, 112–118.
- Laskin, D. L., Gardner, C. R., Price, V. F., and Jollow, D. J. (1995). Modulation of macrophage functioning abrogates the acute hepatotoxicity of acetaminophen. *Hepatology* **21**, 1045–1050.
- Marino, M. W., Dunn, A., Grail, D., Inglesse, M., Noguchi, Y., Richards, E., Jungbluth, A., Wada, H., Moore, M., Williamson, B., Basu, S., and Old, L. J. (1997). Characterization of tumor necrosis factor-deficient mice. *Proc. Natl. Acad. Sci. USA* **94**, 8093–8098.
- Michael, S. L., Pumford, N. R., Mayeux, P. R., Niesman, M. R., and Hinson, J. A. (1999). Pretreatment of mice with macrophage inactivators decreases acetaminophen hepatotoxicity and the formation of reactive oxygen and nitrogen species. *Hepatology* **30**, 186–195.
- Morio, L. A., Chiu, H., Sprowles, K. A., Zhou, P., Heck, D. E., Gordon, M. K., and Laskin, D. L. (2001). Distinct roles of tumor necrosis factor- α and nitric oxide in acute liver injury induced by carbon tetrachloride in mice. *Toxicol. Appl. Pharmacol.* **172**, 44–51.
- Rikans, L. E., DeCicco, L. A., Hornbrook, K. R., and Yamano, T. (1999).

HARSTAD AND KLAASSEN

- Effect of age and carbon tetrachloride on cytokine concentrations in rat liver. *Mech. Ageing Dev.* **108**, 173-182.
- Roland, C. R., Naziruddin, B., Mohanakumar, T., and Flye, M. W. (1996). Gadolinium chloride inhibits Kupffer cell nitric oxide synthase (iNOS) induction. *J. Leukocyte Biol.* **60**, 487-492.
- Sauer, J. M., Waalkes, M. P., Hooser, S. B., Kuester, R. K., McQueen, C. A., and Sipes, I. G. (1997). Suppression of Kupffer cell function prevents cadmium induced hepatocellular necrosis in the male Sprague-Dawley rat. *Toxicology* **121**, 155-164.
- Stacey, N. H., Cantilena, L. R., and Klaassen, C. D. (1980). Cadmium toxicity and lipid peroxidation in isolated rat hepatocytes. *Toxicol. Appl. Pharmacol.* **53**, 470-480.
- Taniguchi, T., Takata, M., Ikeda, A., Momotani, E., and Sekikawa, K. (1997). Failure of germinal center formation and impairment of response to endotoxin in tumor necrosis factor alpha-deficient mice. *Lab. Invest.* **77**, 647-658.
- Tartaglia, L. A., Ayres, T. M., Wong, G. H., and Goeddel, D. V. (1993). A novel domain within the 55 kd TNF receptor signals cell death. *Cell* **74**, 845-853.
- Witthaut, R., Farhood, A., Smith, C. W., and Jaeschke, H. (1994). Complement and tumor necrosis factor-alpha contribute to Mac-1 (CD11b/CD18) up-regulation and systemic neutrophil activation during endotoxemia in vivo. *J. Leukocyte Biol.* **55**, 105-111.
- Yamano, T., DeCicco, L. A., and Rikans, L. E. (2000). Attenuation of cadmium-induced liver injury in senescent male fischer 344 rats: Role of Kupffer cells and inflammatory cytokines. *Toxicol. Appl. Pharmacol.* **162**, 68-75.
- Yamano, T., Shimizu, M., and Noda, T. (1998a). Age-related change in cadmium-induced hepatotoxicity in Wistar rats: Role of Kupffer cells and neutrophils. *Toxicol. Appl. Pharmacol.* **151**, 9-15.
- Yamano, T., Shimizu, M., and Noda, T. (1998b). Comparative effects of repeated administration of cadmium on kidney, spleen, thymus, and bone marrow in 2-, 4-, and 8-month-old male Wistar rats. *Toxicol. Sci.* **46**, 393-402.

Gadolinium Chloride Pretreatment Prevents Cadmium Chloride-Induced Liver Damage in Both Wild-Type and MT-Null Mice¹

Eric B. Harstad² and Curtis D. Klaassen

Department of Pharmacology, Toxicology, and Therapeutics, University of Kansas Medical Center, Kansas City, Kansas 66160-7140

Received November 5, 2001; accepted February 19, 2002

Gadolinium Chloride Pretreatment Prevents Cadmium Chloride-Induced Liver Damage in Both Wild-Type and MT-Null Mice. Harstad, E. B., and Klaassen, C. D. (2002). *Toxicol. Appl. Pharmacol.* 180, 178–185.

The heavy metal cadmium (Cd) causes hepatotoxicity upon acute administration. Kupffer cells, the resident macrophages of the liver, have been suggested to play a role in Cd-induced hepatotoxicity. Gadolinium chloride (GdCl₃) may prevent Cd-induced hepatotoxicity by suppressing Kupffer cells. However, GdCl₃ also induces the Cd-binding protein, metallothionein (MT). Therefore, this study was conducted to determine whether GdCl₃ prevents Cd-induced hepatotoxicity via the induction of MT. Hepatic MT and Kupffer cell counts were analyzed 24 h after wild-type (WT) mice were administered saline or 10, 30, or 60 mg GdCl₃/kg. GdCl₃ induced MT in a dose-dependent manner without affecting nonprotein sulfhydryl content. All examined doses of GdCl₃ were effective at eliminating Kupffer cells from the liver. To examine the hepatoprotective effects of GdCl₃, WT and MT-null mice were pretreated with saline or 10, 30, or 60 mg GdCl₃ 24 h prior to a hepatotoxic dose of Cd (2.5 mg Cd/kg). Blood and livers were removed 16 h later and analyzed for hepatotoxicity as well as MT, Cd, and Kupffer cell content. Hepatotoxicity was alleviated in both WT and MT-null mice that were pretreated with 30 or 60 mg GdCl₃/kg, indicating that MT induction is not required for the hepatoprotective effects of GdCl₃. Hepatic Cd content was not decreased by GdCl₃, demonstrating that GdCl₃ does not negatively affect Cd distribution to the liver. Kupffer cells were depleted at all three doses of GdCl₃, whereas hepatoprotection was only observed at doses of 30 and 60 mg GdCl₃/kg. This does not rule out Kupffer cells in the mechanism of Cd-induced hepatotoxicity, but it does suggest that GdCl₃ exerts hepatoprotective effects on the liver aside from depleting Kupffer cells. In summary, these data substantially rule out MT induction and decrease the importance of Kupffer cells as mechanisms of GdCl₃-induced protection from Cd-induced hepatotoxicity. © 2002 Elsevier Science (USA)

Key Words: cadmium; liver; hepatotoxicity; necrosis; gadolinium chloride; metallothionein; MT-null.

The heavy metal cadmium (Cd) is an industrial and environmental pollutant. It is toxic to several tissues, most notably

causing hepatotoxicity upon acute administration and nephrotoxicity upon chronic exposure. Histological evaluation of liver injury reveals that acute toxicity is comprised of hepatocellular swelling, sinusoidal congestion, pyknosis, and karyorrhexis (Dudley *et al.*, 1982). In a time-course study on Cd-induced hepatotoxicity, early cellular changes occur in the rough endoplasmic reticulum and nucleus (Dudley *et al.*, 1984). Later alterations include swelling of mitochondria and endoplasmic reticulum, loss of ribosomes, and appearance of fibrillar material within the cytoplasm. These cellular changes may result in both apoptosis and necrosis (Habeebu *et al.*, 1998).

Kupffer cells, the resident macrophages of the liver, have been suggested to play a role in Cd-induced hepatotoxicity. Kupffer cell activation by Cd was first noted by the identification of cytoplasmic vacuolization (Hoffmann *et al.*, 1975) and increased colloidal carbon clearance, which is indicative of increased phagocytic activity (Sauer *et al.*, 1997; Yamano *et al.*, 2000). Elimination of Kupffer cells with gadolinium chloride (GdCl₃) alleviated Cd-induced hepatotoxicity in rats (Sauer *et al.*, 1997; Yamano *et al.*, 2000) but failed to protect cultured hepatocytes (Badger *et al.*, 1997). Several reports have examined increases in cytokine expression in response to Cd administration. Tumor necrosis factor- α (TNF- α) is weakly induced by Cd both *in vitro* (Szuster-Ciesielska *et al.*, 2000; Marth *et al.*, 2000) and *in vivo* (Kayama *et al.*, 1995b). Pretreatment with antibodies against TNF- α abrogated Cd-induced expression of acute phase proteins but did not decrease Cd-induced hepatotoxicity (Kayama *et al.*, 1995b). In contrast, Yamano *et al.* (2000) determined that TNF- α was not increased within 24 h of Cd administration. Other cytokines, including interleukin-1 α (IL-1 α), interleukin-1 β (IL-1 β), interferon- γ (IFN- γ), and interleukin-6 (IL-6) (Kayama *et al.*, 1995a) are also increased after Cd treatment, although their relevance is less well defined (Kayama *et al.*, 1995b; Liu *et al.*, 1999; Yamano *et al.*, 2000; Marth *et al.*, 2000). Taken together, these data strongly suggest a role for Kupffer cells in Cd-induced hepatotoxicity.

Glutathione (GSH), the primary cellular nonprotein thiol, plays a role in the detoxication of Cd, although the exact role GSH plays in Cd-induced hepatotoxicity is not accurately defined. Depletion of GSH with phorone, diethyl maleate, or

¹ This study was supported by NIH Grant ES-01142.

² This author was supported by NIH Grant ES-07079.


GdCl₃ PREVENTS CdCl₂-INDUCED HEPATOTOXICITY

FIG. 1. Photomicrographs of wild-type mouse livers 24 h after GdCl₃ administration. Liver sections were prepared as described under Materials and Methods and visualized with either routine H & E staining (A and B, 200 \times) or immunohistochemical visualization of Kupffer cells (C and D, 400 \times). Photomicrographs are representative of mouse liver 24 h after administration of saline (A and C) or 60 mg GdCl₃ (B and D). Brown-staining cells were positive for the F4/80 antigen and were counted as Kupffer cells (C, inset, 1000 \times , thin arrow). Other cell types represented are endothelial cells (C, inset, arrowhead) and parenchymal cells (C, inset, thick arrow).

buthionine sulfoximine increases sensitivity to both Cd-induced lethality and hepatotoxicity (Dudley and Klaassen, 1984; Singhal *et al.*, 1987). Glutathione is also critical for the biliary elimination of Cd from the liver (Dijkstra *et al.*, 1996; Sugawara *et al.*, 1996). However, the protective effect may also be due to the reduction of Cd-induced oxidative stress by GSH (Shaikh *et al.*, 1999). Therefore, GSH plays an important and perhaps multifunctional role in the protection against Cd-induced liver injury.

Metallothioneins (MT) are small, cysteine-rich proteins that bind Cd with high affinity (Klaassen *et al.*, 1999). Pretreatment with low doses of zinc or Cd provides protection against subsequent hepatotoxic doses of Cd (Goering and Klaassen, 1984a,b). This protection appears to be due to induction of MT and subsequent redistribution of Cd to the cytoplasm, where it is sequestered by MT and thus detoxified (Goering and Klaassen, 1983). MT-null mice are not protected from subsequent administration of Cd, thus confirming the requirement of MT in the mechanism of this protective effect (Liu *et al.*, 1996).

GdCl₃ has also been shown to induce MT, although the significance of this induction is unknown (Sauer *et al.*, 1997).

These observations demonstrate that GdCl₃ exerts a variety of effects on the cells within the liver, including depletion of Kupffer cells as well as induction of MT. It is unknown which of these effects protects against Cd-induced hepatotoxicity. Therefore, this study was designed to determine whether GdCl₃ prevents Cd-induced hepatotoxicity by depletion of Kupffer cells or induction of MT.

MATERIALS AND METHODS

Animals. Wild-type (WT) (129SvIm/J, Jackson Labs, Bar Harbor, ME) and mutant mice with disrupted MT-I and MT-II genes (MT-null) (Masters *et al.*, 1994) were used throughout the study. Mice were housed in an AAALAC-certified facility at 70 \pm 2 $^{\circ}$ F with a 12-h light/dark cycle and were fed laboratory mouse chow (Purina, St. Louis, MO) and water *ad libitum*.

Chemical treatment of mice. To determine the hepatic effects of GdCl₃, male WT mice (8 weeks old, approximately 25 g, $n = 5$) were administered saline or various doses of 10, 30, or 60 mg GdCl₃ into the tail vein in a volume

of 10 ml/kg or approximately 250 μ l per mouse. Blood and liver were collected 24 h after dosing. Blood was processed for serum enzyme activity analyses. Livers were removed and a portion of the left lobe of each liver was fixed in 10% neutral buffered formalin. After 24 h in formalin, the fixed liver samples were switched to 70% ethanol for storage. The remainder of the liver was snap frozen in liquid nitrogen and stored at -80°C . To examine the hepatoprotective effects of GdCl_3 , male and female WT and MT-null mice (8 weeks old, approximately 25 g, $n = 7-8$) were injected with saline or 10, 30, or 60 mg GdCl_3 iv 24 h prior to a hepatotoxic dose of CdCl_2 (2.5 mg Cd/kg iv). Sixteen hours after Cd administration, blood and liver were collected and processed as described for the first experiment.

Serum enzyme activity assays. Biochemical evaluation of liver injury was performed by quantifying serum activities of alanine aminotransferase (ALT) and sorbitol dehydrogenase (SDH) spectrophotometrically, using Sigma test kits (Sigma Chemical Company, St. Louis, MO) according to the manufacturer's instructions.

Histopathology. Liver samples were taken consistently from the left lobe of the liver and processed by standard histological techniques. Briefly, samples were fixed in formalin for 24 h and then switched to ethanol for storage. Liver sections were then processed routinely and embedded in paraffin blocks. Slides were prepared (5 μm) and stained with hematoxylin and eosin. The slides were blinded and analyzed by light microscopy for liver injury. The grade of liver injury was analyzed semiquantitatively with six scores of severity per liver section: 0 = none; 1 = minimal (>2 foci of single cell necrosis); 2 = mild (at least 5 areas of focal necrosis); 3 = moderate (at least five foci of zonal necrosis); 4 = severe (lobular damage, with many viable lobules); 5 = global (severe lobular damage, few areas of viability).

Immunohistochemistry. Liver samples were fixed and embedded as described above. Liver sections were cut (5 μm) and mounted on Superfrost Plus slides (Fisher Scientific, Fair Lawn, NJ). Briefly, slides were incubated with proteinase K (20 $\mu\text{g/ml}$) for 15 min at room temperature for antigen retrieval. Endogenous peroxidase activity was quenched by treatment of the slides with 3% H_2O_2 for 10 min. Treating the slides for 30 min with 5% normal rabbit serum in PBS blocked nonspecific binding. Kupffer cells were then detected with a primary antibody to the F4/80 macrophage surface antigen (MCPAP497, Serotec, Inc., Raleigh, NC). A biotinylated anti-rat secondary antibody (BA-4001, Vector Laboratories, Burlingame, CA) specifically identified the primary antibody. The liver sections were then incubated with avidin and biotinylated horseradish peroxidase (HRP) and identified using the HRP substrate diaminobenzidine and counterstained with hematoxylin. Brown-staining cells with appropriate nuclear morphology and sinusoidal location were counted as Kupffer cells. Kupffer cells were quantified as the average number of F4/80-positive cells per field (400 \times magnification) from at least 20 randomly selected fields per section.

Metallothionein assay. Liver samples were prepared by homogenizing 1:3 (w/v) in 10 mM Tris-HCl buffer (pH 7.4), centrifuged at 13,000g, and the supernatant fractions were retained. Protein concentration in supernatants was determined by the bicinchoninic acid method using a BCA kit (Pierce, Rockford, IL). Supernatants were assayed and hepatic MT content was determined using the Cd-hemoglobin method as previously described (Eaton and Toal, 1982).

Hepatic nonprotein sulfhydryl (NPSH) assay. Hepatic NPSH was quantified using a method described previously (Ellman, 1959). Briefly, 100-mg liver samples were homogenized using a Kinematica Polytron homogenizer (Littau, Switzerland) in 5% TCA/EDTA and separated by centrifugation at 13,000g, and the supernatants were retained for NPSH analysis. Each reaction contained 176 μl of 0.1 M PO_4^{3-} buffer (pH 8), 16 μl of supernatant, and 8 μl of 5 mM 5,5'-dithio-bis(2-nitrobenzoic acid). Absorbance was quantified using a Biotek microtiter plate spectrophotometer (Winooski, VT) at 405 nm (analytical wavelength) and 690 nm (reference wavelength) and compared with a standard curve of known GSH concentrations.

Hepatic Cd assay. Liver samples (~ 1.0 g) were digested 1:3 (w/v) in concentrated HNO_3 at 100°C for 60 min. The digested samples were assayed

for Cd content using a Perkin-Elmer atomic absorption spectrophotometer (Norwalk, CT) with an analytical wavelength of 228.8 nm. Sample Cd content was determined by comparing absorbance values to a standard curve of Cd solutions.

Statistics. One-way ANOVA was used to analyze data from the GdCl_3 dose-response study. Two-way ANOVA followed with a Duncan's multiple range test was used for all data from Cd-treated groups. For all experiments, the acceptable level of significance was chosen to be $p < 0.05$.

RESULTS

Assessment of hepatic effects of GdCl_3 . Doses of 10, 30, and 60 mg GdCl_3/kg caused no overt toxicity in WT mice. Serum markers of hepatotoxicity, ALT and SDH were not increased or decreased after administration of any dose of GdCl_3 studied (data not shown). In addition to serum markers of hepatotoxicity, livers were examined for histological evidence of liver injury. After saline treatment, liver was histologically normal except for some mild hydropic change (Fig. 1A, 200 \times). After 60 mg GdCl_3 , liver morphology was histologically indistinguishable from saline treatment (Fig. 1B, 200 \times). Kupffer cells were identified immunohistochemically and counterstained with hematoxylin (Figs. 1C and 1D). Nonparenchymal cells that were positive for the F4/80 antigen (brown-staining cells) and displayed appropriate nuclear morphology (inset, 1000 \times) were counted as Kupffer cells. Nuclear morphology of Kupffer cells can be contrasted with endothelial (inset, arrowhead) and parenchymal cell nuclei (inset, curved arrow). Kupffer cells were quantified as the number of F4/80-positive cells per high-power field (400 \times). Kupffer cells were abundant after saline administration (Fig. 1C, 400 \times). Similar analysis of liver sections after 60 mg/kg identified no F4/80-positive cells (Fig. 1D, 400 \times).

Quantitation of Kupffer cells after GdCl_3 administration. Kupffer cells were quantified as the number of F4/80-positive cells per field from at least 20 randomly selected fields per section. Kupffer cells were identified only in saline-treated control rats (Fig. 2A). At all doses of GdCl_3 , no cells were positive for the F4/80 antigen. Nonparenchymal cells were also qualitatively analyzed by examining nuclear morphology to determine whether Kupffer cells were still present but not antigenic. However, no Kupffer cells could be positively identified morphologically.

Effect of GdCl_3 administration on hepatic MT and NPSH contents. Livers were processed and analyzed for metallothionein by the cadmium-hemoglobin radioassay (Eaton and Toal, 1982). Hepatic metallothionein increased in a dose-dependent manner after GdCl_3 administration (Fig. 2B). In saline-treated mice, MT was approximately 3 $\mu\text{g/g}$ liver. In GdCl_3 -treated mice, MT increased to 8, 18, and 42 $\mu\text{g/g}$ liver at doses of 10, 30, and 60 mg GdCl_3/kg , respectively. These represent 3-, 6-, and 15-fold increases in hepatic MT concentration, respectively. In addition, liver was processed and analyzed for NPSH by the DTNB method (Ellman, 1959). NPSH


GdCl₃ PREVENTS CdCl₂-INDUCED HEPATOTOXICITY

FIG. 2. Quantification of Kupffer cells, MT, and NPSH content 24 h after GdCl₃ administration. Wild-type mice (129SvIm/J) were administered saline or various doses of GdCl₃ (10, 30, or 60 mg GdCl₃/kg). Livers were collected 24 h after dosing. Livers were processed and analyzed for (A) Kupffer cells, (B) MT, and (C) NPSH content as described under Materials and Methods. (A) *Livers from GdCl₃-treated mice contained significantly fewer Kupffer cells than livers from saline-treated control mice ($p < 0.05$). (B) *Livers from GdCl₃-treated mice contained significantly greater MT than livers from saline-treated control mice ($p < 0.05$). Data are presented as means \pm SE ($n = 5$).

concentrations were not significantly changed with any dose of GdCl₃ (Fig. 2C).

Histopathology of Cd-induced hepatotoxicity after saline or GdCl₃ pretreatment in WT and MT-null mice. Liver sections were stained with H & E and scored semiquantitatively for liver injury as described. Cadmium administration resulted in congestion and multifocal hepatic necrosis 16 h after 2.5 mg Cd in WT mice with saline pretreatment (Fig. 3A). Cadmium caused more severe hepatotoxicity 16 h after 2.5 mg Cd in MT-null mice with saline pretreatment (Fig. 3B). In MT-null mice, necrosis was multifocal and consisted of larger lesions with more extensive congestion. Treatment with 60 mg GdCl₃

24 h prior to 2.5 mg Cd completely abrogated the cadmium-induced hepatotoxicity in WT mice (Fig. 3C). In all samples pretreated with 60 mg GdCl₃, there was an absence of both congestion and necrosis after Cd administration. A notable histological change is consistent with hydropic change that was not confined to any specific zone of the liver. This hydropic change was not considered deleterious because hydropic change is reversible and there were no increases in serum indices of toxicity. Treatment with 60 mg GdCl₃ 24 h prior to 2.5 mg Cd also prevented hepatotoxicity in MT-null mice (Fig. 3D). Pretreatment with GdCl₃ protected MT-null mice in a manner that was histologically indistinguishable from the same treatment in WT mice.

Serum indices of Cd-induced hepatotoxicity after GdCl₃ pretreatment in WT and MT-null mice. After saline pretreatment, Cd caused marked increases in serum indices of hepatotoxicity in WT mice (Figs. 4A and 4B). Pretreatment with 10 mg GdCl₃/kg did not significantly decrease ALT or SDH activities in Cd-treated WT mice. However, pretreatment with 30 and 60 mg GdCl₃/kg significantly decreased ALT (95 and 98%, respectively) and SDH (93 and 97%, respectively) activities. In saline-pretreated MT-null mice, Cd administration also caused extensive hepatotoxicity. Pretreatment with 10 mg GdCl₃/kg tended to decrease ALT activity (35%) and SDH activity (55%), although these decreases were not significant; whereas pretreatment with 30 mg GdCl₃/kg resulted in significantly lower serum ALT (90%) and SDH (85%) activities. Pretreatment with 60 mg GdCl₃/kg completely abrogated Cd-induced liver injury in both WT and MT-null mice.

Histological analysis of Cd-induced liver injury and Kupffer cell depletion after GdCl₃ administration in WT and MT-null mice. The histological grade of liver injury paralleled the serum markers of hepatotoxicity (Fig. 5A). F4/80-positive Kupffer cells only were noted in liver sections from saline-pretreated mice (Fig. 5B). Compared to WT mice that were administered saline only, saline-pretreated WT and MT-null mice had significantly fewer Kupffer cells (approximately seven per field). In both WT and MT-null mice that were pretreated with all doses of GdCl₃ prior to Cd administration, very few F4/80-positive cells were noted.

Hepatic Cd content after GdCl₃ and Cd administration. Liver samples were also analyzed for hepatic Cd content (Fig. 6A). Livers from WT mice pretreated with saline, 10, 30, and 60 mg GdCl₃/kg prior to 2.5 mg Cd/kg contained 14, 17, 17, and 24 μ g Cd/g liver, respectively. Only WT mice pretreated with 60 mg GdCl₃/kg prior to 2.5 mg Cd/kg had higher hepatic Cd content than WT mice that were pretreated with saline prior to 2.5 mg Cd/kg. Livers from MT-null mice pretreated with saline, 10, 30, and 60 mg GdCl₃/kg prior to 2.5 mg Cd/kg contained 11, 10, 14, and 11 μ g Cd/g liver, respectively. GdCl₃ pretreatment did not significantly alter Cd accumulation in livers from MT-null mice. Overall, hepatic Cd content was




FIG. 3. Photomicrographs showing Cd-induced hepatotoxicity after $GdCl_3$ pretreatment in WT and MT-null mice. Wild-type and MT-null mice were administered saline or various doses of $GdCl_3$ (10, 30, or 60 mg $GdCl_3/kg$) 24 h prior to a hepatotoxic dose of Cd (2.5 mg Cd/kg). Livers were collected 16 h after Cd administration, processed routinely, and stained with hematoxylin and eosin. Photomicrographs (200 \times) are representative of mouse liver from wild-type (A and B) and MT-null (C and D) mice 24 h after administration of Cd after pretreatment with saline (A and C) or 60 mg $GdCl_3$ (B and D).

slightly higher in WT mice than MT-null mice, although this difference was significant only at 10 and 60 mg $GdCl_3/kg$.

Hepatic MT content after $GdCl_3$ and $CdCl_2$ administration. Hepatic metallothionein was induced in all Cd-treated WT mice (Fig. 6B). Livers from WT mice pretreated with 10 or 30 mg $GdCl_3/kg$ had MT concentrations (91 and 100 $\mu g/g$ liver, respectively) that were similar to saline-pretreated controls (100 $\mu g/g$ liver), whereas livers from WT mice pretreated with 60 mg $GdCl_3/kg$ prior to Cd administration had significantly more MT (115 $\mu g/g$ liver) than mice administered saline plus 2.5 mg Cd/kg (91 $\mu g/g$ liver). In MT-null mice, hepatic MT content was not increased and remained unchanged in all treatment groups.

DISCUSSION

Mounting evidence supports the theory that Kupffer cells play a role in Cd-induced hepatotoxicity. Cadmium administration results in an increase in colloidal carbon clearance,

indicative of a general increase in phagocytic activity (Sauer *et al.*, 1997; Yamano *et al.*, 2000). Elimination of Kupffer cells with $GdCl_3$ alleviates Cd-induced hepatotoxicity in rats (Sauer *et al.*, 1997; Yamano *et al.*, 1998) but does not protect cultured hepatocytes (Badger *et al.*, 1997). However, the effects of $GdCl_3$ on the liver are not limited to Kupffer cell depletion. For example, $GdCl_3$ has been shown to suppress superoxide production, interfere with inducible nitric oxide synthase expression, and induce MT (Iimuro *et al.*, 1994; Roland *et al.*, 1996; Sauer *et al.*, 1997).

Metallothionein is a low-molecular-weight, cysteine-rich protein that has been shown to be highly inducible and to protect the liver from Cd-induced hepatotoxicity. Cd and other metals, such as zinc, also induce MT, thus conferring resistance to Cd-induced hepatotoxicity (Goering and Klaassen, 1984a,b). This protective effect is absent in MT-null mice, confirming the requirement of MT for protection (Liu *et al.*, 1996). Gadolinium is a pleiotropic metal that, in addition to eliminating Kupffer cells from the liver, induces MT (Sauer *et*


GdCl₃ PREVENTS CdCl₂-INDUCED HEPATOTOXICITY

FIG. 4. Assessment of Cd-induced hepatotoxicity after GdCl₃ pretreatment in WT and MT-null mice. Wild-type (solid bars) and MT-null mice (hatched bars) were administered saline or various doses of GdCl₃ (10, 30, or 60 mg GdCl₃) 24 h prior to a hepatotoxic dose of Cd (2.5 mg Cd/kg). Blood was collected 16 h after Cd administration. Hepatotoxicity was assessed by measuring serum indices of hepatotoxicity, namely (A) ALT and (B) SDH activities. *GdCl₃-treated mice exhibited significantly less hepatotoxicity than respective saline-treated control mice ($p < 0.05$). Data are presented as means \pm SE ($n = 7-8$).

al., 1997). It is therefore reasonable to hypothesize that GdCl₃ may increase hepatic MT and thus protect the liver from Cd-induced liver injury.

To determine the dose-dependent effects of GdCl₃ on hepatic MT, WT mice were administered various doses of GdCl₃. The highest dose of GdCl₃ used was 60 mg/kg because higher doses began to cause seizures and death. In these studies, the LD₅₀ was approximately 100 mg GdCl₃/kg. All mice survived doses up to 60 mg GdCl₃/kg. Twenty-four hours after GdCl₃ administration, livers were examined for toxicity, thiol alterations, and depletion of Kupffer cells. Both biochemical and histological examinations were utilized to assess liver injury. Administration of saline, 10, 30, or 60 mg/kg GdCl₃ iv did not cause liver injury as measured by quantifying serum indices of hepatotoxicity (data not shown). Livers from both saline- and GdCl₃-treated mice were histologically normal, except for some hydropic changes (Figs. 1A and 1B). Kupffer cells were identified immunohistochemically (Figs. 1C and 1D) and found to be eliminated from the liver at all doses of GdCl₃ (Fig. 2A). These data indicate that GdCl₃ is selectively toxic to Kupffer cells and does not appear to exert toxicity on other cell types of the liver.

Metallothionein and glutathione comprise the primary protein and nonprotein sulfhydryls that participate in hepatoprotection.

A wide variety of chemicals, including metals such as Cd and Zn, are known to induce MT. In this study, GdCl₃ also increased hepatic MT in a dose-dependent manner (Fig. 2B). Although GdCl₃ induced MT significantly, it was neither as potent or effective as Zn or Cd at inducing MT. However, the degree to which GdCl₃ induced MT is enough to protect liver from Cd-induced liver injury. Nonprotein sulfhydryls were also examined and were found to be unaffected by GdCl₃ treatment (Fig. 2C). Therefore, it appears that, although glutathione is not altered by GdCl₃, MT induction may account for the hepatoprotective effects of GdCl₃.

While it is not a novel finding that GdCl₃ is effective at depleting Kupffer cells from the liver, it is interesting that lower doses of GdCl₃ (10 mg GdCl₃) are required to deplete Kupffer cells than to induce MT (30 and 60 mg GdCl₃/kg) in WT mice. This dose-related difference can be exploited to discern the mechanism of GdCl₃-induced hepatoprotection. Therefore, Cd-induced hepatotoxicity was examined in mice after pretreatment with saline or GdCl₃. This experiment was conducted in both WT and MT-null mice to eliminate MT as an experimental variable, in addition to comparing the dose re-




FIG. 5. Histological analysis of Cd-induced liver injury and Kupffer cell depletion after GdCl₃ pretreatment. Wild-type (solid bars) and MT-null mice (hatched bars) were administered saline or various doses of GdCl₃ (10, 30, or 60 mg GdCl₃) 24 h prior to a hepatotoxic dose of Cd (2.5 mg Cd/kg). Livers were collected 16 h after Cd administration and prepared as described under Materials and Methods. (A) Liver sections were visualized with standard hematoxylin and eosin and then scored semiquantitatively for liver injury. (B) Kupffer cells were identified using a primary antibody to the F4/80 macrophage surface antigen (Serotec) and quantified as the number of F4/80-positive cells per high-power field (400 \times). *Livers from GdCl₃-treated mice contained significantly fewer Kupffer cells than livers from saline-treated control mice ($p < 0.05$). Data are presented as means \pm SE ($n = 7-8$).

HARSTAD AND KLAASSEN




FIG. 6. Assessment of hepatic Cd and MT content after GdCl₃ and CdCl₂ administration. WT (solid bars) and MT-null mice (hatched bars) were administered saline or various doses of GdCl₃ (10, 30, or 60 mg GdCl₃) 24 h prior to a hepatotoxic dose of Cd (5 mg Cd/kg). Livers were collected 16 h after Cd administration. Liver samples were prepared and analyzed for Cd (A) and MT (B) content as described under Materials and Methods. ^aLivers from GdCl₃-treated mice contained significantly greater Cd or MT than livers from saline-treated control mice ($p < 0.05$). ^bLivers from MT-null mice contained significantly less Cd or MT than livers from WT mice after similar chemical treatment ($p < 0.05$). Data are presented as means \pm SE ($n = 7-8$).

sponses of hepatotoxicity, MT induction, and Kupffer cell depletion. Wild-type and MT-null mice were pretreated with saline or the same doses of GdCl₃. Subsequently, all mice were administered a hepatotoxic dose of Cd (2.5 mg Cd/kg). Serum indices of hepatotoxicity (Figs. 4A and 4B) were confirmed by histological evaluation (Fig. 5A). After saline pretreatment, Cd caused extensive liver injury in both WT and MT-null mice. Pretreatment of WT and MT-null mice with 30 mg GdCl₃/kg 24 h prior to the hepatotoxic dose of Cd alleviated most of the Cd-induced hepatotoxicity. Administration of 60 mg GdCl₃/kg effectively abrogated Cd-induced hepatotoxicity in both WT and MT-null mice. These data demonstrate that, in WT mice, GdCl₃ induces MT in a dose-dependent manner that is consistent with hepatoprotection. Similarly, MT-null mice also benefit from the hepatoprotective effects of GdCl₃. Therefore, while GdCl₃ induces MT in WT mice, GdCl₃ does not prevent Cd-induced hepatotoxicity by inducing MT.

Kupffer cells were identified immunohistochemically and counted in an effort to compare the dose responses of Kupffer cell depletion and hepatoprotection. Similar numbers of Kupffer cells were present in WT and MT-null mice after saline pretreatment (Fig. 5B). Again, few Kupffer cells could be identified in GdCl₃-pretreated WT and MT-null mice. How-

ever, the dose-dependent nature of the decrease in Kupffer cells is not consistent with the dose-dependent protection from Cd-induced hepatotoxicity, suggesting that GdCl₃ exerts hepatoprotective effects aside from depleting Kupffer cells. The disparity between dose-response relationships of hepatoprotection and Kupffer cell depletion does not conclusively rule out the involvement of Kupffer cells in the mechanism of Cd-induced hepatotoxicity. However, this is strong evidence that there are other changes in the liver that may account for the hepatoprotective effects of GdCl₃.

Hepatic Cd content was also quantified to determine whether GdCl₃ prevents hepatotoxicity by decreasing the distribution of Cd to the liver (Fig. 6A). In WT mice, the hepatic Cd burden was not decreased with increasing doses of GdCl₃. Rather, hepatic Cd content was actually slightly increased (1.7-fold) at 60 mg GdCl₃/kg despite decreased toxicity. This is most likely due to sequestration of Cd by GdCl₃-induced presynthesized MT. From these data, it can be concluded that GdCl₃ pretreatment does not prevent Cd-induced liver injury by decreasing the absolute quantity of Cd in the liver.

Hepatic MT protein levels were examined in WT and MT-null mice to determine whether MT induction accounts for the increased hepatic Cd content as well as to confirm that MT-null mice are truly deficient in MT. Compared to saline-treated WT mice, hepatic MT was greatly increased after saline or GdCl₃ pretreatment with subsequent Cd administration (Fig. 6B). At the highest dose of GdCl₃ (60 mg/kg), MT was significantly higher than saline pretreatment, indicating that the MT induction by GdCl₃ occurs in addition to Cd-induced MT induction. This pattern is consistent with the increases in hepatic Cd, suggesting that the increases in Cd content are due to sequestration of Cd by presynthesized MT. Hepatic MT was absent in MT-null mice, confirming the genotype of the MT-null mice. This same pattern of hepatoprotection was achieved by pretreatment with subtoxic doses of Zn (Liu *et al.*, 1996). However, the current study contrasts previous studies in which Zn pretreatment did not protect against Cd-induced hepatotoxicity in MT-null mice. In this study, MT-null mice are also protected at the same doses of GdCl₃ that protect WT mice. Therefore, it can be concluded that MT does not account for the hepatoprotective effects of GdCl₃.

Taken together, these data suggest multiple mechanisms for GdCl₃-induced protection from Cd-induced liver injury. These data demonstrate that GdCl₃ is an effective inducer of MT at doses that are consistent with protection against Cd-induced liver injury. However, MT induction has a dispensable cytoprotective effect as seen in MT-null mice. Additionally, GdCl₃ does not alleviate hepatotoxicity by decreasing total Cd accumulation in the liver. Kupffer cells are effectively depleted at doses below those required to prevent Cd-induced liver injury. This does not rule out Kupffer cells in the mechanism of Cd-induced hepatotoxicity, but it does suggest that GdCl₃ exerts hepatoprotective effects in addition to depleting Kupffer cells. In summary, this study substantially rules out MT induc-

GdCl₃ PREVENTS CdCl₂-INDUCED HEPATOTOXICITY

tion in the mechanism of GdCl₃-induced protection from Cd-induced hepatotoxicity.

ACKNOWLEDGMENTS

The authors thank Yaping Liu and Dr. Angela Slitt for their technical assistance.

REFERENCES

- Badger, D. A., Kuester, R. K., Sauer, J. M., and Sipes, I. G. (1997). Gadolinium chloride reduces cytochrome P450: Relevance to chemical-induced hepatotoxicity. *Toxicology* **121**, 143–153.
- Dijkstra, M., Havinga, R., Vonk, R. J., and Kuipers, F. (1996). Bile secretion of cadmium, silver, zinc and copper in the rat: Involvement of various transport systems. *Life Sci.* **59**, 1237–1246.
- Dudley, R. E., and Klaassen, C. D. (1984a). Changes in hepatic glutathione concentration modify cadmium-induced hepatotoxicity. *Toxicol. Appl. Pharmacol.* **72**, 530–538.
- Dudley, R. E., Svoboda, D. J., and Klaassen, C. D. (1982). Acute exposure to cadmium causes severe liver injury in rats. *Toxicol. Appl. Pharmacol.* **65**, 302–313.
- Dudley, R. E., Svoboda, D. J., and Klaassen, C. D. (1984b). Time course of cadmium-induced ultrastructural changes in rat liver. *Toxicol. Appl. Pharmacol.* **76**, 150–160.
- Eaton, D. L., and Toal, B. F. (1982). Evaluation of the Cd/hemoglobin affinity assay for the rapid determination of metallothionein in biological tissues. *Toxicol. Appl. Pharmacol.* **66**, 134–142.
- Ellman, G. L. (1959). Tissue sulfhydryl groups. *Arch. Biochem. Biophys.* **82**, 70–77.
- Goering, P. L., and Klaassen, C. D. (1983). Altered subcellular distribution of cadmium following cadmium pretreatment: Possible mechanism of tolerance to cadmium-induced lethality. *Toxicol. Appl. Pharmacol.* **70**, 195–203.
- Goering, P. L., and Klaassen, C. D. (1984a). Tolerance to cadmium-induced hepatotoxicity following cadmium pretreatment. *Toxicol. Appl. Pharmacol.* **74**, 308–313.
- Goering, P. L., and Klaassen, C. D. (1984b). Zinc-induced tolerance to cadmium hepatotoxicity. *Toxicol. Appl. Pharmacol.* **74**, 299–307.
- Habeebu, S. S., Liu, J., and Klaassen, C. D. (1998). Cadmium-induced apoptosis in mouse liver. *Toxicol. Appl. Pharmacol.* **149**, 203–209.
- Hoffmann, E. O., Cook, J. A., di Luzio, N. R., and Coover, J. A. (1975). The effects of acute cadmium administration in the liver and kidney of the rat: Light and electron microscopic studies. *Lab Invest.* **32**, 655–664.
- Iimuro, Y., Yamamoto, M., Kohno, H., Itakura, J., Fujii, H., and Matsumoto, Y. (1994). Blockade of liver macrophages by gadolinium chloride reduces lethality in endotoxemic rats: Analysis of mechanisms of lethality in endotoxemia. *J. Leukocyte Biol.* **55**, 723–728.
- Kayama, F., Yoshida, T., Elwell, M. R., and Luster, M. I. (1995a). Cadmium-induced renal damage and proinflammatory cytokines: Possible role of IL-6 in tubular epithelial cell regeneration. *Toxicol. Appl. Pharmacol.* **134**, 26–34.
- Kayama, F., Yoshida, T., Elwell, M. R., and Luster, M. I. (1995b). Role of tumor necrosis factor- α in cadmium-induced hepatotoxicity. *Toxicol. Appl. Pharmacol.* **131**, 224–234.
- Klaassen, C. D., Liu, J., and Choudhuri, S. (1999). Metallothionein: An intracellular protein to protect against cadmium toxicity. *Annu. Rev. Pharmacol. Toxicol.* **39**, 267–294.
- Liu, J., Liu, Y., Habeebu, S. S., and Klaassen, C. D. (1999). Metallothionein-null mice are highly susceptible to the hematotoxic and immunotoxic effects of chronic CdCl₂ exposure. *Toxicol. Appl. Pharmacol.* **159**, 98–108.
- Liu, J., Liu, Y., Michalska, A. E., Choo, K. H., and Klaassen, C. D. (1996). Metallothionein plays less of a protective role in cadmium-metallothionein-induced nephrotoxicity than in cadmium chloride-induced hepatotoxicity. *J. Pharmacol. Exp. Ther.* **276**, 1216–1223.
- Marth, E., Barth, S., and Jelovec, S. (2000). Influence of cadmium on the immune system: Description of stimulating reactions. *Cent. Eur. J. Public Health* **8**, 40–44.
- Masters, B. A., Kelly, E. J., Quaife, C. J., Brinster, R. L., and Palmiter, R. D. (1994). Targeted disruption of metallothionein I and II genes increases sensitivity to cadmium. *Proc. Natl. Acad. Sci. USA* **91**, 584–588.
- Roland, C. R., Naziruddin, B., Mohanakumar, T., and Flye, M. W. (1996). Gadolinium chloride inhibits Kupffer cell nitric oxide synthase (iNOS) induction. *J. Leukocyte Biol.* **60**, 487–492.
- Sauer, J. M., Waalkes, M. P., Hooser, S. B., Kuester, R. K., McQueen, C. A., and Sipes, I. G. (1997). Suppression of Kupffer cell function prevents cadmium induced hepatocellular necrosis in the male Sprague-Dawley rat. *Toxicology* **121**, 155–164.
- Shaikh, Z. A., Vu, T. T., and Zaman, K. (1999). Oxidative stress as a mechanism of chronic cadmium-induced hepatotoxicity and renal toxicity and protection by antioxidants. *Toxicol. Appl. Pharmacol.* **154**, 256–263.
- Singhal, R. K., Anderson, M. E., and Meister, A. (1987). Glutathione, a first line of defense against cadmium toxicity. *FASEB J.* **1**, 220–223.
- Sugawara, N., Lai, Y. R., Arizono, K., and Ariyoshi, T. (1996). Biliary excretion of exogenous cadmium, and endogenous copper and zinc in the Eisai hyperbilirubinuric (EHB) rat with a near absence of biliary glutathione. *Toxicology* **112**, 87–94.
- Szuster-Ciesielska, A., Lokaj, I., and Kandefers-Szerszen, M. (2000). The influence of cadmium and zinc ions on the interferon and tumor necrosis factor production in bovine aorta endothelial cells. *Toxicology* **145**, 135–145.
- Yamano, T., DeCicco, L. A., and Rikans, L. E. (2000). Attenuation of cadmium-induced liver injury in senescent male Fischer 344 rats: Role of Kupffer cells and inflammatory cytokines. *Toxicol. Appl. Pharmacol.* **162**, 68–75.
- Yamano, T., Shimizu, M., and Noda, T. (1998). Age-related change in cadmium-induced hepatotoxicity in Wistar rats: Role of Kupffer cells and neutrophils. *Toxicol. Appl. Pharmacol.* **151**, 9–15.