

## Novel Mechanisms of Toxic Bile Salt-Induced Hepatocellular Apoptosis

Byung-Hoon Lee

College of Pharmacy, Wonkwang University

Cholestatic liver injury results from the accumulation of toxic bile salts within the liver. The aim of the present study was to understand the mechanism of bile salts-induced hepatocellular apoptosis in bile duct-ligated (BDL) rats, using Western blot and immunohistochemical analysis. The mechanisms of hepatoprotective effects of *Salvia miltiorrhiza* (SM) was also studied in this model. Apoptotic cell death was increased five-fold after three days of BDL, decreased over two weeks and remained constant thereafter. Total cellular Bax protein was increased three days after BDL and decreased over time thereafter. We observed the translocation of Bax to mitochondria and subsequent release of cytochrome *c*. Nuclear p53 increased three days after BDL, but most p53 was expressed in the cytoplasm after one week. The expression of c-Myc was inhibited by three days, but increased at later stages following BDL. Bcl-2 was increased over time in BDL rats. Fas expression was not changed and activation of caspase 8 did not occur. Fas immunoreactivity was exclusively observed in the cytoplasm of hepatocytes, indicating that Fas expressed in rat hepatocytes is predominantly a soluble form. Thus our data suggest that nuclear p53 regulates the expression of Bax in BDL rats. The translocation of Bax to mitochondria supports the hypothesis that this may lead to release of cytochrome *c* and transduce the signal for apoptotic death of hepatocytes by toxic bile salts. The lack of changes in Fas suggests that Fas may not be playing a role in apoptosis in this model.

*Salvia miltiorrhiza* BUNGE, a traditional chinese herbal medicine, has been commonly used to treat chronic hepatitis and liver fibrosis. However, the mechanism of action is not yet fully understood. The treatment of extract of SM to BDL rat reduced the number of TUNEL positive cells to the control level after 10 days of BDL. The expression of Bax was reduced with time by the treatment of SM. The treatment of SM accelerated the cytoplasmic sequestration of p53. The expression of Bcl-2 was increased, hence the Bax to Bcl-2 ratio was decreased significantly. These results suggest that SM may be used as a hepatoprotective agent in extrahepatic cholestasis model. (This research was supported by a grant (PF002105-01) from Plant Diversity Research Center of 21st Frontier Research Program funded by MOST and Spella Co.)

## *Salvia miltiorrhiza* Inhibits Biliary Obstruction-Induced Hepatocyte Apoptosis by Cytoplasmic Sequestration of p53

Seon-Hee Oh, Ji-Xing Nan,<sup>1</sup> Dong-Hwan Sohn, Youn-Chul Kim, and Byung-Hoon Lee<sup>2</sup>

College of Pharmacy and Medicinal Resources Research Center, Wonkwang University, Iksan, Jeonbuk 570-749 Korea

Received November 26, 2001; accepted January 10, 2002

*Salvia miltiorrhiza* Inhibits Biliary Obstruction-Induced Hepatocyte Apoptosis by Cytoplasmic Sequestration of p53. Oh, S.-H., Nan, J.-X., Sohn, D.-H., Kim, Y.-C., and Lee, B.-H. (2002). *Toxicol. Appl. Pharmacol.* 182, 27–33.

Cholestatic liver injury is caused by hepatocellular apoptosis induced by toxic bile salts. We have studied the effects of a traditional Chinese herbal medicine, *Salvia miltiorrhiza*, on apoptotic cell death in bile duct-ligated (BDL) rats. We also attempted to clarify the molecular mechanisms of the hepatoprotective effects of *S. miltiorrhiza* in this animal model. A water extract of *S. miltiorrhiza* (Sm-X; 200 mg/kg; po) was administered to BDL rats for 10 days. Rats were euthanized and apoptosis was detected in liver tissue by TUNEL staining. Western blot analysis and immunostaining for  $\alpha$ -smooth muscle actin ( $\alpha$ -SMA), Bax, Bcl-2, and p53 were performed. Results show that the treatment of BDL rats with Sm-X significantly improved the liver function parameters, although the expression of  $\alpha$ -SMA, a marker of hepatic stellate cell activation, was not affected. Treatment with Sm-X significantly reduced the number of apoptotic cells. A time-dependent decrease in Bax protein level and an increase in Bcl-2 protein level were observed in BDL rats treated with Sm-X. Immunohistochemical analysis demonstrated that p53 was strongly positive in hepatocyte nuclei of BDL rats but that treatment with Sm-X induced a cytoplasmic sequestration of p53. Taken together, hepatoprotective effects of Sm-X partially ascribe to the antiapoptotic property in hepatocytes. Treatment of Sm-X-induced cytoplasmic sequestration of p53, downregulation of Bax, and upregulation of Bcl-2 protein. This study identifies and delineates signaling factors involved in the antiapoptotic properties of Sm-X and suggests a potential application of *S. miltiorrhiza* in the clinical management of hepatic disease induced by toxic bile salts following biliary obstruction. © 2002 Elsevier Science (USA)

**Key Words:** *Salvia miltiorrhiza*; bile duct ligation; apoptosis; Bax; Bcl-2; p53; cytoplasmic sequestration.

*Salvia miltiorrhiza* BUNGE is a traditional Chinese herbal medicine used for centuries to improve blood circulation. *S. miltiorrhiza* has been widely used in oriental medicine for the

treatment of chronic hepatitis and liver fibrosis. In rats, components of a water-soluble extract of *S. miltiorrhiza* prevent peroxidative damage to liver microsomes, hepatocytes, and erythrocytes (Liu *et al.*, 1992). Lithospermate B, one of the active ingredients of *S. miltiorrhiza*, exerts a protective effect against cell death in experimental hepatitis induced by carbon tetrachloride and D-galactosamine/lipopolysaccharide (Hase *et al.*, 1997). Tanshinone I and tanshinone II-A, other components isolated from the root of *S. miltiorrhiza*, are effective antioxidants that inhibit lipid peroxidation (Cao *et al.*, 1996; Ng *et al.*, 2000). It has also been reported that, by inducing apoptosis of tumor cells and by preventing aflatoxin B1-induced hepatocarcinogenesis in rats, *S. miltiorrhiza* possesses chemopreventative potential (Liu *et al.*, 2000, 2001).

The failure of bile salt excretion in cholestasis leads to retention of hydrophobic bile salts within the hepatocytes (Greim *et al.*, 1972a,b) and causes apoptosis and/or necrosis (Myoshi *et al.*, 1999). Although the mechanism of bile salt-mediated apoptosis is not completely understood, the involvement of the Fas death receptor has been suggested by *in vitro* studies, as well as by *in vivo* studies using an animal model with bile duct ligation (BDL) (Myoshi *et al.*, 1999; Faubion *et al.*, 1999).

Following a death signal, mitochondrial cytochrome *c* is released into the cytoplasm (Gross *et al.*, 1999; Narita *et al.*, 1998), which activates caspase 9, and this subsequently activates caspase 3 in concert with the cytosolic factor Apaf-1 (Li *et al.*, 1997). In Fas-deficient *lpr* mice, hepatocyte apoptosis during extrahepatic cholestasis is associated with an increase in Bax expression and association of Bax with mitochondria (Myoshi *et al.*, 1999). The apoptosis of proliferating bile duct epithelial cells (BEC) induced by the Roux-en-Y biliodigestive anastomosis after BDL is related to *de novo* expression of Bax (Stähelin *et al.*, 1999).

Wild-type p53 is a tumor suppressor protein, which functions as a transcription factor downregulating cell proliferation and inducing apoptosis in response to various types of cellular stress (Levine, 1997; Ko and Prives, 1996). It has been suggested that one of the mechanisms of inhibition of apoptosis is related to the cytoplasmic sequestration of p53 (Kovacs *et al.*, 1996). This type of p53 is functionally inactive, as it fails to

<sup>1</sup> Current address: College of Pharmacy, Yanbian University, Yanji, Jilin 133000, People's Republic of China.

<sup>2</sup> To whom correspondence should be addressed.



OH ET AL.




FIG. 1. Serum biochemical parameters in Sm-X-treated BDL rats. Blood was obtained from sham control (□), BDL control (▨), and Sm-X-treated BDL rats (■) for the determination of aspartate aminotransferase (AST), alanine aminotransferase (ALT), and total bilirubin (TB) using an Autodry Chemistry Analyzer. Data are means  $\pm$  SD of five independent experiments. \* $p < 0.05$ , \*\* $p < 0.01$ , \*\*\* $p < 0.001$ , significantly different from BDL control rats.

bind to its specific DNA target. Thus, nuclear localization of p53 after translation in the cytoplasm is essential for its normal activity (Shaulsky *et al.*, 1991). We have reported previously that a hot water extract of *S. miltiorrhiza* (Sm-X) inhibits fibrosis and lipid peroxidation in BDL rats (Nan *et al.*, 2001). In the present study, we attempted to clarify the molecular mechanisms underlying the hepatoprotective effects of Sm-X in BDL rats. Of particular interest to this investigation was the expression and cellular localization of p53 and Bcl-2 family proteins in terms of the regulation of hepatocellular apoptosis *in vivo*.

## MATERIALS AND METHODS

**Preparation of Sm-X.** The roots of *S. miltiorrhiza* were purchased from Sungboe Farm Co. (Ahsung, Kyunggi-do, Korea) and authenticated by Professor Y.C. Kim, College of Pharmacy, Wonkwang University. Sm-X was prepared by boiling the dried roots with distilled water for 5 h. The extract was filtered, freeze-dried, and kept at 4°C. The extraction yield was approximately 8.1% (w/w). The dried extract was dissolved in distilled water before use.

**Biliary obstruction and animal treatments.** Specific pathogen-free male Sprague-Dawley rats (200–220 g) were obtained from Daehan Laboratory Animal Research Co. (Choongbuk, Korea) and allowed free access to standard chow and tap water. They were kept in temperature-controlled and filter-sterilized animal quarters under a 12-h/12-h light–dark cycle. After animals were anesthetized with pentobarbital sodium (50 mg/kg body wt ip), bile duct ligation was performed by double ligation of the common bile duct and sectioning between the ligatures. After surgery, Sm-X was dissolved in distilled water and administered daily (200 mg/kg po) over the experimental period. Control rats received equal volumes of the solvent and sham-operated animals were used as day 0 controls. Animals were euthanized at 3, 5, and 10 days after BDL.

**Antibodies.** Mouse monoclonal anti- $\alpha$ -smooth muscle actin ( $\alpha$ -SMA) antibody (clone 1A4) was purchased from Sigma Chemical Co. (Poole, UK) and mouse monoclonal antibodies against Bcl-2 and Bax were obtained from Santa Cruz Biotechnology (Santa Cruz, CA). Mouse monoclonal antibodies against p53 (clone PAb 240) and anti-mouse IgG were obtained from Neomarkers (Fremont, CA) and Zymed (San Francisco, CA), respectively.

**Measurement of serum biomarkers.** Blood was obtained by cardiac puncture for serum biochemical analysis. The activities of aspartate aminotransferase (AST) and alanine aminotransferase (ALT), as well as the content of total bilirubin (TB) in the serum, were measured using an Autodry Chemistry Analyzer (Spotchem SP4410, Arkray, Japan).

**In situ nick-end labeling for DNA fragmentation.** Apoptotic cells were detected by the method of terminal deoxynucleotidyl transferase-mediated deoxyuridine triphosphate nick-end labeling (TUNEL, fluorescein), using a commercial *in situ* death-detection kit (Boehringer, Mannheim, Germany). Formalin-fixed liver tissues were dehydrated through increasing concentrations of ethanol and then embedded in paraffin. Sections (4  $\mu$ m) were deparaffinized in xylene. After sections were rehydrated in phosphate-buffered saline, they were incubated with proteinase K (20  $\mu$ g/ml in Tris-HCl) for 10 min at room temperature and processed according to the manufacturer's protocol. The number of TUNEL-positive cells was counted in 20 random microscopic fields (200 $\times$ ) for each specimen using a confocal microscope (Fluoview, Olympus). Indirect TUNEL labeling assay was also performed using a TACS2 TDT-DAB *in situ* apoptosis detection kit (Trevigen, Gaithersburg, TN) to identify the cell types undergoing apoptosis. Color development was performed with 3,3'-diaminobenzidine-tetrahydrochloride and the sections were counterstained with methyl green.

**Immunoblots for  $\alpha$ -SMA, Bax, Bcl-2, and p53.** Isolated liver tissues were homogenized in RIPA buffer (10 mM Tris-HCl, pH 7.5, 150 mM NaCl, 1% Nonidet P-40, 0.1% sodium dodecyl sulfate, 1% sodium deoxycholate, 1 mM PMSF, and 5 mM EDTA). After the homogenates were centrifuged at 4°C (600g for 15 min), the supernatants were collected and the protein concentration was measured. Aliquots of cytosolic protein from each sample were separated onto a 12% acrylamide gel and transferred to a nitrocellulose membrane. Blots were probed with mouse monoclonal anti- $\alpha$ -SMA, anti-Bax, anti-Bcl-2, and anti-p53 (1:500 dilution) for 1 h. The membrane was incubated with secondary antibodies (1:1500 dilution) for 1 h and the blots were visualized by electrochemiluminescence using the Western Blotting Detection System (Amersham, UK).

**Immunohistochemical analysis of  $\alpha$ -SMA, Bax, Bcl-2, and p53.** Section preparation was initially the same as described for TUNEL assay. The sections were then subjected to antigen retrieval by microwave (700 W for 2 min) in citrate buffer (pH 6.0). Proteins were detected by avidin–biotin complex staining. Primary antibodies for  $\alpha$ -SMA (1:300), Bax (1:50), Bcl-2 (1:50), and p53 (1:50) were applied to sections overnight at 4°C. Alternate sections were treated with normal rabbit serum as a control. Sections were further treated with secondary antibody. Peroxidase activity was detected with 3-amino-9-ethylcarbazole before counterstaining sections with hematoxylin.

## RESULTS

### Effect of Sm-X on Liver Function Parameters in Serum

To test the general liver function of sham control, BDL control, and Sm-X-treated BDL rats, serum levels of AST, ALT, and TB were determined utilizing specific enzyme assays. As shown in Fig. 1, serum AST and ALT were increased

ANTIAPOPTOTIC PROPERTY OF *S. Miltiorrhiza* IN BILIARY OBSTRUCTION

at 3 days in BDL rats; the levels decreased thereafter. The TB level was increased and remained constant until 10 days after BDL. Treatment with Sm-X lowered the activities of AST and ALT significantly. The TB level was slightly increased by treatment with Sm-X at day 3, but remained unchanged thereafter.

#### Expression of $\alpha$ -SMA

Nonparenchymal hepatic stellate cells (HSC) are activated in response to liver damage. They produce type I collagen leading to hepatic fibrosis and also express the intracellular microfilament protein  $\alpha$ -SMA. Therefore, the effect of Sm-X on the expression of  $\alpha$ -SMA in BDL rats was determined by Western blot analysis and immunostaining of HSC. In BDL rats, expression of  $\alpha$ -SMA increased with time. However, treatment with Sm-X for 10 days did not affect the activation of HSC (Fig. 2).

#### Prevention of Apoptotic Cell Death by Sm-X

TUNEL staining was performed to analyze hepatocellular apoptosis using the direct fluorescein-labeling cell death detection kit. As shown in Fig. 3A–C, a five- to sevenfold increase in the number of hepatocytes undergoing apoptosis was observed at 3 and 5 days after BDL. Treatment with Sm-X significantly decreased the number of apoptotic cells in BDL rats. To identify the cell types undergoing apoptosis in BDL rats, we performed indirect TUNEL assay using the TACS2 TDT-DAB *in situ* apoptosis detection kit. We observed that only hepatocytes, and not HSC or BEC, were undergoing apoptosis in BDL rats (Fig. 3D). The number of apoptotic hepatocytes was reduced significantly by treatment with Sm-X (Fig. 3E).

#### Expression of Bax and Bcl-2 Proteins by Treatment with Sm-X

The expression of proapoptotic Bax and antiapoptotic Bcl-2 was monitored in BDL control and Sm-X-treated BDL rats. Following BDL, total cellular Bax protein initially increased at 3 and 5 days and then decreased at 10 days (Fig. 4A). Treatment of BDL rats with Sm-X further decreased the expression of Bax to the control level. Parallel to the results of Western blot analysis, the staining intensity of Bax increased in BDL rats and then decreased dramatically after treatment with Sm-X (Fig. 5A–C). Antiapoptotic Bcl-2 protein is known to be expressed in BDL rats as an adaptive mechanism to resist hepatocyte damage induced by toxic bile salts. As was anticipated, the level of Bcl-2 protein increased over time in BDL rats. This increase in Bcl-2 expression then became more pronounced in BDL rats treated with Sm-X (Fig. 4A). The increased ratio of Bax/Bcl-2 in BDL rats was reduced by treatment with Sm-X in a time-dependent manner (Fig. 4B). In BDL rats, most Bcl-2 was expressed in BEC surrounding bile ductules (Fig. 5E).

However, treatment with Sm-X increased Bcl-2 expression in hepatocytes rather than in BEC (Fig. 5F).

#### Cytoplasmic Sequestration of p53 by Sm-X

To test the relationship between Bax and p53 protein and to compare the time course of the apoptotic protein expression, we performed Western blot and immunohistochemical analysis of p53. It has been reported that Bax expression is regulated by p53 in many cell types. Consistent with upregulation of Bax expression in BDL rats, an increase in p53 protein was observed by Western blot analysis (Fig. 4A). In Sm-X-treated rats, however, the expression of p53 was not reduced in the Western blot analysis. This discrepancy between Bax and p53 protein expression in Sm-X-treated rats could be explained by the immunohistochemical analysis. At 3 and 5 days after BDL, strong intranuclear staining was observed, indicating that p53 had accumulated in the nucleus (Figs. 5H and 5I). By 10 days, when the apoptotic index had decreased to 50% of maximum, there was some sequestering of p53 in the cytoplasm and in the perinuclear region (data not shown). However, most p53 still remained in the nucleus.

In the Sm-X-treated BDL rats, we could observe that all of the p53 was sequestered in the cytoplasm (Fig. 5J). From these results, we can conclude that treatment of BDL rats with Sm-X protects hepatocytes from apoptosis by downregulating Bax and upregulating Bcl-2 expression. The time course of Bax expression correlates well with the relocation of p53 from the nucleus to the cytoplasm.

## DISCUSSION

The aim of the present investigation was to demonstrate the effect and to determine the possible molecular mechanisms of Sm-X on hepatocyte injury induced by bile salts in BDL rats. It has been demonstrated that the BDL rat model shows almost no generation of toxic intermediates, no massive hepatocyte necrosis, and no significant inflammation (Muriel *et al.*, 1994). BDL induces a type of liver damage that is etiologically and pathogenetically different from the experimental liver damage induced by poisons such as carbon tetrachloride (Kountouras *et al.*, 1984). The BDL model therefore allows the detection of antiapoptotic drug effects. Using the BDL model or toxic bile salts in primary hepatocyte culture, several researchers have examined the antifibrotic and hepatoprotective effects of apoptosis-modulating agents. Interferon- $\alpha$  and tetrandrine prevent hepatic fibrosis in BDL rats (Bueno *et al.*, 2000; Park *et al.*, 2000). Hydrophilic bile acids, such as ursodeoxycholic acid and tauroursodeoxycholic acid, reduce the incidence of apoptosis in hepatocytes induced by toxic bile salts (Benz *et al.*, 1998, 2000); antioxidants exert the same effect (Yerushalmi *et al.*, 2001).

The results of our present study demonstrate that the treatment of BDL rats with Sm-X prevents hepatocellular apoptosis

OH ET AL.




FIG. 2. Western blot and immunohistochemical analysis of  $\alpha$ -smooth muscle actin ( $\alpha$ -SMA) protein. Liver tissues were obtained from sham control, BDL control, and Sm-X-treated BDL rats. Seventy-microgram proteins were separated onto a 12% acrylamide gel, transferred to a nitrocellulose membrane, and probed with mouse monoclonal anti- $\alpha$ -SMA antibody (A). For immunohistochemical analysis, immunoreactivity was detected by avidin-biotin complex. Primary and secondary antibodies were applied to sections and peroxidase activity was detected with 3-amino-9-ethylcarbazole before counterstaining sections with hematoxylin. Activated stellate cells are seen around the blood vessels ( $\Rightarrow$ ) and proliferating bile ductules ( $\blacktriangleright$ ) of BDL control (B) and Sm-X-treated BDL rats (C). Data are representative of five independent experiments.

FIG. 3. Antiapoptotic effects of Sm-X in BDL rats. Apoptotic cells were detected using the direct fluorescein-labeling cell death detection kit (A, BDL control; B, BDL + Sm-X), and the number of apoptotic cells was counted in 20 random microscopic fields of sham control ( $\square$ ), BDL control ( $\blacksquare$ ), and Sm-X-treated BDL rats ( $\blacksquare$ ). The values are means  $\pm$  SD of five independent experiments ( $*p < 0.001$ ) (C). The cell types undergoing apoptosis were identified using a TACS2 TdT-DAB *in situ* apoptosis detection kit (D, BDL control; E, BDL + Sm-X). Only hepatocytes ( $\blacktriangleright$ ) and not hepatic stellate cells ( $\Leftarrow$ ) or bile duct epithelial cells ( $\Rightarrow$ ) were undergoing apoptosis in BDL rats. TUNEL stainings are representative of at least five independent experiments.

*in vivo*. Although the hepatoprotective and antifibrotic effects of *S. miltiorrhiza* or its constituents have been demonstrated in various cell types and animal models (Nan *et al.*, 2001; Cao *et al.*, 1996; Wang, 2000), the molecular mechanisms by which *S. miltiorrhiza* protects the hepatocyte from toxic bile salt-induced damage are largely unknown.

According to our previous results, the treatment of BDL rats with Sm-X for 4 weeks inhibits hepatic fibrosis and lipid peroxidation (Nan *et al.*, 2001). Sm-X contains a mixture of natural phenolic compounds, which have antioxidant effects *in vitro* and *in vivo* (Cao *et al.*, 1996; Lin *et al.*, 1996; Zhou *et al.*, 1999). It is widely accepted that the severity of hepatic damage

ANTIAPOPTOTIC PROPERTY OF *S. Miltoirrhiza* IN BILIARY OBSTRUCTION




FIG. 4. Western blot analysis of Bax, Bcl-2, and p53 proteins. Liver tissues were obtained from sham control, BDL control, and Sm-X-treated BDL rats. Total cellular protein was separated onto a 12% acrylamide gel, transferred to a nitrocellulose membrane, and immunoblotted with the corresponding antibodies. The blots were visualized by the electrochemiluminescence Western Blotting Detection System (A). The Bax/Bcl-2 ratio was obtained by densitometric analysis of each blot from sham control (□), BDL control (▣), and Sm-X-treated BDL rats (■) (B). Results are representative of at least five independent experiments.

FIG. 5. Immunohistochemical localization of Bax, Bcl-2, and p53 protein. Liver tissues were obtained from sham control (A, D, and G), BDL control (B, E, H, and I), and Sm-X-treated BDL rats (C, F, and J) and incubated with corresponding antibodies. Bax expression increased in hepatocytes (▶) and bile duct epithelial cells (⇨) of BDL liver tissues obtained at 5 days (B) but decreased dramatically after treatment with Sm-X (C). In BDL rats, Bcl-2 was expressed mainly in bile duct epithelial cells (⇨) at 5 days (E). The treatment of Sm-X increased the Bcl-2 expression in hepatocytes (▶) (F). p53 protein had accumulated in the nucleus in BDL control rats at 3 (H) and 5 days (I). Treatment of Sm-X to BDL rats induced cytoplasmic sequestration of p53 (J). (Inset: Higher magnification of the hepatocytes showing cytoplasmic sequestration of p53). Data are representative of at least five independent experiments.

in individuals or in experimental animals with biliary obstruction is causally associated with the extent of intrahepatic oxidative stress (Ljubuncic *et al.*, 2000; Baron and Muriel, 1999). Increased levels or accelerated generation of reactive oxygen species and toxic degradative products of lipid peroxidation induce hepatic cell death. Based on these findings, we have postulated that the antioxidative properties of Sm-X may contribute to the protection of the liver from damage induced by toxic bile salts in BDL rats.

Contrary to the results obtained in the 4-week-treatment study, short-term treatment of BDL rats with Sm-X for 10 days did not affect the expression of  $\alpha$ -SMA, a marker of hepatic stellate cell activation during the fibrogenic process. We have therefore studied the acute protective effects of Sm-X on hepatocyte apoptosis in BDL rats. Treatment with Sm-X for at least 3 days decreased the number of TUNEL-positive cells dramatically. After 10 days of treatment, the number of apoptotic cells had decreased to almost the control level.

Elevation of the Bax protein level is induced in many clinically relevant settings where cell death occurs. Bcl-2 is an apoptosis suppressing factor that heterodimerizes with Bax and neutralizes the protective effects of the latter. When Bcl-2 is present in excess, cells are protected from apoptosis. In contrast, when Bax is in excess and the homodimers of Bax dominate, cells are susceptible to programmed cell death. Therefore, it is the ratio of Bax to Bcl-2 that determines the fate of a cell, rather than the absolute concentrations of either (Gross *et al.*, 1999). In BDL rats, the Bax/Bcl-2 ratio was elevated at 3 and 5 days after BDL, but decreased at 10 days. The time course of changes in the Bax/Bcl-2 ratio correlates very well with the time course of the number of TUNEL-positive cells. According to Greim *et al.* (1972a), the total amount of rat liver bile acids is increased fourfold 3 days after BDL. The level of chenodeoxycholic acid, one of the major toxic bile acids responsible for hepatocyte apoptosis, increases threefold at 3 days after BDL and returns to almost normal at 8–10 days after BDL. Thus, the kinetic change in the concentration of bile acids is in good accord with our present data. Treatment with Sm-X reduced the expression of Bax and increased the expression of Bcl-2 significantly, thereby reducing the Bax/Bcl-2 ratio. Moreover, the expression of Bcl-2 was not restricted to BEC. Under treatment with Sm-X, Bcl-2 was also expressed in hepatocytes. This result indicates that, as well as BEC, with which bile salts are mainly in contact, hepatocytes are also protected from apoptosis by treatment with Sm-X.

Wild-type p53 protein plays a key suppressor role in cell growth and tumor formation and acts as a cell cycle checkpoint after DNA damage induces G1 arrest or apoptosis (Levine, 1997; Ko and Prives, 1996). One of the transcriptional targets of p53 that may be important for apoptosis is Bax. Overexpression of p53 increases Bax expression in several cell types and induces apoptosis (Selvakumaran *et al.*, 1994; Han *et al.*, 1996; Xiang *et al.*, 1998). In addition, Bcl-2, which can inhibit

apoptosis induced by enforced p53 expression, can physically associate with Bax, implying that this oncoprotein interferes with p53-dependent apoptosis by antagonizing Bax function (Oltvai *et al.*, 1993; Moll *et al.*, 1996). Wild-type p53 is a nuclear protein and nuclear localization is essential for its growth-suppressing activity. Cytoplasmic sequestration of wild-type p53 protein has been reported in mammalian models of viral oncogenesis and atherogenesis (Wang *et al.*, 2001) and in certain types of cancer (Moll *et al.*, 1995). Cytoplasmic p53 is functionally inactive because it fails to bind to its specific DNA target. Thus, cytoplasmic p53 protect cells from death, as has been shown in many tumors (Moll *et al.*, 1996; McKenzie *et al.*, 1999). In mammary epithelium and in a neuroblastoma cell line, treatment with retinoic acid or hormones reverses the cytoplasmic sequestration and functional repression of p53 (Kuperwasser *et al.*, 2000; Takada *et al.*, 2001), thereby suggesting that the functional status of p53 can be altered and regulated by specific stimuli. In our immunohistochemical analysis based on a p53-specific monoclonal antibody, reaction products in rats receiving Sm-X were observed only in the cytoplasm. Consistent with the cytoplasmic sequestration of p53, the expression of Bax decreased following treatment with Sm-X.

In cases of extrahepatic cholestasis, the use of *S. miltiorrhiza* may provide an effective prophylactic and therapeutic tool by inhibiting apoptosis in hepatocytes. The elucidation of the properties of the factors involved in the interactions of p53 in hepatocytes may uncover the mechanism of action of the antiapoptotic effects of Sm-X. The results presented in this report may open the window to develop a new therapeutic strategy in cholestatic liver injury.

#### ACKNOWLEDGMENT

This research was supported by a grant (PF002105-01) from the Plant Diversity Research Center of The 21<sup>st</sup> Century Frontier Research Program funded by the Ministry of Science and Technology of the Korean government.

#### REFERENCES

- Baron, V., and Muriel, P. (1999). Role of glutathione, lipid peroxidation and antioxidants on acute bile-duct obstruction in the rat. *Biochim. Biophys. Acta* **1472**, 173–180.
- Benz, C., Angermuller, S., Otto, G., Sauer, P., Stremmel, W., and Stiehl, A. (2000). Effect of tauroursodeoxycholic acid on bile acid-induced apoptosis in primary human hepatocytes. *Eur. J. Clin. Invest.* **30**, 203–209.
- Benz, C., Angermuller, S., Toxicol., U., Kloters-Plachky, P., Riedel, H. D., Sauer, P., and Stremmel, W. (1998). Effect of tauroursodeoxycholic acid on bile-acid-induced apoptosis and cytolysis in rat hepatocytes. *J. Hepatol.* **28**, 99–106.
- Bueno, M. R., Daneri, A., and Armendariz-Borunda, J. (2000). Cholestasis-induced fibrosis is reduced by interferon alpha-2a and is associated with elevated liver metalloprotease activity. *J. Hepatol.* **33**, 915–925.
- Cao, E. H., Liu, X. Q., Wang, J. J., and Xu, N. F. (1996). Effect of natural antioxidant tanshinone II-A on DNA damage by lipid peroxidation in liver cells. *Free Radical Biol. Med.* **20**, 801–806.

ANTIAPOPTOTIC PROPERTY OF *S. Miltiorrhiza* IN BILIARY OBSTRUCTION

- Faubion, W. A., Guicciardi, M. E., Miyoshi, H., Bronk, S. F., Roberts, P. J., Svingen, P. A., Kaufmann, S. H., and Gores, G. J. (1999). Toxic bile salts induce rodent hepatocyte apoptosis via direct activation of Fas. *J. Clin. Invest.* 103, 137-145.
- Greim, H., Trülsch, D., Roboz, J., Dressler, K., Czygan, P., Hutterer, F., Schaffner, F., and Popper, H. (1972a). Mechanism of cholestasis. 5. Bile acids in normal rat livers and in those after bile duct ligation. *Gastroenterology* 63, 837-845.
- Greim, H., Trülsch, D., Czygan, P., Rudick, J., Hutterer, F., Schaffner, F., and Popper, H. (1972b). Mechanism of cholestasis. 6. Bile acids in human livers with or without biliary obstruction. *Gastroenterology* 63, 846-850.
- Gross, A., McDonnell, J. M., and Korsmeyer, S. (1999). Bcl-2 family members and the mitochondria in apoptosis. *Genes Dev.* 13, 1899-1911.
- Han, J., Sabbatini, P., Perez, D., Rao, L., Modha, D., and White, E. (1996). The E1B 19K protein blocks apoptosis by interacting with and inhibiting the p53-inducible and death-promoting Bax protein. *Genes Dev.* 10, 461-477.
- Hase, K., Hasimu, R., Basnet, P., Kadota, S., and Namba, T. (1997). Preventive effect of lithospermate B from *Salvia miltiorrhiza* on experimental hepatitis induced by carbon tetrachloride or D-galactosamine/lipopolysaccharide. *Plant Med.* 63, 22-26.
- Ko, L. J., and Prives, C. (1996). p53: Puzzle and paradigm. *Genes Dev.* 10, 1054-1072.
- Kountouras, J., Billing, B. H., and Scherer, P. J. (1984). Prolonged bile duct obstruction: A new experimental model for cirrhosis in the rat. *Br. J. Exp. Pathol.* 65, 305-311.
- Kovacs, A., Weber, M. L., Burns, L. J., Jacob, H. S., and Vercellotti, G. M. (1996). Cytoplasmic sequestration of p53 in cytomegalovirus-infected human endothelial cells. *Am. J. Pathol.* 149, 1531-1539.
- Kuperwasser, C., Pinks, J., Hurlbut, G. D., Naber, S. P., and Jerry, D. J. (2000). Cytoplasmic sequestration and functional repression of p53 in the mammary epithelium is reversed by hormonal treatment. *Cancer Res.* 60, 2723-2729.
- Levine, A. J. (1997). p53, the cellular gatekeeper for growth and division. *Cell* 88, 323-331.
- Li, P., Nijhawan, D., Budihardjo, I., Srinivasula, S. M., Ahmad, M., Alnemri, E. S., and Wang, X. (1997). Cytochrome c and dATP-dependent formation of Apaf-1/caspase-9 complex initiates an apoptotic protease cascade. *Cell* 91, 479-489.
- Lin, T. J., Zhang, K. J., and Liu, G. T. (1996). Effects of Salviaolic acid A on oxygen radicals released by rat neutrophils and on neutrophil function. *Biochem. Pharmacol.* 51, 1237-1241.
- Liu, G. T., Zhang, T. M., Wang, B. E., and Wang, Y. W. (1992). Protective action of seven natural phenolic compounds against peroxidative damage to biomembranes. *Biochem. Pharmacol.* 43, 147-152.
- Liu, J., Shen, H. M., and Ong, C. N. (2000). *Salvia miltiorrhiza* inhibits cell growth and induces apoptosis in human hepatoma HepG2 cells. *Cancer Lett.* 153, 85-93.
- Liu, Y., Yang, C. F., Wasser, S., Shen, H. M., Tan, C. E., and Ong, C. N. (2001). Protection of *Salvia miltiorrhiza* against aflatoxin B1-induced hepatocarcinogenesis in Fischer 344 rats dual mechanisms involved. *Life Sci.* 69, 309-326.
- Ljubuncic, P., Tanne, Z., and Bomzon, A. (2000). Evidence of a systemic phenomenon for oxidative stress in cholestatic liver disease. *Gut* 47, 710-716.
- McKenzie, P. P., Guichard, S. M., Middlemas, D. S., Ashmun, R. A., Danks, M. K., and Harris, L. C. (1999). Wild-type p53 can induce p21 and apoptosis in neuroblastoma cells but the DNA damage-induced G1 checkpoint function is attenuated. *Clin. Cancer Res.* 5, 4199-4207.
- Moll, U. M., LaQuaglia, M., Benard, J., and Riou, G. (1995). Wild-type p53 protein undergoes cytoplasmic sequestration in undifferentiated neuroblastomas but not in differentiated tumors. *Proc. Natl. Acad. Sci. USA* 92, 4407-4411.
- Moll, U. M., Ostermeyer, A. G., Haladay, R., Winkfield, B., Frazier, M., and Zambetti, G. (1996). Cytoplasmic sequestration of wild-type p53 protein impairs the G1 checkpoint after DNA damage. *Mol. Cell. Biol.* 16, 1126-1137.
- Muriel, P., Suarez, O. R., Gonzalez, P., and Zuniga, L. (1994). Protective effect of S-adenosyl-L-methionine on liver damage induced by biliary obstruction in rats: A histological, ultrastructural and biochemical approach. *J. Hepatol.* 21, 95-102.
- Myoshi, H., Rust, C., Roberts, P. J., Burgart, L. J., and Gores, G. J. (1999). Hepatocyte apoptosis after bile duct ligation in the mouse involves Fas. *Gastroenterology* 117, 669-677.
- Nan, J. X., Park, E. J., Kang, H. C., Park, P. H., Kim, J. Y., and Sohn, D. H. (2001). Anti-fibrotic effects of a hot water extract from *Salvia miltiorrhiza* roots on liver fibrosis induced by biliary obstruction in rats. *J. Pharm. Pharmacol.* 53, 197-204.
- Narita, M., Shimizu, S., Ito, T., Chittenden, T., Lutz, R. J., Matsuda, H., and Tsujimoto Y. (1998). Bax interacts with the permeability transition pore to induce permeability transition and cytochrome c release in isolated mitochondria. *Proc. Natl. Acad. Sci. USA* 95, 14681-14686.
- Ng, T. B., Liu, F., and Wang, Z. T. (2000). Antioxidative activity of natural products from plants. *Life Sci.* 66, 709-723.
- Oltvai, Z. N., Millman, C. L., and Korsmeyer, S. J. (1993). Bcl-2 heterodimerizes in vivo with a conserved homolog, bax, that accelerates programmed cell death. *Cell* 74, 609-619.
- Park, P. H., Nan, J. X., Park, E. J., Kang, H. C., Kim, J. Y., Ko, G. I., and Sohn, D. H. (2000). Effect of tetrandrine on experimental hepatic fibrosis induced by bile duct ligation and scission in rats. *Pharmacol. Toxicol.* 87, 261-268.
- Selvakumaran, M., Lin, H. K., Miyashita, T., Wang, H. G., Krajewski, S., Reed, J. C., Hoffmann, B., and Liebermann, D. (1994). Immediate early up-regulation of bax expression by p53 but not TGF beta 1: A paradigm for distinct apoptotic pathways. *Oncogene* 9, 1791-1798.
- Shaulsky, G., Goldfinger, N., Tosky, M. S., Levine, A. J., and Rotter, V. (1991). Nuclear localization is essential for the activity of p53 protein. *Oncogene* 6, 2055-2065.
- Stähelin, B. J., Marti, U., Zimmermann, H., and Reichen, J. (1999). The interaction of bcl-2 and bax regulates apoptosis in biliary epithelial cells of rats with obstructive jaundice. *Virchows Arch.* 434, 333-339.
- Takada, N., Isogai, E., Kawamoto, T., Nakanishi, H., Todo, S., and Nakagawara, A. (2001). Retinoic acid-induced apoptosis of the CHP134 neuroblastoma cell line is associated with nuclear accumulation of p53 and is rescued by the GDNF/Ret signal. *Med. Pediatr. Oncol.* 36, 122-126.
- Wang, B. E. (2000). Treatment of chronic liver disease with traditional Chinese medicine. *J. Gastroenterol. Hepatol.* 15(Suppl.), E67-E70.
- Wang, J., Belcher, J. D., Marker, P. H., Wilcken, D. E., Vercellotti, G. M., and Wang, X. L. (2001). Cytomegalovirus inhibits p53 nuclear localization signal function. *J. Mol. Med.* 78, 642-647.
- Xiang, H., Kinoshita, Y., Knudson, C. M., Korsmeyer, S. J., Schwartzkroin, P. A., and Morrison, R. S. (1998). Bax involvement in p53-mediated neuronal cell death. *J. Neurosci.* 15, 1363-1373.
- Yerushalmi, B., Dahl, R., Devereaux, M. W., Gumprich, E., and Sokol, R. J. (2001). Bile acid-induced rat hepatocyte apoptosis is inhibited by antioxidants and blockers of the mitochondrial permeability transition. *Hepatology* 33, 616-626.
- Zhou, G. Y., Zhao, B. L., Hou, J. W., Ma, G. E., and Xin, W. J. (1999). Protective effects of sodium tanshinone IIA sulphinate against adriamycin-induced lipid peroxidation in mice heart in vivo and in vitro. *Pharmacol. Res.* 40, 487-491.