

Poster PE-13

Assessment of Malignancy in Brain Tumors by 3T MR Spectroscopy

최보영¹, 전신수², 김범수³, 이재문³, 정성택⁴, 안창범⁵, 오창현⁶, 김선일⁷, 이형구¹, 서태석¹
가톨릭대학교 의과대학 의공학교실¹, 신경과학교실², 방사선과학교실³,
(주)메디너스⁴, 광운대 전기공학과⁵, 고려대 전자 및 정보공학부⁶, 한양대 의공학교실⁷

Purpose : To assess clinical proton MR spectroscopy (MRS) as a noninvasive method for evaluating tumor malignancy at 3T high field system.

Materials and methods : Using 3T MRI/MRS system, localized water-suppressed single-voxel technique in patients with brain tumors was employed to evaluate spectra with peaks of N-acetyl aspartate (NAA), choline-containing compounds (Cho), creatine/phosphocreatine (Cr) and lactate. On the basis of Cr, these peak areas were quantificated as a relative ratio.

Results : The variation of metabolites measurements of the designated region in 10 normal volunteers was less than 10%. Normal ranges of NAA/Cr and Cho/Cr ratios were 1.67018 and 1.16015, respectively. NAA/Cr ratio of all tumor tissues was significantly lower than that of the normal tissues ($p=0.005$). Cho/Cr ratio of high-grade gliomas was significantly higher than that of low-grade gliomas ($P=0.001$). Except 4 meningiomas, lactate signal was observed in all tumor cases.

Conclusion : The present study demonstrated that the neuronal degradation or loss was observed in all tumor tissues. Higher grade of brain tumors was correlated with higher Cho/Cr ratio, indicating a significant dependence of Cho levels on malignancy of gliomas. This results suggest that clinical proton MR spectroscopy could be useful to predict tumor malignancy.

Acknowledgement : This study was supported by a grant of the Mid and Long Term Nuclear R/D Plan Program, Ministry of Science and Technology, Republic of Korea.