

Poster PE-15

Feasibility of Proton Chemical Shift Imaging with a Stereotactic Headframe

백현만¹, 최보영¹, 손병철², 정성택³, 이형구¹, 서태석¹
가톨릭대학교 의과대학 의공학교실¹, 신경과학교실², (주)메디너스,³

Purpose : To prove feasibility of proton chemical shift imaging (1H CSI) during stereotactic procedure, authors performed 1H CSI in combination with a stereotactic headframe and selected targets according to local metabolic information, evaluated the pathologic results

Materials and methods : The 1H CSI directed stereotactic biopsy was performed in five patients. 1H CSI was performed before conventional stereotactic MRI with gadolinium enhancement for stereotactic coordinates. The metabolite images expressed as integral ratios, Cho/Cr and Lac/Cr, were displayed in different colors. The stereotactic target coordinates were correlated with the coordinates from the 1H CSI images.

Results : The final pathologic results obtained were concordant with the local metabolic information from 1H CSI. We believe that 1H CSI-directed stereotactic biopsy has the potential to significantly improve the accuracy of stereotactic biopsy targeting

Conclusion : Metabolic signals derived from 1H CSI could give us more direct clues for stereotactic target selection during the subsequent conventional stereotactic MR imaging. 1H CSI was feasible with the stereotactic headframe in place. The final pathologic results obtained were concordant with the local metabolic information from 1H CSI.

Acknowledgement : This study was supported by a grant of the Center for Functional and Metabolic Imaging Technology, Ministry of Health & Welfare, Republic of Korea (02-PJ3-PG6-EV07-0002).