

**Motor Cortex in Hemiparetic Patients to Deep Intracerebral Hematoma**

백현만<sup>1</sup>, 최보영<sup>1</sup>, 손병철<sup>2</sup>, 정성택<sup>3</sup>, 이형구<sup>1</sup>, 서태석<sup>1</sup>

가톨릭대학교 의과대학 의공학교실<sup>1</sup>, 신경과학교실<sup>2</sup>, (주)메디너스,<sup>3</sup>

**Purpose :** To determine the motor cortex dysfunction in hemiparetic patients due to deep intracerebral hematoma, authors performed proton magnetic resonance spectroscopy (1H MRS) for the evaluation of biochemical changes in the cortex on affected hemisphere according to axonal injury at the level of internal capsule.

**Materials and methods :** Ten control subjects and 14 patients with documentable hemiparesis of varying severity hemiparesis were included. All the hemiparesis was caused by deep intracerebral hematoma (putaminal and thalamic hemorrhage). In vivo 1H MRS study was performed on a 3T MRI/MRS system using STEAM sequence. As a single-voxel technique, Spectral parameters were: 20 ms TE, 2000 ms TR, 128 averages, 2500 Hz spectral width, and 2048 data points.

**Results :** We found that the mean N-acetylaspartate (NAA)/ phosphocreatine (Cr) and NAA/ choline (Cho) ratios were significantly decreased in the motor cortex of the hemiparesis patients compared with control subjects.

**Conclusion :** 1H MRS examinations of the motor cortex might help to differentiate distinct clinical entities of hemiparesis and to monitor pharmacological effects in therapeutic trials, providing a quantitative biological marker for motor neuron dysfunction.

**Acknowledgement :** This study was supported by a grant of the Center for Functional and Metabolic Imaging Technology, Ministry of Health & Welfare, Republic of Korea (02-PJ3-PG6-EV07-0002).