

paper, we discuss the sample size calculation in 2 x 2 crossover design with the log-transformed data.

[OA-3] [ 04/18/2003 (Fri) 14:00 – 14:15 / Orchid ]

### **Protective Effect of Biopectin on 2,3,7,8-Tetrachlorodibenzo-p-dioxin Induced Reproductive System Damage and Its Action Mechanism**

Shim KyooJung<sup>o</sup> Choung SeYoung

Lab. of toxicology, College of Pharmacy, Kyung Hee University

A growing body of scientific research indicates that man-made chemicals (xenobiotics) may interfere with the normal functioning of endocrine, or hormone systems. These endocrine disruptors may cause a variety of problems with development, behavior, and reproduction. Amongst the xenobiotics the World Health Organization classed 2,3,7,8-TCDD as a "known" human carcinogen. Other than carcinogenicity, the dioxins exhibit immunotoxicological, reproductive and developmental effects in mammals.

According to many reports, tissue distribution of TCDD in male Sprague-Dawley rats (240-290 g) was considered complete in 24 hr after TCDD administration. The highest levels of distribution of the chemical were found in liver (5% of dose/g tissue), followed by white fat (1% of dose/g tissue); serum was lowest at 0.01% of dose/ml serum. TCDD was mainly excreted via feces and the biological half-life of TCDD was  $16.3 \pm 3.0$  days in rat.

Treatment of rats with TCDD resulted in a broad range of toxic effects including growth suppression, hepatomegaly, hypercholesterolemia, thymic atrophy, and increased microsomal enzyme activities.

The dioxin-induced toxic manifestations described above were reduced by water soluble biopectin administration. The hypocholesterolemic effects of pectins were studied in Sprague-Dawley rats and pectins significantly reduced plasma cholesterol levels: supplementation of pectin lowers LDL cholesterol and in some cases VLDL cholesterol rather than HDL cholesterol. The raised ratios of liver/body weight and elevated EROD activity in SD rats by TCDD were return to normal levels. Antiestrogenic effect of pectin was shown in E-screening method and trace of androgenic activity was observed in established A-screening test. Therefore, it was concluded that the lipophilic property of TCDD enhanced its adhesion to cholesterol and lowering characteristic of soluble biopectin discharge TCDD out of the body. In addition, trace of androgenic property of biopectin might help recovery of the TCDD induced male reproductive organ damage.

[OA-4] [ 04/18/2003 (Fri) 14:15 – 14:30 / Orchid ]

### **Bisphenol A-induced overall immune downregulation in mice.**

Byun JungA<sup>o</sup>, Pyo MyoungYun

College of Pharmacy, Sookmyung Women's University, Seoul

This study was undertaken to assess overall effects of bisphenol A, a monomer widely used in manufacturing polycarbonate plastics or epoxy resin, exposure on immune system of mice. For in vitro evaluation, serial concentration of BPA was added into culture of various immune cells from normal female ICR mice, and for in vivo or ex vivo assessment, mice were orally exposed to BPA dissolved in olive oil as doses of 500, 1000, 2000 mg/kg b.w. for acute exposure or 100, 500, 1000 mg/kg/day b.w. 5 days a week for subacute exposure. thereafter we conducted evaluation