

[PA4-41] [ 04/17/2003 (Thr) 14:00 – 17:00 / Hall P ]

### Acute effects of 2-bromopropane and 1,2-dibromopropane on hepatotoxic and immunotoxic parameters

Kim NamHee<sup>o</sup>, Hyun SunHee<sup>\*\*</sup>, Kim ChunHwa, Lee SangKyu, Lee DongWook, Jeon TaeWon, Lee JaeSung<sup>\*\*</sup>, Lee EungSeok, Chae WhiGun<sup>\*</sup>, Jeong TaeCheon

College of Pharmacy, Yeungnam University, Kyungsan, 712-749, Korea; <sup>\*</sup>School of Medicine, Taegu Catholic University, Taegu, Korea; <sup>\*\*</sup>Department of Food Science and Technology, College of Natural Resource, Yeungnam University, Kyungsan, 712-749, Korea

2-Bromopropane (2-BP) is a major component of the mixture of SPG-6AR and Solvent 5200 that is a substitute of chlorofluorocarbon. Many female workers exposed to 2-bromopropane in a Korean electronic company were found to have amenorrhea and male workers were diagnosed with oligospermia. In the present studies, immunotoxic effects of 2-BP and an analog, 1,2-dibromopropane (1,2-DBP), were investigated in female BALB/c mice. The mice were treated po with either 2-BP at 2000 and 4000 mg/kg or 1,2-DBP at 300 and 600 mg/kg once. The mice were immunized ip with sheep red blood cells (SRBCs). The spleen and thymus weights were reduced by 1,2-DBP. The number of splenic cells was decreased by 1,2-DBP. In addition, the antibody response to SRBCs was suppressed by the treatment with either 2-BP or 1,2-DBP. Meanwhile, these parameters were not significantly changed by the treatment with much lower doses of 2-BP. In a subsequent study, the time course effects of 2-BP and 1,2-DBP on the hepatotoxic and immunotoxic parameters were compared in mice. When mice were treated po with either one of these chemicals once for 6, 12, 24 and 48 hr, the activities of serum alanine aminotransferase and aspartate aminotransferase were significantly elevated only by 1,2-DBP 24 hr after the treatment. The hepatic content of glutathione was reduced by 1,2-DBP. The present results suggest that 1,2-DBP contained in the Solvent 5200 may contribute to the immunotoxicity, although 2-BP is a major component. (Supported by grant from Korea Science and Engineering Foundation (grant No. RO1-2000-00182)).

[PA4-42] [ 04/17/2003 (Thr) 14:00 – 17:00 / Hall P ]

### Cytochrome P450 발현 및 기능 조절제의 탐색 및 응용에 관한 연구

Kim NamHee<sup>o</sup>, Jeon TaeWon, Hyun SunHee, Kim ChunHwa, Lee SangKyu, Lee DongWook, Jeong TaeCheon

College of Pharmacy, Yeungnam University, Kyungsan, 712-749, Korea

To develop cytochrome P450 (P450) modulators with low toxicity from natural products, we have evaluated more than 20 compounds in mouse and rat models. Among those, the effects of 1,8-cineole and diallyl sulfide (DAS) were most profound in modulating P450 expression. In this presentation, the effects of 1,8-cineole and diallyl sulfide (DAS) on P450 expression were introduced. In addition, these two compounds were applied to investigate the role of metabolic activation in thioacetamide-induced hepatotoxicity and/or immunotoxicity in animal models. When rats were treated orally with 200, 400 and 800 mg/kg of 1,8-cineole for 3 consecutive days, the liver microsomal activities of pentoxoresorufin- and benzyloxyresorufin-O-dealkylases and erythromycin N-demethylase were dose-dependently induced. The Western immunoblotting analyses clearly indicated the induction of P450 2B1/2 and P450 3A1/2 proteins by 1,8-cineole. When rats were pre-treated orally with 800 mg/kg of 1,8-cineole for 3 days, thioacetamide-induced hepatotoxicity and immunotoxicity were significantly potentiated. When rats were treated orally with 100, 200 and 400 mg/kg of DAS in corn oil for three consecutive days, the liver