

In this study, effect of CAD on heterologous passive cutaneous anaphylaxis (PCA) were studied by the method of Levine and Vaz(1970). Anti-serum was prepared from OA-sensitized male Balb/c mouse at two weeks after the last challenge of ovalbumin (OA) and alumina gel, and these serum diluted with HEPES buffer by means of the heterologous PCA titer, i.e. the highest dilution inducing PCA in rats. Heterologous PCA test in rats were carried out to determine the width and contents of pigment leaked in the dorsal skin 30 minutes after i.v. injection of 0.2 ml of 1 % egg albumin and 1 % Evans blue mixture (1 : 1). It shows that all of CAD inhibited the heterologous PCA : Quinic acid, ferulic acid and coumaric acid at a dose of 25 mg/kg, and sinapinic acid and cinnamic acid at a dose of 50 mg/kg., inhibited significantly the contents of leaked pigments and width of pigmented skin as compared with control, respectively ($p < 0.01$). These activity were less than that of prednisolone acetate at a dose of 10 mg/kg. Quinic acid has more inhibitory activity of PCA than chlorogenic acid. These results indicated that CAD inhibited dose-dependently the anaphylactic hypersensitivity, and also the more hydrogen radical in benzene ring of CAD have, the more inhibitory activity on PCA have.

[PB2-7] [04/18/2003 (Fri) 09:30 - 12:30 / Hall P]

Inhibitory Action of Cinnamic Acid Derivatives on Reversed Cutaneous Anaphylaxis and Hemolysin Titer

Kim TaeDoo^o, Lee JiYun, Kim YounJoung, Lee JinHee, Sim SangSoo, Kim ChangJong

Department of Pathophysiology, College of Pharmacy, Chung Ang University

Cinnamic acid derivatives(CAD) originating from vegetable kingdom have some biological activity. Effect of CAD on reversed cutaneous anaphylaxis (RCA) and hemolysin (HY) titer were studied in rats. Experiments were carried out to determine RCA as the skin edema induced at 2 hours after injection of 0.05ml/site of anti-rat serum rabbit serum. Drugs were orally administered at one hour before antigen challenge. HY titer determined the hemolysis of sRBC to spleen cells. Two weeks after sensitization of i.p. injection of sRBC (4×10^8) cells, mice were challenged by i.p. injection of sRBC. On day 5 after sRBC challenged, spleen cells were isolated by cytosieve. HY titer exhibited as $\log_2 X$ (X is the highest dilution). Drugs were orally administered one hour before the last challenge of antigen. It shows that all of CAD have generally the dose-dependently inhibitory action on RCA and HY titer. Sinapinic acid at a dose of 12.5 mg/kg, and coumaric acid, quinic acid, chlorogenic acid, cinnamic acid and ferulic acid at a dose of 50 mg/kg inhibited significantly the RCA as compared with control, respectively ($p < 0.01$). Quinic acid at a dose of 12.5 mg/kg inhibited significantly HY titer as compared with control ($p < 0.01$), and its activity was one half of prednisolone acetate. Sinapinic acid and chlorogenic acid at a dose of 25 mg/kg have the significant inhibition of HY tier. These results indicated that the more methoxy or hydroxyl radical in benzene ring of CAD have, the more inhibitory activity of RCA and HY titer have.

Poster Presentations - Field B3. Neuroscience

[PB3-1] [04/18/2003 (Fri) 09:30 - 12:30 / Hall P]

ROLE OF ERK1/2 IN 6-HYDROXYDOPAMINE-INDUCED APOPTOSIS IN SK-N-SH HUMAN NEUROBLASTOMA CELLS