

bacteria. However, the purification of the glycosidases related to the metabolism of ginsenosides except β -glucosidase from K-110 have not been studied. We purified and characterized ginsenoside hydrolyzing β -glucosidase and α -arabinofuranosidase from *Bifidobacterium cholerium* K-103, a human intestinal bacterium. The specific activity of the homogeneously purified β -glucosidase was 0.65 $\mu\text{mole}/\text{min}/\text{mg}$. Molecular weight of the purified β -glucosidase was determined to be 360 kDa and the enzyme was composed of four identical subunits. Optimal pH range of β -glucosidase was 5.0~5.5 in phosphate buffer and the enzyme activity was inhibited by Cu^{++} . This enzyme transformed ginsenoside Rc to compound O. Specific activity of the homogeneously purified α -arabinofuranosidase was 0.76 $\mu\text{mole}/\text{min}/\text{mg}$. Molecular weight of the purified enzyme was determined to be 173 kDa and it has two identical subunits. Optimal pH range 5.5~6.0 in phosphate buffer and the enzyme activity was inhibited by addition of Cu^{++} . α -L-Arabinofuranosidase from *Bifidobacterium* K-103 transformed ginsenoside Rc to Rd but it didn't hydrolyze ginsenoside Rb2.

[PC2-3] [04/18/2003 (Fri) 09:30 - 12:30 / Hall P]

Antiallergic Activities of Daidzein, a Metabolite of Puerarin and Daidzin Produced by Human Intestinal Microflora

Park EunKyung^o, Choo MinKyung, Kim DongHyun

College of Pharmacy, Kyung Hee University

To evaluate the antiallergic activities of puerarin and daidzin from the rhizome of *Pueraria lobata*, *in vitro* and *in vivo* inhibitory activities of these compounds and their metabolite daidzein were measured. Daidzein exhibited potent inhibitory activity on the β -hexosaminidase release induced by DNP-HSA and potently inhibited the PCA reaction in mice. Daidzein administered intraperitoneally showed the strongest inhibitory activity and significantly inhibited the PCA reaction at doses of 25 and 50 mg/kg with inhibitory activity of 37% and 73%, respectively. The inhibitory activity of intraperitoneally administered daidzein was stronger than those of intraperitoneally and orally administered puerarin and daidzin. Therefore we believe that puerarin and daidzin in the rhizome of *Pueraria lobata* are prodrugs, which have antiallergic activities, produced by intestinal microflora.

[PC2-4] [04/18/2003 (Fri) 09:30 - 12:30 / Hall P]

Protective Effect of Kakkalide from *Puerariae Flos* on Ethanol-Induced Lethality and Hepatic Injury Is Expressed by Human Intestinal Microflora

Han YeoOk, Han MyungJoo, Park SungHwan^o, Kim DongHyun

Department of Food Science and College of Pharmacy, Kyung Hee University

The inhibitory effect of kakkalide isolated from *Puerariae flos* on ethanol-induced lethality and hepatic injury were investigated. Intraperitoneally treated Kakkalide was weakly reduced the mortality associated with administration of ethanol and did not reduce alcohol hepatotoxicity. However, orally administered kakkalide and intraperitoneally administered irisolidone significantly reduced the mortality and potently reduced serum ALT and AST activities on liver-injured mice by ethanol. When kakkalide (200mg/kg) was orally administered to rat, a main compound was irisolidone and kakkalide was not detected in blood and urine. Kakkalide was metabolized to irisolidone via kakkalidone when incubated with human intestinal microflora. Based on these

findings, it is suggested that kakkalide is a prodrug, and irisolidone is a bioactive agent for counteracting the effect of drinking and hepatic injury.

[PC2-5] [04/18/2003 (Fri) 09:30 – 12:30 / Hall P]

Diarylheptanoid Isolated from *Alpinia officinarum* Inhibits Pancreatic Lipase

Shin Ji-Eun^o, Song Myoung-Chong, Han MyungJoo, Baek NamIn, Kim Dong-Hyun

Department of Food Science, and College of Pharmacy, Kyung Hee University

Pancreatic lipase-inhibitory activity of the rhizome of *Alpinia officinarum* (AO) and its antihyperlipidemic activity were measured. When water extract of AO was stepwise fractionated with organic solvents, ethylacetate fraction exhibited the most potent inhibition. From it, Diarylheptanoid was isolated as an inhibitor of pancreatic lipase and we investigated its *in vitro* inhibitory effect of lipase activity and *in vivo* antihyperlipidemic effect. AO, ethylacetate fraction and Diarylheptanoid significantly inhibited lipase activity, and serum TG level in corn oil feeding-induced mice and serum TG and cholesterol in Triton WR-1339-induced hyperlipidemic mice.

[PC2-6] [04/18/2003 (Fri) 09:30 – 12:30 / Hall P]

Optimal culture conditions for production of Escherichia coli Adhesin protein coupled to Escherichia coli Heat Labile Enterotoxin A2B in Escherichia coli TB1.

Lee YongHwa Rhee DongKwon Pyo SuhkNeung

College of pharmacy, Sungkyunkwan University, Suwon, 440-746, Kyunggi-do, Korea

The FimH subunit of type 1-fimbriated *Escherichia coli* has been determined as a major cause of urinary tract infection. To produce a possible vaccine antigen against urinary tract infection, the fimH gene was genetically linked to the ltxa2b gene, which was then cloned into the pMAL-p2E expression vector. The chimaeric construction of pMALfimH/ltxa2b was transformed into *Escherichia coli* TB1 and its N-terminal amino acid sequence was analyzed. Fusion protein, the adhesin fused to the *Escherichia coli* heat labile enterotoxin A2B (LTXA2B), was induced for 4 hr with 0.3 mM isopropyl- β -D-thiogalactopyranoside (IPTG) at 37C, to yield soluble fusion protein. The expressed fusion protein was confirmed by SDS-PAGE, western blotting, and GM1-ganglioside ELISA using antibodies for maltose binding protein (MBP) and cholera toxin subunit B(CTXB). The results indicate that the purified fusion protein is an Adhesin/LTXA2B protein containing GM1-ganglioside binding activity of LTXB. The Adhesin/LTXA2B protein may be used as a candidate antigen for oral immunization against uropathogenic *E.coli*.

[PC2-7] [04/18/2003 (Fri) 09:30 – 12:30 / Hall P]

Inhibitory effects of Kimchi lactic acid bacteria on harmful enzymes of human intestinal bacteria

Han SeungBae, Kim DongHyun

Barumin Co. Ltd., and College of Pharmacy, Kyung Hee University