

microsomes.

[PD4-29] [04/18/2003 (Fri) 13:30 - 16:30 / Hall P]

Alternation of plasma F₂-isoprostanes before and after hemodialysis in patients with atherosclerosis in end-stage renal disease

Kim KyungMee^o, Jung ByungHwa, Kim SangWook, Paeng KiJung, Chung BongChul

Korea Institution of Science and Technology, Yonsei University, Korea University

F₂-isoprostanes are derived *in vivo* principally from the formation of positional peroxy radicals of arachidonic acid, endocyclization to prostaglandin G₂-like structures, and reduction to PGF₂-like compounds. F₂-isoprostanes have been proposed as biomarkers of lipid peroxidation, oxidative stress status, and oxidation of low density lipoprotein (LDL) in atherogenesis. Especially, elevated 8-iso-PGF_{2a} is known as a useful clinical biomarker of oxidative stress in end-stage renal disease (ESRD) patients receiving hemodialysis (HD). Therefore, we studied about alternation of plasma F₂-isoprostanes on before and after hemodialysis in patients with atherosclerosis in ESRD using gas chromatography-tandem mass spectrometry (GC/MS/MS). Plasma concentrations F₂-isoprostanes were significantly higher in post-hemodialysis than pre-hemodialysis in atherosclerosis ($P < 0.05$). This may suggest that F₂-isoprostanes is affected with oxidative stress by hemolysis treatment in atherosclerosis.

[PD4-30] [04/18/2003 (Fri) 13:30 - 16:30 / Hall P]

Methyl carboxylates of naphtho[1,2-b]furan and naphtho[1,2-b]pyran from the roots of *Rubia akane*

Jung SoonJa^o, Jung JiHyun, Cao LiHua, Shin YoungHee, Son JongKeun*, Lee ChongSoon**, Woo MiHee

College of Pharmacy, Catholic University of Daegu, *College of Pharmacy, Yeungnam University, **Department of Biochemistry, Yeungnam University.

Rubia akane (Rubiaceae) is a perennial vine plant and generally grows at the mountain, field and fence of Korea, Japan, China and Taiwan. This plant grows about two meters and has empty stem and red roots. The roots of this plant have been used as a Korean traditional medicine for the treatment of cough, the stone in the bladder and kidney, inflammation of the joints, uterine hemorrhage and uteritis.

From the screening experiment, *Rubia akane* exhibited anti-cancer activity. MeOH extract of the roots of this plant was subsequently fractionated into four parts: methylene chloride, ethyl acetate, n-butanol and water fractions. Methylene chloride fraction showed cytotoxicity against HT-29 and MCF-7 cell lines, and DNA Topoisomerase I inhibitory activity at the concentration of 5 $\mu\text{g}/\text{ml}$ and DNA Topoisomerase II inhibitory activity at the concentration of 50 $\mu\text{g}/\text{ml}$. Ethyl acetate fraction showed DNA Topoisomerase I inhibitory activity at the concentration of 5 $\mu\text{g}/\text{ml}$ and DNA Topoisomerase II inhibitory activity at the concentration of 50 $\mu\text{g}/\text{ml}$.

We isolated methyl carboxylates of naphtho[1,2-b]furan and naphtho[1,2-b]pyran from the methylene chloride fraction.