

macrophages Raw 264.7. The compound inhibited not only LPS-induced NO production with an IC<sub>50</sub> value of 6.5  $\mu$ M but also TNF and IL-1 productions with IC<sub>50</sub> values of 17.2  $\mu$ M and 10.0  $\mu$ M, respectively. Benzoxathiol LYR-71 compound inhibited the IL-6 signaling with an IC<sub>50</sub> value of 4.3  $\mu$ M but did not inhibit TNF or IL-1 signaling at all. Therefore, benzoxathiol LYR-71 compound would be beneficial for treatment of inflammatory and immune diseases.

[PC1-15] [ 2003-10-10 09:00 - 13:00 / Grand Ballroom Pre-function ]

### **Anti-inflammatory and Antinociceptive Effects of Methanol Extract from the Fomes fomentarius**

**Young-Mi Park**<sup>o</sup>, Kyoung-Min Shin, In-Tae Kim, Hee-Juhn Park, Jong-Won Choi, Kyoung-Tae Lee  
*College of Pharmacy, Kyung Hee University, Sangji University, Kyungsoong University*

As an attempt to search for bioactive natural products exerting antiinflammatory activity, we have evaluated the anti-inflammatory and antinociceptive activities of the methanol extract from the Fomes fomentarius (MEFF). MEFF (50, 100 mg/kg/day, p.o.) significantly reduced an acute paw edema induced by carrageenan in rats. When analgesic activity was measured by acetic acid-induced writhing test and hot plate test, MEFF showed a dose-dependent inhibition in animal models. In addition, MEFF potently inhibited the LPS-induced production of NO, PGE<sub>2</sub> and TNF- $\alpha$  production of macrophages. Consistent with these observations, the expression level of iNOS and COX-2 enzyme was decreased by MEFF in a concentration-dependent manner. These results suggest that methanol extract from the Fomes fomentarius exert anti-inflammatory effects by inhibiting NO and PGE<sub>2</sub> induction.

[PC1-16] [ 2003-10-10 09:00 - 13:00 / Grand Ballroom Pre-function ]

### **Fast visible dye staining of proteins in one- and two-dimensional sodium dodecyl sulfate-polyacrylamide gels compatible with matrix-assisted laser desorption/ionization-mass spectrometry**

Jin Li-Tai , Hwang Sun-Young , Yoo Gyurmg-Soo, Choi Jung-Kap  
*College of Pharmacy, Chonnam National University, Kwangju 500-757, South Korea, .*

A fast and matrix-assisted laser desorption/ionization-mass spectrometry compatible protein staining method in one- and two-dimensional sodium dodecyl sulfate-polyacrylamide gel electrophoresis is described. It is based on the counterion dye staining method that employs oppositely charged two dyes, Zincon and Ethyl Violet to form an ion-pair complex. It is safe to use since the methanol used previously in staining solution was replaced with ethanol, which is not toxic. The protocol including fixing, staining and quick washing steps can be completed in 1 to 1.5 h depending upon gel thickness. It has the sensitivity comparable with the colloidal Coomassie Brilliant Blue G (CBBG) stain using phosphoric acid as a component of staining solution (4-8 ng). The counterion dye stain does not induce protein modifications like in CBBG stain using trichloroacetic acid and methanol as components of staining solution, which complicates interpretation of peptide mapping data from mass spectrometry. Considering the safety, sensitivity, speed and compatibility with mass spectrometry, the counterion dye stain may be more practical than any other dye-based protein stains for routine proteomic researches.

[PC1-17] [ 2003-10-10 09:00 - 13:00 / Grand Ballroom Pre-function ]

### **Curcumin Inhibits Phorbol Ester-induced Expression of Cyclooxygenase-2 in Mouse Skin through Suppression of Extracellular Signal-Regulated Protein Kinase Activity and NF- $\kappa$ B Activation**

**Chun Kyung-Soo**<sup>o</sup>, Surh Young-Joon  
*College of pharmacy, Seoul National University*

Recently, there have been considerable efforts to search for naturally occurring substances for the intervention of carcinogenesis. Curcumin, a yellow coloring ingredient of turmeric (*Curcuma longa* L., Zingiberaceae), has been

shown to inhibit experimental carcinogenesis and mutagenesis, but molecular mechanisms underlying its chemopreventive activities remain unclear. In the present work, we found that curcumin inhibited 12-O-tetradecanoylphorbol-13-acetate (TPA)-induced expression of COX-2 in female ICR mouse skin when applied topically 30 min prior to TPA as determined by both immunoblot and immunohistochemical analyses. Multiple lines of evidence support the role of the eukaryotic transcription factor NF- $\kappa$ B in regulation of COX-2 expression. In agreement with this notion, the NF- $\kappa$ B inhibitor pyrrolidine dithiocarbamate suppressed not only NF- $\kappa$ B activation but also induction of COX-2 in mouse skin. Curcumin treatment attenuated TPA-stimulated epidermal NF- $\kappa$ B activation, which was associated with its blockade of degradation of the inhibitory protein I $\kappa$ B $\alpha$  and subsequent translocation of p65 subunit to nucleus. TPA treatment resulted in rapid activation via phosphorylation of extracellular signal-regulated protein kinase (ERK)1/2 and p38 mitogen-activated protein kinase (MAPK), which are upstream of NF- $\kappa$ B. The MEK1/2 inhibitor U0126 strongly inhibited NF- $\kappa$ B activation, while p38 MAPK inhibitor SB203580 failed to block TPA-induced NF- $\kappa$ B activation in mouse skin. Furthermore, U0126 blocked the TPA-induced I $\kappa$ B $\alpha$  phosphorylation by TPA, thereby blocking the nuclear translocation of NF- $\kappa$ B. Curcumin inhibited the catalytic activity of ERK1/2 in mouse skin. Taken together, suppression of COX-2 expression by inhibiting ERK activity and NF- $\kappa$ B activation may represent molecular mechanisms underlying previously reported anti-tumor promoting effects of this phytochemical in mouse skin tumorigenesis.

[PC1-18] [ 2003-10-10 09:00 - 13:00 / Grand Ballroom Pre-function ]

### **p38 MAP kinase and Akt regulate Bax translocation from mitochondria during ceramide-mediated apoptosis**

**Hae-Jong Kim**<sup>o</sup>, Seung-Koo Kang, Young-Jin Chun, Mic-Young Kim

*Department of Immunology, College of Pharmacy, Chung-Ang University, Seoul 156-756, Korea*

Ceramide is an important lipid messenger involved in mediating a variety of cell functions including apoptosis. Previously, we have shown that ceramide induces Bax translocation which is associated with cytochrome c release from the mitochondria. In this study, we show that p38 MAP kinase is involved in ceramide-induced Bax translocation. In human leukemic cells, ceramide stimulated the phosphorylation of p38 MAP kinase. Preincubation of cells with SB203580, a specific inhibitor of p38 inhibited DNA fragmentation induced by cell-permeable ceramide. Protection from apoptosis by SB203580 inhibited activation of caspase-3 and translocation of Bax. Furthermore, expression of dominant negative mutant of p38 attenuated ceramide-induced Bax translocation and apoptosis, indicating that p38 activation is required for Bax-mediated apoptosis induced by ceramide. In respect to PI3 kinase pathway, expression of constitutively active Akt reduced cell death and Bax translocation. We also found that expression of dominant-negative p38 suppressed ceramide-induced Akt dephosphorylation, indicating crosstalk between the two signaling pathways. Our results show that both the p38 and Akt pathways are involved in ceramide-mediated apoptosis by regulating Bax translocation.

[PC1-19] [ 2003-10-10 09:00 - 13:00 / Grand Ballroom Pre-function ]

### **Potent inhibition of human cytochrome P450 1 enzymes by SY-081**

**Kim Yongmo**<sup>o</sup>, Lee Sang Kwang, Kim Mie Young, Kim Sanghee, Chun Young Jin

*Department of Immunology, College of Pharmacy, Chung-Ang University, Seoul 156-756, Korea, Natural Products Research Institute, Seoul National University*

Recently we have reported that various hydroxystilbenes show strong inhibition of human cytochrome P450 1 enzyme activities. A series of synthetic trans-stilbene derivatives were prepared and their inhibitory potentials were evaluated with the bacterial membrane of recombinant human cytochrome P450 1A1, 1A2 and 1B1 coexpressed with human NADPH-P450 reductase to find a new inhibitor of cytochrome P450 enzymes. Of the compounds tested, SY-081 exhibited a potent inhibition of human cytochrome P450 1B1 with an IC<sub>50</sub> value of 2.6 nM. SY-081 also showed the inhibition of cytochrome P450 1A1 with IC<sub>50</sub> value of 47.6 nM and cytochrome P450 1A2 with IC<sub>50</sub> value of 116.6 nM. SY-081 showed 18-fold selectivity for cytochrome P450 1B1 over 1A1 and 45-fold selectivity for cytochrome P450 1B1 over 1A2. We also have investigated the inhibition kinetics of cytochrome