A0416

Calving Production from Hanwoo (Korean Cattle) IVM/IVF/IVC Blastocysts: Direct Transfer of Vitrified and Quick One-Step Diluted Hanwoo Blastocysts

<u>Sae Young Park</u>, Deok Im Kim, Jin Cheol Tae, Deok Im Kim¹, Sae Young Park, Eun Young Kim, Won Don Lee², Sepill Park and Jin Ho Lim²

Maria Infertility Hospital Medical Institute/Maria Biotech,

¹Hanwoo Improvement Center, NLCF, ²Maria Infertility Hospital

In this study we examined whether vitrified Hanwoo (Korean cattle) IVM/IVF/IVC blastocysts can survive *in vitro/in vivo* by a quick one-step dilution method and these embryos result in live births. Blastocysts produced *in vitro* were vitrified by serial exposure to glycerol (G) and/or ethylene glycol (EG) mixtures of 10% (v/v) G for 5 min, 10% G plus 20% EG (v/v) for 5 min, and 25% G plus 25% EG (v/v) for 30 sec. The blastocysts were then loaded in straw, placed in cold nitrogen vapor for 3 min and plunged into LN₂ (-196°C). One-step dilution within the straw was done in 25°C and/or 36°C water baths for different times (from 1 min to 3.5 min). *In vitro* survival of vitrified embryos was $76.9 \sim 93.5\%$ after the one-step dilution. However, 48 h after thawing, the *in vitro* development rates in the beyond hatching or hatched state in 1 min dilution group (the quick one-step dilution method; 73.3 and 56.7%, respectively) were better than those of the other treatment groups (55.6~61.3% and 8.1~36.1%, respectively). Direct transfer of quick one-step diluted embryos into recipient cows resulted in an overall pregnancy incidence of 33.3% (12/36). The good pregnancy incidence was obtained when the recipients estrus cycle was one day earlier than the age of the transferred embryos (53.3 vs. 25.0~27.3%), irrespective of synchronization methods or condition of the corpus luteum. We obtained nine offspring from 12 pregnant cows. Thus, quick one-step dilution and direct transfer of vitrified bovine IVM/IVF/IVC blastocysts could be applied as an efficient method for the field trials.

Key words: Hanwoo IVM/IVF/IVC blastocysts, Vitrification, One-step dilution, Direct transfer