

# OPTIMIZATION OF BIPHENILE CHLOROMETHYLATION PROCESS

V.V. Pak\*, R.K. Karimov\*, Kh.M. Shakhidoyatov\*, Deawha Soh\*\*

\*Institute of Plant Substance Chemistry, Uzbekistan Academy of Sciences, \*\*Myongji University  
dwhsoh@mju.ac.kr


**Abstract**— Optimization of the biphenile chloromethylation process with paraphormaldehyde has been done in the presence of ZnCl<sub>2</sub> with HCl gas by the Box-Wilson method of mathematical planning of experiment. The 4,4'-(dichloromethyl)-biphenile yield dependence on the biphenile – paraphormaldehyde ratio, temperature and reaction duration has been studied. A mathematical model of the process has been developed and optimal conditions for the biphenile chloromethylation procedure has been determined

## I. INTRODUCTION

The well known chlorometilation reaction is of great importance, since chlorine derivatives of aromatic compounds are intermediate products of synthesis of medicines, as well as chemicals protecting plants, dyes and others [1]. There is no any publication concerning procedure for producing 4,4'-(dichloromethyl)-biphenile. For this reason the biphenile chloromethylation reaction has been studied.

A 4,4'-(dichloromethyl)-biphenile is an intermediate component used in production of various classes of compounds which are in wide use in human life. Particularly, they are used in production of optical bleaches [2, 3] consumption of which in the world reaches to about 60-80 thousand tons per year [4]. It is also used in production of cleaning powder [5], thermoresistant and chemically resistant polymers [6]. Therefore, optimization and development of technological process to synthesize a 4,4'-(dichloromethyl)-biphenile is of practical interest.

In the papers [7] and [8] there were described procedures for production of 4,4'-(dichloromethyl)-biphenile in the presence of zinc chloride adding HCl gas in the reaction mixture (63% yield) and with aqueous HCl solution in the trichloroacetic acid (80% yield and 90% purity).



Our experiments on biphenile chloromethylation (I) with heating in concentrated HCl [9] have shown that

monochloromethylphenile (III) was produced with 90% yield. Add of HCl gas in the reaction mixture and in the presence of zinc chloride, the yield of the 4,4'-(dichloromethyl)-biphenile (II) was 90% with 95% purity. Therefore, to develop a technological scheme to produce a 4,4'-(dichloromethyl)-biphenile it was necessary to make optimization of the process.

## II. Results and Discussion

To optimize a synthesis of a 4,4'-(dichloromethyl)-biphenile we used the Box-Wilson method of mathematical modeling of experiments [10]. There were performed preliminary monofactor experiments to study a dependence of the 4,4'-(dichloromethyl)-biphenile yield on temperature, reaction duration and amount of waterless zinc chloride. A biphenile chloromethylation reaction has been performed at the constant molal ratio of the biphenile-paraphormaldehyde-ZnCl<sub>2</sub> which was 1.5:3.3:1, and different temperatures – 40, 45, 50, 55, 60, 65, 70 °C; duration of the reaction was 9 hours (Fig.1).




Fig. 1. Dependence of 4,4'-(dichloromethyl)-biphenile on the reaction temperature

At the temperature of 40 and 45°C the reaction was not completed fully, and residual monochloromethylbiphenile up to 6% presented in the reaction chemical mixture. At the temperature of 50°C, the reaction completed fully with a maximum yield of

90.3% and purity of 95%. Further temperature increasing resulted in decreasing the 4,4'-(dichloromethyl)-biphenile yield due to formation of by-products up to 15%. In this connection we have selected optimum temperature of 50°C and studied yield of dichloromethyl derivative at different amounts of zinc chloride. Duration of the reaction was 9 hours. The results of the experiments have been shown in Fig.2.




Fig.2. Dependence of 4,4'-(dichloromethyl)-biphenile on the amount of  $ZnCl_2$

From the Fig.2 it can be seen that increasing amount of waterless zinc chloride results in decreasing the reaction product yield. The experiments have also shown that when the portion of  $ZnCl_2$  was less than 50% of the initial amount the reaction failed.

The next set of experiments was aimed at the study of duration of the reaction process on the 4,4'-(dichloromethyl)-biphenile yield. Duration of the biphenile chloromethylation reaction depends on the amount of HCl-gas coming in during reaction process. In this connection we have determined the mean rates of HCl-gas entering at different reaction duration, namely, 9, 14 and 18 hours. The results are shown in Fig. 3.




Fig.3. Dependence of the mean rate of HCl entering on the reaction duration

It was revealed that increase of mean rate of HCl coming in the reaction mixture leads to decreasing duration of the reaction process. Dependence of the biphenile chloromethylation reaction product yield on the 1.5:3.3:1 molal ratio of the biphenile-paraphormaldehyde- $ZnCl_2$  at the 50°C is shown in Fig. 4 (a). There also are shown changes of 4-chloromethylbiphenile (b) and biphenile (c).




Fig.4. Dependence of 4,4'-(dichloromethyl)-biphenile on the reaction process duration

It can be seen from this figure that HCl gas coming in the reaction mixture longer than 9 hours does not increase the 4,4'-(dichloromethyl)-biphenile yield. A similar relation was observed at the mean HCl-gas rates during 14 and 18 hours reaction process, respectively.

On the base of these experiments, the main parameters having effect on the biphenile chloromethylation process can be considered  $x_1$  = temperature of the reaction (°C),  $x_2$  = duration of the reaction process and  $x_3$  = the waterless zinc chloride portion relative to its initial amount.

Table 1. Parameters and variation ranges

Parameters	Variation levels			Variation ranges
	-1	0	+1	
$x_1$	50	60	70	10
$x_2$	9	13.5	18	4.5
$x_3$	1	1.25	1.5	0.25

For optimization of the process a full factor experiment  $2^3$  with expanded planning matrix was used. Results of the experiments carried out are presented in Table 2.

Table 2. Planning matrix and the results of the experiments

No	$x_0$	$x_1$	$x_2$	$x_3$	$x_1x_2$	$x_1x_3$	$x_2x_3$	$x_1x_2x_3$	$y_1$	$y_2$	$y$	$S^2$	$y$	$(y-y)^2$
1	+	-	-	-	+	+	+	-	90.1	90.5	90.3	0.08	88.68	2.62
2	+	+	-	-	-	-	+	+	79.0	79.2	79.1	0.02	79.28	0.03
3	+	-	+	-	-	+	-	+	66.9	68.5	67.7	1.28	69.30	2.56
4	+	+	+	-	+	-	-	-	61.2	59.0	60.1	2.42	59.90	0.04
5	+	-	-	+	+	-	-	+	77.2	76.2	76.7	2.25	77.14	0.19
6	+	+	-	+	-	+	-	-	68.9	73.5	71.2	10.5	72.18	0.96
7	+	-	+	+	-	-	+	-	61.1	55.3	58.2	8.41	57.76	0.96
8	+	+	+	+	+	+	+	+	55.0	52.6	53.8	2.88	52.80	1.00

This equation was used for the program of sharp rising on the surface of response function in order to determine optimum conditions for the maximum 4,4'-(dichloromethyl)-biphenile yield. The variables in these experiments were temperature, reaction duration and amount of waterless zinc chloride. Intervals (or steps) of the variables were taken  $b_j \times \Delta x_j / 20$ :

$$b_1 \times \Delta x_1 = 3.59 \times 10 = 35.9$$

$$b_1 \times \Delta x_1 / 20 = 1.795 \cong 2^\circ\text{C}$$

$$b_2 \times \Delta x_2 = 9.69 \times 4 = 38.76$$

$$b_2 \times \Delta x_2 / 20 = 1.938 \cong 2 \text{ h}$$

$$b_3 \times \Delta x_3 = 4.66 \times 0.25 = 1.165$$

$$b_3 \times \Delta x_3 / 20 = 0.058 \cong 0.006\%$$

The results of the sharp rise experiments are given in the Table 3.

Table 3. Results of the sharp rise experiments

N	$x_1$	$x_2$	$x_3$	$y_1$	$y_2$	$y$
9	60	13.5	1.25	86.8	89.4	88.1
10	58	11.5	1.19	88.6	89.8	89.2
11	56	9.5	1.13	90.1	89.9	90.0
12	54	7.5	1.07	87.6	89.4	88.5

It can be seen from the table that the best result has been received in the experiment No1. Parameter variation in the sharp rise experiments did not improve the reaction yield. This can be explained by the fact that increasing the reaction duration leads to increasing mean HCl –gas rate coming in the reaction chemical mixture; a high HCl-gas rate leads to getting away the solvent, that leads to decreasing a solubility of the reagents at the initial stage of the reaction, and, therefore, leads to lowering the 4,4'-(dichloromethyl)-biphenile yield. On the other hand, increase of waterless zinc chloride leads to increasing intermediate layer that lessens the reaction product yield.

### III. Conclusion

The carried out experiments and use of the Box – Wilson mathematical model made it possible to make optimization of the biphenile chloromethylation process and determine optimal conditions for this process. Optimal values of the main parameters responsible for the maximum 4,4'-(dichloromethyl)-biphenile yield have been determined to be:

- mole ratio of the biphenile-paraphormaldehide-ZnCl<sub>2</sub> 1.5:3.3:1;
- temperature 50°C;
- duration of the reaction process 9 hours.

### REFERENCES

- [1] A.I. Belenkiy, Yu.B. Volkenshtein, N.B. Karmanov. New data on chloromethylation reaction for aromatic and heterocyclic compound. – Uspehi Himii, 1977, Vol. 46, No 9, P. 1699-1719 [Russian].
- [2] USA Patent, Cl. 260-240 CA, (C 07 D 263/56), No 3926964, Dec 16, 1975. Appl. July 30, 1973, No 383737.
- [3] Swiss Patent, Cl. C 07 B 27/00, C 08K 5/00, No 583668, Jan 14, 1977. Appl. Jun 30, 1972, No 9853/72.
- [4] A.G. Emelyanov. Optical bleaches and their application in textile industry. Moscow, 1971.
- [5] U.S. U.S. 5,326,491 (Cl. 252-95; C11D7/34), 05 Jul 1994, CH Appl 89/1,629, 28 Apr 1989; 9p.
- [6] B. Tarasyuk, W. Podkoscielny, W. Majewski. – Angew Mokromol. Chem. 1997, 251, 13-21 [Eng].
- [7] Swiss Patent, Cl. C 07C 15/12, C 07C 143/38, No 55481, Oct 15, 1974. Appl. Aug 20, 1968
- [8] Jpn. Kokai Tokkyo Koho Jp 1077, 238 [9877, 238] (Cl. C 07C22/04), Mar 24, Appl 96/233, 792, Sep 4 1996; 9 pp (Japan).
- [9] V.V. Pak, R.K. Karimov, Kh.M. Shakhidoyatov. Chemistry of natural compounds. Special issue. 1999. P. 101-102.
- [10] S.L. Akhazarov, V.V. Kafarov. Optimization of experiment in chemistry and chemical technology. Moscow: Visshaya Shkola. 1978. P. 158-178.