

***Heterorhabditis exomali* n. sp (Rhabdiyida:
Heterorhabditidae) from White Grub Cadavers**

Ho Yul Choo, Dong Woon Lee¹ and Chang Dae Song

Division of Applied Life Science, Gyeongsang National University,
Institute of Agriculture & Life Sciences, Gyeongsang National University

¹Institute of Agricultural Science, Sangju National University

Heterorhabditis exomali n. sp. was isolated from cadavers of oriental beetle (*Exomala orientalis*) larvae from Daegu Golf Club in Gyeongsangbukdo province in 1996. Adult female ovaries paired, reflexed, vulva located at midbody with slightly protruding lips, excretory pore not located on anterior to basal bulb., Hermaphroditic female length 3557.66 ± 322.10 , greatest body width 196.26 ± 20.92 , stoma length 16.31 ± 4.21 , stoma width 21.88 ± 2.99 , anterior end to nerving 138.50 ± 16.66 , oesophagus length 170.23 ± 12.86 , tail length 49.36 ± 9.89 , body width at anus 39.73 ± 8.65 , anterior end to vulva 1671.75 ± 187.67 . Amphimictic female length 1360.86 ± 368.09 , greatest body 81.22 ± 22.58 , stoma length 4.63 ± 1.03 , stoma width 6.76 ± 1.71 , anterior end to nerving 83.73 ± 19.17 , oesophagus 111.36 ± 22.10 , tail length 51.67 ± 8.95 , body width at anus 24.08 ± 4.42 , anterior end to vulva 666.43 ± 194.51 . Male length 888.39 ± 50.44 , greatest body 46.73 ± 2.95 , stoma length 4.12 ± 1.01 , stoma width 5.11 ± 0.94 , anterior end to nerving 80.63 ± 4.46 , oesophagus length 105.67 ± 10.94 , tail length 23.88 ± 2.37 , body width at anus 20.51 ± 1.32 , spicule length 42.51 ± 2.37 , spicule width 6.02 ± 0.79 , gubarnaculum length 21.80 ± 1.76 , gubarnaculum width 2.04 ± 0.33 . larvae length 560.62 ± 20.42 , greatest body width 21.12 ± 0.89 , stoma length 2.76 ± 0.48 , stoma width 1.80 ± 0.39 , anterior end to nerving 102.43 ± 8.49 , oesophagus length 115.76 ± 7.60 , tail length 55.30 ± 3.68 , body width at anus 12.52 ± 1.12 .