

[S-6]**Pulmonary Toxicity of Single Walled Carbon Nanotubes**

Vincent Castranova

National Institute for Occupational Safety and Health, Morgantown, West Virginia, USA

Single walled carbon nanotubes (SWCNT) are, in their purest form, a single layer of carbon atoms in a cylindrical arrangement. Individual SWCNT are 1.5 nm in diameter and up to a millimeter or more in length. SWCNT are of commercial interest because of their great tensile strength, high conductivity, high surface area, unique electronic properties and high adsorption capacity. However, little is known concerning the potential adverse effects of SWCNT in the lung. In the present study, SWCNT, produced by the HiPco process and purified to remove contaminating iron, were evaluated. Mice were exposed to 10, 20, or 40 µg of SWCNT by pharyngeal aspiration. Saline was used as the vehicle control. Pulmonary responses were evaluated 1-60 days post-exposure. Aspiration of SWCNT was mimiced by nebulization of this material. Nebulized SWCNT exhibited two distinct morphologies, i.e., aggregates and dispersed SWCNT, as determined by transmission electron microscopy. Aspiration of SWCNT resulted in dose dependent oxidant stress (depletion of GSH), pulmonary damage (protein and LDH in bronchoalveolar lavage fluid), and inflammation (neutrophils, IL-1 β , TNF- α , TNF- α and TGF- β 1 in bronchoalveolar lavage samples), which peaked 1-7 days post-exposure before declining toward baseline. Sites of deposition of aggregates of SWCNT developed granulomatous lesions within 7 days post-exposure with the development of fibrosis, which progressed through 60 days post-exposure. Alveolar regions distal to SWCNT aggregates developed interstitial fibrosis, which progressed from 7-60 days post-exposure. Gold labeling of SWCNT indicated that this interstitial fibrotic reaction was associated with the deposition of nanotubes or nanoropes rather than aggregates of SWCNT. These data indicate that acute pulmonary exposure of rats to SWCNT results in a dose-dependent fibrotic response, which develops rapidly and progresses with time post-exposure. Therefore, inhalation of SWCNT may be a concern.

Pulmonary Toxicity of Single Walled Carbon Nanotubes

Anna Shvedova, PhD
Robert Mercer, PhD
Andrew Maynard, PhD
Vincent Castranova, PhD

National Institute for Occupational Safety and Health
Morgantown, West Virginia, USA

Background

Single Walled Carbon Nanotubes (SWCNT)

- A. Long tube-like configuration of carbon molecules
- B. Single layer of carbon atoms in a cylindrical arrangement
- C. Nanotube \approx 1.5 nm in diameter, up to 1 mm in length
- D. High tensile strength, high surface area, unique electronic properties, high adsorption capacity
- E. Used in electronics, structural materials, etc

Issue

- A. Potentially wide commercial applications
- B. Little information is available concerning the potential adverse effects of inhalation of SWCNT

Objective







- A. Characterize the pulmonary responses to SWCNT
- B. Determine the dose-dependence of the responses
- C. Determine the duration of responses post-exposure







Test Material







- A. SWCNT were produced by the high-pressure carbon monoxide process (HiPCO)
- B. Unpurified SWCNT - contain 30% metal catalyst (ultrafine Fe, or Fe/Ni)
- C. Purified SWCNT - acid treated; < 0.2 wt% metals
- D. Suspended material supplied by NASA





Methods

- A. Exposure
 - 1) Pharyngeal aspiration of mice
 - 2) PBS or SWCNT (10, 20, or 40 μ g/mouse)
 - 3) Sacrifice 1 - 60 days post-exposure
- B. Endpoints
 - 1) Damage - BAL protein and LDH
 - 2) Inflammation - BAL cells and cytokines
 - 3) Oxidant stress - lung GSH
 - 4) Histology - particle deposition, inflammation, granulomas, and fibrosis









Summary

A. Nebulized SWCNT dispersed as aggregates and nanotubes

B. Aspiration causes transient oxidant stress, damage and inflammation, peaking by 7 days post-exposure

C. Histology visualizes aggregates in the terminal bronchiols and proximal alveoli with no visible material in distal alveoli

D. Size of aggregates doesn't change with time

E. Rapid fibrosis – begins in 7 days and progresses through 60 day post-exposure

- 1) Fibrosis in granulomatous lesions containing aggregates
- 2) Diffuse interstitial fibrosis in distal alveolar walls with no visible SWCNT

F. Used silver enhancement of gold-labeled SWCNT

- 1) See aggregates in proximal alveoli and terminal bronchiols
- 2) See nanoropes in walls of distal alveoli

Conclusions

A. See granulomatous lesions at deposition sites of aggregates.

B. See interstitial fibrosis in sites of deposition of nanoropes.