

# Two New Species and Two New Records of Cuculliinae (Lepidoptera, Noctuidae) from Korea

Hui Lin Han and Kyu Tek Park

Center for Insect systematics, Kangwon National University, Chuncheon, 200-701

The first record of the subfamily Cuculliinae from Korea was made by Leech (1889), reporting a species, *Cucullia perforata* Bremer, from Wonsan, N. Korea. Following after Draudt (1934), Wu (1938), Bryk (1948), Pak (1959), Park and Lee (1977), Nam (1982), Ronkay and Park (1993), Ahn *et al.* (1994, 1996) and some others, Kononenko *et al.* (1998) reviewed 85 species belonging to 31 genera, describing two new species of the subfamily, *Meganephria parki* and *Eupsilia silla*. In this study, two new species, *Lithophane kononenkoi* sp. n. and *Valeria gongjuensis* sp. n., are described, and two species, *Cucullia hostilis* Boursin and *Eupsilia boursini* Sugi, are reported for the first time from Korea. On the other hand, the genus *Valeria* Stephens, 1829, which is mainly distributed in S. China, Turkey, Ussuri region and Central Asia, is known for the first time in the Muchurian subregion.

**Key words:** Taxonomic, Lepidoptera, Noctuidae, fauna, Korea, Mt. Changbai