Isolation and characterization of marine microorganisms producing cellulase

<u>Kang-Ick Jo</u>, Bo-Hwa Lee, Hae-Young Jo and Jin-Woo Lee, Sung-Koo Kim¹, Soo-Wan Nam²

Department of Biotechnology, College of Nature Resources and Life Science, Dong-A University,
Busan

¹Department of Biotechnology and Bioengineering, Pukyung National University, Busan ²Department of Biotechnology and Bioengineering, Dong-Eui University, Busan TEL: +82-51-200-5749 FAX: +82-51-20-6536

Marine microorganisms from the seashore of the Kyungsang province in Korea to produce functional biopolymers were isolated. Seventy microorganisms isolated in this study were found to produce functional biopolymers. Some of them hydrolyzed cellulose, carboxymethyl cellulose(CMC) or skim milk. Microorganism to show the activity to hydrolyze cellulose and CMC were incubated in marin broth and the other liquid medium containing 2.0% (w/v) glucose, 0.25% yeast extract, 0.5% K₂HPO₄, 0.1% NaCl, 0.02% MgSO₄·7H₂O and 0.06% (NH₄)₂SO₄ at 30°C for 72 hr under aerobic conditions 200rpm. Cellulase and CMCase activities of supernatants were assayed. One microorganism cultivated in marine broth showed higher activity to hydrolyze cellulose than *Bacillus amyloliquefaciens* DL-3, which was known as a cellulase-producing strain, whereas six microorganisms cultivated in the other liquid medium showed higher activities of cellulase and CMCase than *B. amyloliquefaciens* DL-3. Optimization of media to produce cellulase by these strains isolated form seawater will be studied.

References

- 1. Chio, H. J., M. J. Jung, Y. K. Jeong (2000), Isolation and Identification of a Marine Bacterium Producing an Immunostimulant, J. Korean Soc Food Sci. Nutr. 29(6), 1145-1150.
- Kim, Y. C., S. S. Yoo, Y. A. Oh, S. J. Kim (2003), Isolation and Characteristics of Trichoderma harzianum FJ1 Producing Cellulase and Xylanase, J. Microbiol. Biotechnol. 13(1), 1-8.