

Study on Sansevieria removing of the generation of Benzene

Ji-Seok Oh, Hae-Jeong Moon, Chang-Ho Park

College of Environment and Applied Chemistry, Kyung Hee University, Yongin-si, Kyunggi-do
449-701, Korea

TEL: +82-31-201-2531, FAX: +82-31-202-1946

Abstract

This research investigates the efficiency of removing Benzene by plant. The crude protein extracted from plant decreases the Benzene's concentration partially. The obtained results show that proteins of various plant have each different degradation rate of Benzene and the protein of sansevieria is more favorable for removing activity than those of other plants. The important point of this view is "plant has an new important removal mechanism of Benzene"

References

1. Xuan YJ, Endo Y, Fujimoto K (2002), Oxidative degradation of bisphenol a by crude enzyme prepared from potato, *J Agric Food Chem.* 50, 6575-6578.
2. Converse BM, Schroeder ED, Iranpour R, Cox HH, Deshusses MA (2003), Odor and volatile organic compound removal from wastewater treatment plant headworks ventilation air using a biofilter, *Water Environ Res.* 75(5), 444-454.
3. Lu C, Chu W, Lin MR (2000), Removal of BTEX vapor from waste gases by a trickle bed biofilter, *J Air Waste Manag Assoc.* 50(3), 411-417.