

Characterization of an extracellular β -agarase from *Agarivorans* sp. JA-1

Jong-Geun Jung, Jin-Wook Kim, Chi-Un Joo, Song-Ae Lee,
Jae-Woong Hwang, and Jae-Hwa Lee*

Department of Bioscience and Biotechnology, Silla University, Busan 617-736, Korea
TEL: +82-2-999-5748, FAX: +82-2-999-5636

Abstract

Agar, a polysaccharide present in the walls of some red algae, can be degraded by several bacterial strains from marine environments and other sources. Some of the bacterial isolates have been assigned to the genera *Bacillus*, *Paenibacillus*, *Streptomyces*, *Pseudoalteromonas*, *Pseudomonas*, *Alteromonas*, *Microscilla*, *Vibrio*, *Cytophaga* and *Zobellia*. β -agarase is unique in its ability to break down the agarose polysaccharide core made up of repeating 1,3-linked β -D-galactopyranose and 1,4-linked 3,6-anhydro- α -L-galacto-pyranose into neoagarobiose oligosaccharides. In this study, we describe the properties of β -agarase from *Agarivorans* sp. JA-1. The strain was cultivated in 300 ml flask with shaking incubator at 30 °C, 250 rpm and the culture broth was centrifuged to obtain clear supernatant which was used for all assay. Temperature, pH, and substrate concentration were tested for investigating of the optimal β -agarase activity.

References

1. Ohta Y, Hatada Y, S Ito, K Horikoshi, High-level expression of a neoagarobiose-producing β -agarase gene from *Agarivorans* sp. JAMB-A11 in *Bacillus subtilis* and enzymatic properties of the recombinant enzyme. (2004), *J. Biotechnology. & Applied. Biochemistry*, **10**, 1042/BA20040083.
2. Yasushi S, N Hideki, I Koji, Y Takeshi, T Ichiro, Y Yoshinari, Production and characteristics of some new β -agarases from a marine bacterium, *Vibrio* sp. strain JT0107, (1995) *J. Fermentation and Bioengineering*, **79**, 549-554.
3. Hisashi S, S Yoshinori, S Tohru, K Keiich, Purification and characterization of an extracellular β -agarase from *Bacillus* sp. MK03, (2003) *J. Bioscience and Bioengineering*, **95**, 328-334.