

## Antioxidant responses of *Pseudomonas putida* BCNU 106 to toluene stress

Ju Soon Yoo,<sup>3</sup>Ja Young Moon, <sup>4</sup>Dong-Wan Kim,

<sup>5</sup>Young Kee Jeong, and <sup>2</sup>Woo Hong Joo

Institute of Genetic Engineering, <sup>2</sup>Department of Biology, <sup>3</sup>Department of  
Biochemistry and Health Sciences, <sup>4</sup>Department of Microbiology,  
Changwon National University, Changwon 641-773, Korea

<sup>5</sup>Department of Life science and Biotechnology, Dong-Eui University,  
Pusan 614-714, Korea

*Pseudomonas putida* BCNU 106 was used to examine the links between the toluene exposure, reactive oxygen species generation and oxidative stress response. Treatment of *Pseudomonas putida* BCNU 106 with 10% toluene revealed decreased generation of reactive oxygen species assessed by 2',7'-dichlorofluorescein oxidation. Antioxidative defence systems were investigated by measuring the activity of superoxide dismutase, glutathione peroxidase, glutathione reductase, glutathione S-transferase and catalase. Increased activity of superoxide dismutase, glutathione peroxidase, glutathione reductase, glutathione S-transferase and catalase were perceived after the treatment with 10% toluene. The results indicate that the tolerance of *Pseudomonas putida* BCNU 106 to different organic solvents was correlated with the reactive oxygen species generation in the cells and with the efficiency of antioxidant defence systems.

### Reference

1. W. jakubowski, T. bilinski, G. bartosz (2000) Oxidative stress during aging of stationary cultures of the yeast *Saccharomyces Cerevisiae*. *Free Radical Biology & Medicine*, 28(5), 659-664, 2000.