

In silico discovery and characterization of novel fish epoxide hydrolase from Zebrafish, *Danio rerio*

Soo Jung Lee¹, Eun Yeol Lee, Hee Sook Kim

Department of Food Science and Technology, Kyungshung University, Busan 608-736, Korea; ¹Recipient of BB21 scholarship

Enantiopure epoxides are important intermediates for producing optically active compounds. In this study, novel fish epoxide hydrolase (EH) from *Danio rerio* was accessed and compared based on the multiple sequence alignments, phylogenetic analysis and homology modelling. *D. rerio* EH was cloned and expressed in *Escherichia coli* for the first time. The recombinant *E. coli* expressing the enantioselective EH gene of *D. rerio* was used for producing enantiopure styrene oxide by asymmetric resolution of racemic styrene oxide. The recombinant *E. coli* exhibited the enantiopreference toward (*R*)-enantiomer, and has a maximum hydrolytic activity as 8.3 $\mu\text{mol}/\text{mg cell} \cdot \text{min}$. We obtained enantiopure (*S*)-styrene oxide with high optical purity ($> 99\%$ *ee*) and a yield of 31.3 % (theoretically 50 % maximum yield) at 60 min from 60 mM of racemic styrene oxide using the recombinant *E. coli*.

Acknowledgement : This work was supported by Marine and Genome Research Center Program, Ministry of Marine Affairs and Fisheries, Republic of Korea.